Originality Report

Similarity Found: 2%

Date: Wednesday, January 23, 2019
Statistics: 188 words Plagiarized / 7808 Total words
Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

Vol. 24 No. (4) 2018 also de v eloped b y scimago: SCIM A GO INSTI TUTIONS RANKINGS Scimago Journal & Countr y Rank Home Journal Rankings Countr y Rankings Viz T ools Help About Us E cology , Envir onment and Conser v ation Countr y India - SIR Ranking of India 11 H Index Subject Ar ea and Categor y Agricultur al and Biological Sciences E cology , E v olution, Beha vior and Systematics Envir onmental Science E cology Natur e and Landscape Conser v ation Publisher EM International Publication type Journals ISSN 0971765X Co v er age 1997-ongoing Scope ECOL OGY , ENVIRONMEN T AND CONSER V A TION is one of the leading International envir onmental journal.

It is widely subsribed in India and abr oad b y Institutions and Individuals in education and r esear ch as well as b y Industries, Go vt. Depar tments and Resear ch Institutes. Homepage Contact Join the conv ersation about this journal Enter Journal T itle, ISSN or Publisher Name Quar tiles The set of journals ha v e been r ank ed accor ding t o their SJR and divided int o four equal gr oups, four quar tiles.

Q1 (gr een) comprises the quar ter of the journals with the highest v alues, Q2 (y ellow) the second highest v alues, Q3 (or ange) the thir d highest v alues and Q4 (r ed) the lowest v alues. Categor y Y ear Quar tile E cology 1999 Q4 E cology 2000 Q3 E cology 2001 Q3 E cology 2002 Q3 SJR The SJR is a siz e-independent pr estige indicat or that r anks journals b y their ' a v er age pr estige per ar ticle '. It is based on the idea that ' all citations ar e not cr eated equal'.

SJR is a measur e of scienti c in uence of Citations per document This indicat or counts

the number of citations r eceiv ed b y documents fr om a journal and divides them b y the t otal number of documents published in that journal. The char t shows the e v olution of the a v er age number of 1999 2001 2003 2005 2007 2009 2011 2013 2015 2017 E cology E cology , E v olution, Beha vior and Systematics Natur e and Landscape Conser v ation journals that accounts for both the number of citations r eceiv ed b y a journal and the impor tance or pr estige of the journals wher e such citations come fr om It measur es the scienti c in uence of the a v er age ar ticle times documents published in a journal in the past two, thr ee and four y ears ha v e been cited in the curr ent y ear . The two y ears line is equiv alent t o journal impact fact or ™ (Thomson Reuters) metric.

Cites per document Y ear V alue Cites / Doc. (4 y ears) 1999 0.043 Cites / Doc. (4 y ears) 2000 0.101 Cites / Doc. (4 y ears) 2001 0.075 Cites / Doc. (4 y ears) 2002 0.084 Cites / Doc. (4 y ears) 2003 0.054 Cites / Doc. (4 y ears) 2004 0.068 Cites / Doc. (4 y ears) 2005 0.054 Cites / Doc. (4 y ears) 2006 0.062 Cites / Doc. (4 y ears) 2007 0.055 Cites / Doc (4 y ears) 2008 0 064 T otal Cites Self-Cites E v olution of the t otal number of citations and journal' s self-citations r eceiv ed b y a journal' s published documents during the thr ee pr e vious y ears.

Journal Self-citation is de ned as the number of citation fr om a journal citing ar ticle t o ar ticles published b y the same journal. Cites Y ear V alue S lf Cit 1999 0 External Cites per Doc Cites per Doc E v olution of the number of t otal citation per document and external citation per document (i.e. journal self- citations r emo v ed) r eceiv ed b y a journal' s published documents during the thr ee pr e vious y ears.

External citations ar e calculated b y subtr acting the number of self-citations fr om the t otal number of citations r eceiv ed b y the journal’ s documents. Cit Y V l % International Collabor ation International Collabor ation accounts for the ar ticles that ha v e been pr oduced b y r esear chers fr om se v er al countries. The char t shows the r atio of a journal' s documents signed b y r esear chers fr om mor e than one countr y; that is including mor e than one countr y addr ess.

Y ear International Collabor ation 1999 0.00 2000 0 00 Citable documents Non-citable documents Not e v er y ar ticle in a journal is consider ed primar y r esear ch and ther

efor e " citable ", this char t shows the r atio of a journal' s ar ticles including substantial r esear ch (r esear ch ar ticles, conf er ence papers and r e views) in thr ee y ear windows vs.

those documents other than r esear ch ar ticles, r e views and conf er ence papers. Documents Y ear V alue Cited documents Uncited documents Ratio of a journal' s items, gr ouped in thr ee y ears windows, that ha v e been cited at least once vs. those not cited during the following y ear .

Documents Y ear V alue Uncited documents 1999 89 Uncited documents 2000 158 Uncited documents 2001 201 Uncited documents 2002 225 ? Show this widget in y our own website Just cop y the code below and paste within y our html code: < a h r e f = " h t t p s : / / w w w . s c i m a g 0.21 0.28 0 0.06 0.12 0.18 0.24 0.3 1999 2002 2005 2008 2011 2014 2017 0 200 400 1999 2002 2005 2008 2011 2014 2017 0 0.2 0.4 1999 2002 2005 2008 2011 2014 2017 0 2.5

5 1999 2002 2005 2008 2011 2014 2017 0 600 1.2k 1999 2002 2005 2008 2011 2014 2017 0 600 1.2k S h o p p in g B a g (I t e m s) Download Sample Paper ISSN: 0971-765X Chief Editor Prof.(Dr .) R.K.T rivedy , Pune, India Editorial Board Download Sample Issue Subscription Rates Submit Article Click here for scimagojr .com SCOPUS - H Index - 1 1 NAAS Rating - 4.89 Home International Journals Books Environmental Consulting About Us Contact E c o l o g y , E n v i r o n m e n t a n d C o n s e r v a t i o n UGC, NEW DELHI,INDIA APPROVED JOURNAL B r i e f A b o u t E c o l o g y , E n v i r o n m e n t a n d C o n s e r v a t i o n P u b l i s h e d Q u a r t e r l y S i n c e 1 9 9 5 .

E c o l o g y , E n v i r o n m e n t a n d C o n s e r v a t i o n i s p u b l i s h e d i n M a r c h , J u n e , S e p t e m b e r a n d D e c e m b e r e v e r y y e a r . E C O L O G Y , E N V I R O N M E N T A N D C O N S E R V A T I O N i s o n e o f t h e l e a d i n g I n t e r n a t i o n a l e n v i r o n m e n t a l j o u r n a l .

I t i s w i d e l y s u b s r i b e d i n I n d i a a n d a b r o a d b y I n s t i t u t i o n s a n d I n d i v i d u a l s i n e d u c a t i o n a n d r e s e a r c h a s w e l l a s b y I n d u s t r i e s , G o v t . D e p a r t m e n t s a n d R e s e a r c h I n s t i t u t e s . E c o l o g y , E n v i r o n m e n t a n d C o n s e r v a t i o n i s i n M a s t e r J o u r n a l L i s t o f I S I (T h o m s

onReuters,U.S.

A.).Ecology,EnvironmentandConservationisabstracted/covered in:ChemicalAbstracts,U.S.A.SCOPUSEBSCOPublishing,U.S.A.CambridgeScienceAbstractsEcologyAbstractsPollutionAbstractsEco-DiscCDRomGeologicalAbstractsInternationalDevelopmentAbstractsOceanographicLiteratureReviewZoologicalRecordsIndianScienceAbstracts,Niscair,IndiaElsevier'sCompendexElsevier'sCurrentAwarenessinBiologicalSciencesElsevier'sEncompassElsevier'sGeobaseEcology,EnvironmentandConservationjournalisaccreditedwithNationalAcademyofAgriculturalsciences,NAAS,India.Ecology,EnvironmentandConservationjournaliscoveredbySCOPUS.

JournalIssuesVol 24, Issue 3 2018 Vol 24, Issue 2 2018 Vol 24, Issue 1 2018 Vol 24, March Suppl. Issue 2018 Vol 24, Feb. Suppl. Issue 2018 Vol 23, Issue 4, 2017 Vol 23, Nov . Suppl. Issue 2017 Vol 23, Sept. Suppl. Issue 2017 Vol 23, Issue 3, 2017 Vol 23, Issue 2, 2017 Vol 23, Issue 1, 2017 Vol 23, Feb 2017 Suppl. Issue Vol 22, Dec 2016 Suppl.

Issue Vol 22, Issue 4, 2016 Vol 22, Sept. Suppl. Issue , 2016 Vol 22, Issue 3, 2016 Vol. 22, June Suppl. Issue 2016 Vol 22, Issue 2, 2016 Vol. 22, April Suppl. Issue 2016 Vol 22, Issue 1, 2016 Vol 21, Issue 4, 2015 Vol. 21 Dec. 2015 Suppl. Issue Vol. 21 Nov . 2015 Suppl. Issue Vol 21, Issue 3, 2015 Vol. 21 Suppl.Issue August 2015 Vol 21, Issue 2, 2015 Vol 21.Suppl.Issue June 2015 Vol 21, Issue 1, 2015 Supplement Issue, Dec.

2014 Special Issue-2014 Vol 20, Issue 4, 2014 Vol 20, Issue 3, 2014 • • • • • • • • • • • • • • • • • • ShoppingBag(Items) Become a fan on Facebook Follow us on Twitter Home International Journals Books Environmental Consulting About Us Contact Ecology,EnvironmentandConservationEditorialAdvisoryBoardChiefEditorProf.(Dr.)R.K.

Trivedy,Pune,IndiaEDITORIALADVISORYBOARD1. Dr . Teresa

Ferreira, Portugal 2. Dr . Michael Ukwuru, Nigeria 3. Dr . Moses Inbaraj, Chennai 4. Dr . D.J. Lee, T aiwan 5. Dr . Christial Paul P .delacruz, Phillipnes 6. Dr . T . Bahorun, Mauritius 7. Dr . Linda Blackwell, Australia 8. Dr . G. Zellner , Netherlands 9. Dr . Wilson S. T isera, Kupang, Indonesia 10. Dr . M.F . Hamoda, Kuwait 1 1. Dr . H.A.Abrahamse, South Africa 12. Dr . Arulmozhiyal R., Salem 13. Dr . Hassan Ibrahim Ali, Sudan 14. Dr . A.R.Ghosh, Burdwan, India 15. Prof. M. Zaman, Bangladesh 16. Dr .

Marcantonio Bragdin, V enice, Italy 17. Dr . Z. Fuat T opark, T urkey 18. Dr . Z. Li. Bonn, Germany 19. Dr . A. Olawale, Nigeria 20. Dr . Ing. Agr . Mario Ridardo Sabbatini, Argentina 21. Dr . Philip C. Reid, U.K. 22. Dr . Bajcinovci, B. Kosovar , Bosnia 23. Dr . Mohd. Y usuf, Malaysia 24. Dr . Oswaldo A. Feernandez, Argentina 25. Dr . Ms. Mirela T ulik, W arsaw , Poland 26. Dr . L.L. Chukwu, Nigeria 27. Dr . Azni Idris, UPM, Malaysia 28. Dr . G.

Suresha, Saudi Arabia 29. Dr . Amresh Chandra Pandey , Jharkhand, India 30. Dr . Shambhu Sharan Kumar , Ranchi, India 31. Dr . A.K. Panigrahi, Berhampur , India 32. Dr . Ahmed EI Mahmoudi, Saudi Arabia 33. Dr . Seyed Mohammad T ajbakhsh, Iran 34. Dr . Amin L. Setyo, Indonesia 35. Dr . Francis Gbogbo, Ghana 36. Dr . S.

Shabanlou, Iran Back to EEC Journal Details Home | International Journals | Books | Environmental Consulting | About Us | Contact Us | Submit Paper | © EM International 2012 | ECOLOGY, ENVIRONMENT AND CONSERVATION VOL. 24 (4) : 2018 CONTENTS 1467–1474 Sustainable fisheries resource management based on local knowledge of pole and line fishers (Case study, Murante Villages, Luwu Regency, South Sulawesi, Indonesia) —Amilludin, Yusran NurIndar, Aris Baso, Najamuddin and Andi Adri Arief 1475–1481 Economic development of the republic Mari El on the basis of natural resource-based multicluster —Nina I. Larionova, Tatyana V. Yalyalieva and Dmitry L.

Napolskikh 1482–1491 Function of a temporary wetland for wintering anatidea and coot: Mekhada Marsh (North East Algeria) —Rima Bendjeddou, Zahra Benyacoub Brahmia and Slim Benyacoub 1492–1500 Review and Implementation of the accident analysis methods (AAM) in the Oil and GAS Upstream sector —Nikolaos Georgakopoulos 1501–1507 Determining crop planting pattern and sorting base on priority farmers' subjective —Parisa Shahinrokhsar, Amin Alizadeh, Hossein Ansary and

Mohamad Ghorbany 1508–1514 Soil contamination and assessment of heavy metals neartwo landfills area due to leachate migration —Tafë Veselaj, Rifat Morina, Afrim Berisha, Valdet Gashi and Fatbardh Sallaku 1515–1522 Ecotourism model based on social asset (Halal Tourism) as the strenghthening of economy of Lombok community, Indonesia —Wildan and Sukardi 1523–1529 Riple harmonization of transcontinental allometric of Picea spp. and Abies spp.

forest stand biomass —Vladimir Andreevich Usoltsev, Seyed Omid Reza Shobairi and Viktor Petrovich Chasovskikh 1530–1534 Preliminary study single stage micro gas turbine integrated with South Sumatera Indonesia low rank coal gasification —Fajri Vidian, Hasan Basri, Helmy Alian, Elian Zhafran and Thorik Aziad 1535–1540 Results of preliminary studies on the production and agrucultural crops research of "JAMB- 70" polycomponent mineral fertilizers —Zh. N. Rakhmanberdiyeva, K. T.

Zhantasov, Z. K. Myrzaliyeva, B. R. Naimanbayev and D.M. Zhantasova 1541–1547 Spatial modeling vocational education development to support regional potential —Agus Wiyono, Abdul Hakim, Endah Setyowati and Agus Suharyanto 1548–1556 Evaluation of conditions for effective agricultural land-use as a basis for sustainable development of plant-growing production in the Omsk Region —Yuri Mikhailovich Rogatnev, Olga Nikolaevna Dolmatova, Vitaliy Viktorovich Aleschenko, Marina Nikolaevna Veselova and Yulia Stanislavovna Yusova 1557-1562 The monitoring of San Jose scale (Homoptera, Diaspididae: Quadraspidiotusperniciosus Comstock, 1881) in the southeast of Kazakhstan —Z.B. Beknazarova, B. Lozowicka, B.K.

Kopzhasarov and R.A. Iskendirova 1563–1567 Multiplet interrelations in Oil – Water compositions —Vinera Bekbayeva, Ashimhan Toktasynovich Kanaev, Galina Metaxa and Zy Losinq 1568–1575 Domestic and industrial waste as the basis for formation of compound compost —I. S. Belyuchenko II CONTENTS Eco. Env. & Cons.

24 (4) : 2018 1576–1586 The estimation of the sevastopol bays ecological state on basic chemical and microbiological criteria —Tikhonova E. A., Burdiyan N. V., Soloveva O. V., Doroshenko Yu. V. 1587–1591 The effect of biological fertilizers on productivity of clover on saline soils of rice systems of the Aral Sea Region, Kazakhstan —Zhanar Shadibekovna Zhumadilova, Ibadulla Aigalievich Tautenov and Eric Zharylkasynovich

Shorabaev 1592–1601 Feasibility study of cantrang (Danish Trawl): fisheriesbiology perspective —Sri Suro Adhawati, Achmar Mallawa, Aris Baso and A.

Adri Arief 1602–1608 Immune -phytopathological assessment of resistance of wheats to stem rust in conditions of the Southeast of Kazakhstan —Z. Amangeldikyzy, A.S. Kochorov, Aziz Karakaya, A.I. Morgunov, R.D. Karbozova and M.A. Gabdulov 1609–1616 Changing the properties of cultivated land is a real basis for the formation of new environmental niches —I.S. Belyuchenko and O.A.

Melnik 1617–1623 The influence of seeding time on growth development and productivity of sunflower in the dry steppe area —Beybit Nasiyev, Nurbîlat Zhanatalapov and Alexander Bushnev 1624–1629 Effects of human pressure on the distribution of Macaca sylvanus in the sector of Ait ouabane, National Park of the Djurdjura, Algeria — Mohammed Oudahmane, Aissa Moali and Farid Bekdouche 1630–1633 Carbon dioxide (CO 2) absorption of several landscape plants through photosynthetic route —F.

Fathurrahman, Nizam Mohd Said 3,4, Wan Juliana Wan Ahmad, and Che Radziah Che Mohd Zain 1634–1639 Assessment of lead content in soil and plants as a consequence of continuous application of fertilizer systems —R. F. Baibekov, A. N. Esaulkov, M. S. Sigida, Y. I. Grechishkina and A. V. Voskoboinikov 1640–1644 The predation efficacy of Coccinella septempunctata L.

(Coccinellidae: Coleoptera) against cotton aphid Aphis gossypii glover (Homoptera: Aphididae) —Ali K. R. Al-Shujairi and Sindab S. J. Al-Dahwy 1645–1650 The selection of parental material for winter barley breeding Inboharic direction —Zhanbolat Serkebaevich Mussabaev, Galiolla Tulendinovich Meirmanov, Anarbai Kainarovich Ortaev and Hristofor Kirchev 1651–1656 Assessment of recovery of medicinal plants of the Kurtidistrict of the Almaty region —A.S. Seilkhan, R.À.

Mirzadinov, A. Aksoy, A. Abulgaziyev and G. Kanat 1657–1663 The study of the importance of cytogenetic and molecular genetics in the diagnosis of fish diseases — Mohsen Shokohinia 1664–1668 Studying the technology and methods of increasing the yield of cultivated plants on strongly saline soils —Aliya Ainabekovna Ismailova, Ashimhan Toktasynovich Kanaev, Nariman Zhalgassuly, Mahaya Aisijiang and Alexander

Vladimirovich Kogut 1669–1676 Assessment of recovered organic waste quality for application to sandy soils in the United Arab Emirates —Maitha Al Majid Almheiri, James Peter Terry and Munawwar Ali Khan 1677–1683 Characteristic of stands of forest Fringillidae in the El Kala region (Northeastern Algeria) —Siouane Noureddine, Ziane Nadia and Slim Benyacoub Eco. Env. & Cons.

24 (4) : 2018 CONTENTS III 1684–1689 Discrimination of tropical Mangrove species with hyperspectral leaf reflectance —Pandi Selvam Pandi, Ramesh Ramachandran, Purvaja Ramachandran and Srinivasalu Sechachalam 1690–1695 Zero waste management and green steel by replacement of steel mill scale in Eco friendly construction —P. Ganeshprabhu and P.

Chandrasekaran 1696–1701 Ecological potential of highloft textiles from jute blended nonwovens for sound and thermal insulation —Bharanitharan Ramanathan, K. Thangamani and S. Sundaresan 1702–1711 Environmental effects of exposure to cellphone radiations and its impacts on human health —A. Adappa Sholapuri and Shrinivas Rao 1712–1719 Economic and environmental impacts of producer Company in Coimbatore District, Tamil Nadu —V.

Jagadeesh Pandian and Madhavi Ganesan 1720–1725 Improving energy proficiency and trust worthiness for power plant monitoring in WSN —Sathiyaseelan Rathinavel 1726–1731 Study on utilization of waste plastic fibres as a construction entity —S. Karthikeyan and G. Vennila 1732–1737 Performance of different capsicum hybrids for earliness and yield under various protected structures —Daljit Singh Gill1,3, Ramesh Kumar Sadawarti1 and Madhu Sharma2 1738–1753 Implication of woody species diversity on the aboveground biomass in subtropical sacred groves of Manipur, north east India —Chongtham Sanjita, TH. Binoy Singh and TH. Rojen Singh 1754–1759 Degradation of reactive dyes in textile dyeing effluent by fenton and electro fenton processes —A.

Latha, P. Partheeban and R. Ganesan 1760–1766 Problem associated with breeding of two commercially important catfish Clarias magur (Hamilton, 1822) and Heteropneustes fossilis (Bloch, 1974): alternative approaches for solution —Pradip Kumar Maurya, Shubham Gupta, Arvind Kumar Verma and Sullip Kumar Majhi 1767–1775 In-vitro efficacy of fungicides, bioagents and plant leaf extracts against Sclerotium rolfsii causing

collar rot disease of french bean (Phaseolus vulgaris L.) —Ashwini Kumar, N.

Kudada and Amit Saurabh 1776–1779 Mushroom as food and medicine for Wellness and Wellbeing —A.S. Deshmuknh 1780–1784 Effect of sewage irrigation on physio-biochemical characterization of two Mulberry varieties (Morus alba and Morus serrata) —M. Aslam Abdullah, G.S. Nirmala, Aruna Singh, Rajan Jain, Sindhu Manchikanti and I.

Ciurcanu 1785–1793 Impact of sub watershed implementation at Pandoga in swan river catchment area of shivalik foothills H.P, India on life form and biological spectrum — Bindu Sharma, Sujata Bhattacharya and Sunil Puri 1794–1806 Investigation of physicochemical parameters to assess groundwater quality due to solid waste dumping - A case study —P. Vijayalakshmi and Marykutty Abraham IV CONTENTS Eco. Env. & Cons.

24 (4) : 2018 1807–1810 Sexual dimorphism of body size variations and mating behaviour in natural populations of Drosophila melanogaster —Veer Bhan 1811–1818 Mathematical modeling and analysis of domestic wastewater treatment in Phagwara using moving bed biofilm reactor —Bimenyimana Hubert and Misbar Bashir 1819–1823 Effect of city sewage wastewater irrigation on the nutrient and heavy metal status of soil and crops —Afroza Akhter, Abdul Rashid Dar, Irfana Amin and Arif Inam 1824–1826 Weed biodiversity and phytosociological composition of Bhadohi District of Uttar Pradesh —Ranjit Singh and P.N.

Dongre 1827–1829 Consequence of changeable concentration of Cr (VI) on a bacterial isolate —Ravneet Chug and Manishita Das Mukherji 1830–1836 Behaviour and the influence of meteorological parameters towards air pollutant dispersion modelling studies, Manali Industrial area, Chennai —J. Karthick and R. Samuel Devadoss 1837–1842 Isolation and identification of bioactive protein from fish by products —Indira N., Valivittan K. and P.

Dhasarathan 1843–1846 Modelling of hybrid energy systems for renewable energy conservation and pollution control —S. Arulanantha Samy, S. Venkatesh, S.Vasanth and P. Satyanathan 1847–1854 Madhav national park: A potential site for vulture in Bundelkhand region —Ruby Yadav, Adesh Kumar and Amita Kanaujia 1855–1858

Isolation, screening and identification of cellulose producing bacteria from rotten rruits —N.C. Tharavathy and M.H.

Attiya Ilham 1859–1865 Physico-chemical analysis of natural water sources at Thengaipattinam, Kannyakumari District, Tamilnadu, India —Indira N., Arunthathi R., Valivittan K. and P. Dhasarathan 1866–1874 Diversity and abundance of coral reef fish species in the coastal waters of salahutu sub- District, central Moluccas regency – Indonesia —Hasan Tuaputty, Marleny Leasa and John Rafafy Batlolona 1875–1878 Comparative evaluation of wet Lignocellulosic biomass for biocrude oil production —R.

Divyabharathi, P. Subramanian, S. Pugalendhi, S. Avudainayagam and D. Manohar Jesudas 1879–1884 Case study of biogas production from various feedstocks —Shivani and Bashir Misbah 1885–1894 Soil Co 2 flux and temperature sensitivity (Q 10) in natural and human impacted ecosystems of senapati District, Manipur, India —Ng Niirou and Asha Gupta 1895–1902 Biodiversity status and conservational requirements of tropical Tasar (Antheraea mylitta D.)

— A review —Khasru Alam, Debjoy Bhattacharryay, Soumen Saha and Prasanta Kumar Kar 1903–1913 Bacterial community structure (V3 Metagenomics) of Ashtamudi estuarine sediment, Kerala, India, by 16s RRNA amplicon sequencing —R. R. Reshmi and Salom Gnana Thanga, Vincent 1914–1923 Modelling the relationship between land use and surface water quality —Mahdi Mahmoodi, Mahdi Honarmand, Farzin Naseri and Sedigheh Mohammadi Eco. Env. & Cons.

24 (4) : 2018 CONTENTS V 1924–1930 Augmenting water stress tolerance in Tomato by silicon supplementation —Chanchal Malhotra, Riti Thapar Kapoor, Deepak Ganjewala and N. B. Singh 1931–1939 Assessing public awareness level on the preservation of coral reefs (The case study in Biak Numfor, Papua, Indonesia) —Lis Melissa Yapanto and Meilinda Lestari Modjo 1940–1944 Model multivariate adaptive regression spline on lead exposure found within the hair of Petrol station's workers in Gorontalo City, Indonesia —Herlina Jusuf 1945–1953 Flood disaster preventive measures using GIS and multicriteria technique in the water sheds area of Ambasamuthiram Town —M.

Seenirajan, M. Natarajan, R. Thangaraj and S. Shanmugasundaram 1954–1959 Ranging

and feeding ecology of Javan langur (Trachypithecus auratus) rehabilitation in coban Talun protected forest, Batu, East Java, Indonesia —Emanuel Naitio, Amin Setyo Leksono and Bagyo Yanuwiadi 1960–1965 Inter-dependency of forest diversity and service towards the potency of ecotourism development in Pegunungan Arfak nature reserve — Lukas Yowel Sonbait, Hermanus Warmetan, Hotlan Manik and Reinardus Liborius Cabuy 1966–1972 Triple harmonization of transcontinental allometric of Picea spp. and Abies spp.

forest stand biomass —Vladimir Andreevich Usoltsev, Seyed Omid Reza Shobairi and Viktor Petrovich Chasovskikh Article-70 Eco. Env. & Cons. 24 (4) : 2018; pp. (1940-1944) Copyright@ EM International ISSN 0971–765X Model multivariate adaptive regression spline on lead exposure found within the hair of Petrol station's workers in Gorontalo City, Indonesia Herlina Jusuf Faculty of Public Health, Universitas Negeri Gorontalo, Indonesia (Recieved 5 June, 2018; accepted 20 August, 2018) ABSTRACT MARS (Multivariate Adaptive Regression Spline) is one of the non-parametric regression models that employs modified recursive partitioning algorithm.

This study applied MARS model in the data of Lead (Pb) exposure in the workers' hair at the petrol station in Gorontalo City, Indonesia. The study is intended to investigate the content of Lead in one's body. The results of the study reveal that the MARS model in the Lead exposure in the workers' hair at the petrol station in Gorontalo City consists of Y = 1.93593 - 0.219669 * BF8 - 0.0220152 * BF13.

Based on that MARS model, there are only three variables out of 10 assumed to affect the Lead exposure in the workers' hair at the petrol station in Gorontalo City including the working time, disease symptoms, and age. Key words: Multivariate, Adaptive, Regression, Spline, Lead exposure, Hair. Introduction Time series modelling is usually found in suffi- ciently stationary and linear data by application of Autoregressive Integrated Moving Average (ARIMA) method for prediction (Arsyad, 1999). This method is highly effective, and model that is highly suitable will be achieved when the two re- quirements above are met.

If the data could not ful- fil the assumption, then the obtained model would not be complete in describing the pattern of the sys- tem behaviour, hence, unsuitable for

forecasting. Development of non-linear time series has been made through the introduction of several flexible methods in various applications, where Multivari- ate Adaptive Regression Splines (MARS) is one of them (Friedman, 1991).

Fluctuated and non linear data and time series modelling to obtain the value of given period in the future was proposed through MARS approach, which was considered better than the stochastic model (Buja, Duffy, Hastie, Tibshirani, 2001). One of the chemical elements that are dangerous to human health is the heavy metal. There are sev- eral heavy metals that are toxic and acts as a pollut- ant in the environment such as mercury (Hg), lead (Pb), Cadmium (Cd), and Cuprum (Cu).

The lead that polluted the air can come from the burning resi- due of additive substance from the motor vehicles' fuel. The lead particles in the air can also come from other sources such as Allkil Pb factory and Pb-oxide, coal burning, etc (Sunu, 2001). The lead polluted air can enter the body through inhalation process. Most of the inhaled lead will enter the blood vessels in the lungs.

Level of lead's absorption is heavily influ- enced by the size of the lead particle and the inhaled blood volume during the breathing process. The lead inhaled into the lungs will be absorbed and HERLINA JUSUF 1941 chained with blood in the lungs to further be distrib- uted to all tissues and organs. In tissues or organs, the inhaled lead will be accumulated within the bone. Regardless, to the small amount of inhaled lead, this metal can be hazardous.

This was due to the lead compounds that are toxic to many func- tions of the body organs (Palar, 1994). The other mechanism for the lead to enter the human body is through oral or direct contact with the skin surface. About 40% of lead within the body which comes from inhalation process is absorbed into the respira- tory system.

About 5-10% of the lead compounds were absorbed into the gastrointestinal tract (Naria, 2005). The danger of the inhaled motored-vehicles' emission has a long-term effect on human body. The emission that contained lead is found in vehicles that used leaded-fuels as anti-knock compounds.

The lead element in the body can be deposited into the soft tissues (bone marrow, nerve systems, kid- ney, and liver) and hard tissues (bones, teeth, nails, and hair). The lead element within the soft tissues is toxic within that soft tissues itself (Ardiyanto, 2005), whereas the intrusion of a large quantity of lead into the human body could cause kidney failure (Soesanto, 1998).

It can also cause cancer, problems in peripheral and central nervous system, digestion system, and reproduction system (Ardyanto, 2005). Also, in a pregnant woman, lead can penetrate the placenta and enter the blood circulation of the fetus. The lead will be disposed of with the breast milk after the baby was born (Palar, 1994). There has been extensive research on the level of lead within the body.

However, the focus of this in- vestigation is on the petrol's station workers in Gorontalo regency. Therefore, a research to find out the level of lead accumulated within the petrol sta- tion workers' body is needed. Lead Found in Hair Hair is a solid structure which consists of solid kera- tin cells from made from epidermis follicles which shaped like a bulb that grows into the dermis layer. Normal and healthy hair looks shiny, elastic, and not easily broken, as well as absorbs water. Hair composition consists of 50.60% of carbon, 6.36% of hydrogen, 17.14% of nitrogen, 5.0% of Sulphur, and 20.80% of oxygen (Pusponegoro, Erdina H.D. 2002).

Hair is also one of the skin adnexa that spread along the body except for the palm of the hands, soles of the feet, nails, and lips (Soepardiman, Lily. 2002). There are two generic types of hair in the human body, terminal hair, the pigmented hair that grows on the head, brow, eye lashes, and in the pubic re- gions, and vellus hair, the least pigmented hair that grows in almost all body parts. (Soepardiman, 2002).

In the human body, the lead will be disposed through the various mechanisms, including through hair. Considering that hair reflects more on the level of a heavy lead pollutant that has entered the body (Kamal, 2007). This is because hair contains a lot of structural proteins that composed of cysteine amino acid that contains disulphide bonds (- S - S -) and cysteine containing sulfhydryl groups (- SH) with the ability to bind the heavy metals that enter the human body.

The level of tolerated lead in human hair is 12? g/g, in this level the toxin within the lead is not harmful (Palar, 1994). Factors that influence the level of lead in human hair are the length of exposure, age, genetic, and nutrition, hence, hair can be used as an indicator for the level of lead pollution in human and animal (Ashraf et al. 1995).

Multivariate Adaptive Regression Spline MARS is one of the non-parametric regression model which uses modified-recursive partitioning algorithm. The MARS modelling is preceded by three things below (Friedman & Silverman, 1989). i. Base Function (BF) is a function defined from each region. A maximum number of recom- mended base functions (BF) are 2-4 times of the number of the predictor variable. ii.

Maximum interaction (MI) is a number of inter- action that happens on the model. The number of maximum interaction (MI) are 1, 2, and 3, if the maximum interaction used is more than 3, then the CGV value will increase and the model used will be more complex. If the maximum interac- tion used is 1, it means there is no interaction among variables within the model.

If the maxi- mum interaction used is 2, then there is an inter- action between two variables within the model. If the maximum interaction used is 3, it means that some interaction that happens within the model is mostly among three variables. iii.Minimum observation (MO) is a number of mini- mum observations among knots. A number of maximum interactions (MI) are 0,1,2, and 3, be- yond this figure, the CGV will increase.

Establishment of the value of the base function 1942 Eco. Env. & Cons. 24 (4) : 2018 (BF), maximum interaction (MI) and minimum ob- servation will influence the MARS model that will be developed. The multivariate adaptive regression splines (MARS) can be written in the following for- mula ?? ?? ?? 0 , 1 1 ˆ .

m KM m km km v km m k f x S x t ?? ?? ? ?????? ? ? Research Method This study is an observational research with cross- sectional approach. This study is conducted on petrol stations' workers that work in seven petrol stations in Gorontalo city with a total number of 48 workers. The samples are taken between 07.00 – 11.00 am because that period is the peak time for vehicles to fill their petrol tanks in these stations.

The testing of the hair samples is conducted in Balai Pengawasan Mutu Perikanan (fisheries quality control office, henceforth called as BPMP) of Gorontalo city. The population in this study is all the workers of petrol stations in Gorontalo city. Samples are taken using purposive sampling method based on several criteria. The samples' criteria are as follow: (1) Age 20-40 years old; (2) Hair length minimum 5 cm; (3) Work tenure ? 2 years and (4) Volunteer to become respondents Research Variables The were three variables using of this research. These research Variables are: (1) Work Tenure.

It is about the length of time where respondents, in their line of work have been exposed to lead. The longer the working tenure, the higher the risk to be ex- posed to lead. The objective criteria are short work tenure (? 2 years) and long work tenure (> 2 years); (2) Level of Lead (Pb). Number of lead accumulated in respondents' body.

The tolerated lead exposure on human hair is ? 12 µg/g, hence, the toxicity is not severe (WHO, 1994). The objective criteria are nor- mal, when the lead level is ? 12 µg/g and abnormal when the level of lead is > 12 µg/g and (3) Health problems on petrol stations' workers. Health con- cerns are often found in people who have been ex- posed to lead.

The target of this study is the lead in the hair of petrol stations' workers in the area of Gorontalo city. Primary data are obtained through an interview with additional observation sheet that contains the questions such as the identity of the respondents (name, age, and respondents' work history, medical history).

The respondents' hair specimens are col- lected and tested on the level of lead in respondents' hair is conducted in the laboratory. Secondary data used in this study are data from BLHRD (Provincial Environment and Research Agency) of Gorontalo province, Samsat Office (Vehicle Registration office) of Gorontalo city, and Badan Pusat Statistik (statisti- cal bureau) of Gorontalo Province.

Tool, Technique and Research Procedure The method used in analysing the specimens in this study is Atomic Absorption Spectrophotometry (AAS). The samples are tested at BPMP laboratory of Gorontalo city. i. Tools and equipment • Hair scissor used to cut the respondents' hair • Plastic bag • Observation sheet • Acetone and water • beaker glass •

volumetric flask • pipet • mortar and pestle • vial polyethylene • PC computer • Digital Scale • Kooling Module (KMS) or water cooling system • Atomic Absorption Spectrophotometry as tool of analysis • Blower • Petrol stations workers' hair • Condensed Nitric Acid (HNO3) • Perchlorate Acid • Aqua dest ii.

Hair specimen collection technique Each respondent hair was taken 0.5 – 1 cm, then the hair is put into a labelled plastic bag. Respondents' data are also collected, such as the name, age, and work tenure. i. Preparation Technique and AAS analysis Prior to hair specimens' analysis, the specimens are washed with 100 mL of acetone and rinsed three times with water.

After that, the hair is washed us- ing the acetone to get rid of the fat and other con- taminants that might interfere other elements in hair. The specimens then dried. The specimen then destroyed to obtain a homogenous result. The result is then put into a vial, each specimen is given a la- bel. HERLINA JUSUF 1943 Research Findings and Discussion In Mars modeling process, there are three things to be considered: the base function (BS), maximum in- teraction (MI), and minimum observation (MO). The base function (BF) is a function defined from each region.

The maximum number of recom- mended base function (BF) is 2-4 times of the num- ber of the predictor variable (Friedman, 1991). The predictor variables that are suspected to influence the lead exposure are ten variables, hence, the num- ber of base function that will be combined in the establishment of the model are 20, 30, and 40. Maximum interaction (MI) is the number of inter- action that happens on the model.

The number of maximum interaction (MI) is 1, 2, and 3 (Friedman, 1991). If the maximum interaction used is more than 3, then the CGV value will increase and the model used will be more complex. Minimum observation (MO) is number of mini- mum observation among knots.

Number of maxi- mum interaction (MI) is 0, 1, 2 and 3, beyond this figure, the CGV will increase. The establishment of MARS model is conducted through trial and error for all combination of the BF, MI, and MO values that have been previously deter- mined. The possible models based on those combi- nations are 36 models.

From each of these modellings, GCV values will be produced and cre- ated predictor variables included into the model. The result is presented in Table 1 below. Based on the table above, the smallest GCV val- ues are found in model number 11, 23, and 35. Be- cause these three models yield the same result, the model with the smallest MI and MO values are se- lected.

Hence, the chosen model is model number 11, where the value of GVC is 0.08952 with the com- bination of BF=20, MI=3 and MO=2. The MARS model equation is Y = 1.93593 - 0.219669 * BF8 - 0.0220152 * BF13 where, BF6 = (×5 in (1)); BF8 = (×10 in (2)) * BF6; BF13 = max (0, 39 - X1) * BF8; From this model, it is clear that out of ten vari- ables that are assumed to have an influence on the model, only three variables that are proven to have an influence on lead exposure.

Because the best model is a model with maximum two interactions, it is suspected that there are factors that are interact- ing with one another. This model consists of one intercept (base function master) and three base func- tions that consist two level-2 interactions. To examine the variables that influence the lead exposure, interpretation of the MARS model is pre- sented below.

Interpretation of this MARS model is quite tricky because the base function in this model consists of not only one variable, but there is also the interaction between variables. • BF8 = (X10 in (2)) * BF6, where BF6 = (X5 in (1) It means that BF* coefficient will be meaningful if respondents were indicated to have some com- plaints and the length of work shift in petrol station was less than eight hours, then each unit increase in base function BF8 will decrease the lead exposure Table 1.

Trial and Error Result for All BF, MI, and MO Combinations BF MI MO GCV 1 20 1 0 0.11958 2 20 1 1 0.11905 3 20 1 2 0.11888 4 20 1 3 0.11905 5 20 2 0 0.11518 6 20 2 1 0.09693 7 20 2 2 0.10564 8 20 2 3 0.09838 9 20 3 0 0.11553 10 20 3 1 0.10139 11 20 3 2 0.08952 12 20 3 3 0.09931 13 30 1 0 0.11933 14 30 1 1 0.11871 15 30 1 2 0.11856 16 30 1 3 0.11892 17 30 2 0 0.11518 18 30 2 1 0.09693 19 30 2 2 0.10564 20 30 2 3 0.09838 21 30 3 0 0.09838 22 30 3 1 0.10139 23 30 3 2 0.08952 24 30 3 3 0.09931 25 40 1 0 0.11933 26 40 1 1 0.11871 27 40 1 2 0.11856 28 40 1 3 0.11892 29 40 2 0 0.11518 30 40 2 1

0.09693 31 40 2 2 0.10564 32 40 2 3 0.09838 33 40 3 0 0.11553 34 40 3 1 0.10139 35 40 3 2 0.08952 36 40 3 3 0.09931 1944 Eco. Env. & Cons. 24 (4) : 2018 by 0.219669 percent. • BF13 = max (0.

39 - X1) * BF8, where BF8 = (×10 in (2) * BF6 and BF6 = X5 in (1) It means that coefficient BF8 =13 will be mean- ingful if the respondent's age were less than 39 years old and indicated health complaint and length of work shift in petrol station was less than 8 hours, then each unit increase in base function BF8 will decrease lead exposure by 0.0220152 percent.

Mars modelling in this study showed that there are interactions among predictor variables, which in turn influence the responds' variables as shown in the following table Table 2 in general shows the significant interac- tion among variables in each base function. The three influential predictor variables indicate that each variable contributes to the responds' variable. Level of variable's importance is the extent of 0.219669 * BF8 - 0.0220152 * BF13.

From this model it is obtained that out of 10 variables that are sus- pected to have an influence on the model, there are only three variables that have the influence on lead exposure namely, work tenure in a petrol station, health complaint indication, and age. Tabel 3. Level of Predictor's Variable Importance Variable Level of Importance GCV Work tenure 100.00000 0.14486 Indicated health complaint 100.00000 0.14486 Age 38.43010 0.08373 Table 2.

Interaction in Base Function BF Interaction 8 Indicated to have health complaint and length of work shift in petrol station 13 Age indicated to have health complaint and length of work shift in petrol station predictor's variable contribution toward the MARS model. Level of predictor's variable importance in grouped function is calculated by the increase in GCV value.

The increase of GVC value is due to the migrations of considered variables within the model. The importance level of predictor's variable is shown in the following table. Table 3 shows that work tenure in petrol stations and health complaint indication have the biggest contribution by 100% in determining the lead expo- sure risk.

The second contributed variable by 38.43% is the age of the respondents. Conclusion Based on the findings and analysis discussed above, the following things are concluded in this study: MARS equation model obtained is:Y = 1.93593

INTERNET SOURCES:

1% - http://sarwonohm.com/wp-content/uploads/2017/11/Ijaer-v9-n22-2014-factor-analysis-on-causal-w-sjr.pdf
<1% - https://secure.in.gov/sba/files/rev_forecast_20151217_Economic_Outlook.pdf
<1% - http://www.envirobiotechjournals.com/journal_details.php?jid=3
<1% - https://www.texasgop.org/wp-content/uploads/2018/06/PLATFORM-for-voting.pdf
<1% - https://www.uschamber.com/sites/default/files/legacy/international/mideast/files/uschamberiraqbusinessinitativemission.pdf
<1% - http://www.fao.org/docrep/017/i3225e/i3225e.pdf
<1% - https://www.researchgate.net/publication/229118655_Landscape_evolution_space_and_the_relative_importance_of_geomorphic_processes_and_controls
<1% - http://www.envirobiotechjournals.com/EEC_editorial_board.php
1% - https://www.researchgate.net/publication/329402772_TRIPLE_HARMONIZATION_OF_TRANSCONTINENTAL_ALLOMETRIC_OF_PICEA_SPP_AND_ABIES_SPP_FOREST_STAND_BIOMASS
<1% - http://europepmc.org/articles/PMC2999420
<1% - https://sullipkm.weebly.com/publications.html
<1% - http://www.yspuniversity.ac.in/mpp/thesis/mpp-thesis-index.htm

CERTIFICATE OF ORIGINALITY

To Whom It May Concern:

This is to certify that the following document has been checked for originality with premium plagiarism checker. The result is as follows:

Originality Report	
Document Title	Model multivariate adaptive regression spline on lead exposure found within the hair of Petrol station's workers in Gorontalo City, Indonesia
Author(s)	Herlina Jusuf
Similarity Found	2%
Statistics	188 words Plagiarized / 7808 Total words
Remark(s)	Low Plagiarism detected
Internet Sources	
1% - http://sarwonohm.com/wp-content/uploads/2017/11/Ijaer-v9-n22-2014-factor-analysis-on-causal-w-sjr.pdf <1% - https://secure.in.gov/sba/files/rev_forecast_20151217_Economic_Outlook.pdf <1% - http://www.envirobiotechjournals.com/journal_details.php?jid=3 <1% - https://www.texasgop.org/wp-content/uploads/2018/06/PLATFORM-for-voting.pdf <1% - https://www.uschamber.com/sites/default/files/legacy/international/mideast/files/uschamberiraqbusinessinitativemission.pdf <1% - http://www.fao.org/docrep/017/i3225e/i3225e.pdf	

<1% -
https://www.researchgate.net/publication/229118655_Landscape_evolution_space_and_the_relative_importance_of_geomorphic_processes_and_controls
<1% - http://www.envirobiotechjournals.com/EEC_editorial_board.php
1% -
https://www.researchgate.net/publication/329402772_TRIPLE_HARMONIZATION_OF_TRANSCONTINENTAL_ALLOMETRIC_OF_PICEA_SPP_AND_ABIES_SPP_FOREST_STAND_BIOMASS
<1% - http://europepmc.org/articles/PMC2999420
<1% - https://sullipkm.weebly.com/publications.html
<1% - http://www.yspuniversity.ac.in/mpp/thesis/mpp-thesis-index.htm

Date: Wednesday, January 23, 2019

Novriyanto Napu, M.AppLing., Ph.D.
Director