




Professional Services Software

Advanced Planning Tools for Efficient Project Resource Allocation.

Wrike

OPEN

Agrivita

Country	Indonesia -  SIR Ranking of Indonesia
Subject Area and Category	Agricultural and Biological Sciences Agronomy and Crop Science
Publisher	University Of Brawijaya
Publication type	Journals
ISSN	01260537
Coverage	2013-ongoing
Scope	The aims of the journal are to publish and disseminate high quality, original research papers and article review in plant science i.e.: -agronomy -horticulture -plant breeding -soil sciences -plant protection -other pertinent field related to plant production
	Homepage
	How to publish in this journal
	Contact
	Join the conversation about this journal

5

H Index


Linguistics Journal

Cutting-Edge Research in Linguistics


degruyter.com

OPEN


Agronomy and Crop Science




SJR




Citations per document




Total Cites Self-Cites




External Cites per Doc Cites per Doc




% International Collaboration



Citable documents Non-citable documents



Cited documents Uncited documents



Agrivita

Q3

Agronomy and Crop Science

best quartile

SJR 2018

0.22

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

``

AGRIVITA, Journal of Agricultural Science

AGRIVITA *Journal of Agricultural Science* is a peer-reviewed, scientific journal published by Faculty of Agriculture Universitas Brawijaya Indonesia in collaboration with Indonesian Agronomy Association (PERAGI). The aims of the journal are to publish and disseminate high quality, original research papers and article review in plant science i.e. agronomy, horticulture, plant breeding, soil sciences, plant protection and other pertinent field related to plant production. AGRIVITA is published three times per year.

The Journal has been indexed in SCOPUS, Scimago Journal Ranks (SJR), Emerging Source Citation Index (ESCI-Web of Science), EBSCO, ProQuest, Google Scholar and others international indexing.

AGRIVITA is accredited first grade (Sinta 1/S1) for five years (2018-2023) based on Decree No: 30/E/KPT/2018 by Ministry of Research, Technology and Higher Education (Ristek Dikti), The Republic of Indonesia.

We accept submission from all over the world. All submitted articles shall never been published elsewhere, original and not under consideration for other publication.

Announcements

News: DOI for 2012-2015 Edition

For citation purpose of 2012-2015 edition, we inform you to use DOI that stated in the document (not in website page).

If you have question please feel free to email us at agrivita@ub.ac.id

Posted: 2019-07-08

[More...](#)

News: AGRIVITA June 2019, Coming Soon

AGRIVITA *Journal of agricultural science* Vol.41 No 2 June 2019 Edition will be published.

Posted: 2019-05-27

News: IMPORTANT ANNOUNCEMENT !

AGRIVITA *Journal of Agricultural science* is Accredited First Grade

Posted: 2019-02-10

[More...](#)

[More Announcements...](#)

About Agrivita

- [Aim and Scope](#)
- [Editorial Team](#)
- [Publication Ethics](#)
- [Visitor Statistics](#)
- [Reviewer Acknow](#)

Issues

- [Current Issue](#)
- [Back Issues](#)
- [Article in Press](#)
- [Accepted Papers](#)

Information For

- [Author Guideline:](#)
- [Template \(docx\)](#)
- [Template \(pdf\)](#)

User

Username

Password

Remember me

Login

Notifications

- [View](#)
- [Subscribe](#)

Indicators



[HOME](#) | [ABOUT](#) | [LOGIN](#) | [REGISTER](#) | [SEARCH](#) | [CURRENT](#) | [ARCHIVES](#) | [ARTICLE IN PRESS](#) | [ANNOUNCEMENTS](#)

Home > About the Journal > Editorial Team

Editorial Team

Editor in Chief

1. Kuswanto Kuswanto, Prof. , Faculty of Agriculture Universty of Brawijaya (Scopus ID: 57192702058), Indonesia

International Editorial Board

1. Widiatmaka Widiatmaka, Prof. , Department of Soil Science and Land Resources Bogor Agricultural University (Scopus ID: 56962708400), Indonesia
2. Jintana Unartngam, Ph.D , Department of Plant Pathology Faculty of Agriculture at Kamphaeng sane Kasertsat University (Scopus ID: 8528493500), Thailand
3. Dono Wahyuno, Dr. , Indonesian Spice and Medicinal Crops Research Institute (Scopus ID: 6507969735), Indonesia
4. Seca Gandaseca, Ph.D , Universiti Putra Malaysia, Department of Forest Production, Serdang, Malaysia (Scopus ID: 35071255800), Malaysia
5. Mohammad Reza Alizadeh, Dr. , Department of Agricultural Engineering, Rice Research Institute of Iran (RRII), Rasht, Iran (Scopus ID: 16642292100), Iran, Islamic Republic of
6. Markus Anda, Ph.D , Indonesian Centre for Agricultural Land Resource Research and Development (Scopus ID: 23024287000), Indonesia
7. Anoop Kumar Srivastava, Prof. , National Research Centre For Citrus Amravati Road, Nagpur, Maharashtra, India (Scopus ID: 56688989000), India
8. Mohammad Ali Shariati, Dr. , Orel State Agrarian University (Scopus ID: 56096567300), Russian Federation
9. Mohammad Valipour, Dr. , Islamic Azad University, Kermanshah Branch, Tehran, Iran (Scopus ID: 55877297968), Iran, Islamic Republic of
10. B. Mohan Kumar, Dr. , Nalanda University (Scopus ID: 55435104500), India
11. Hiran Anjana Ariyawansa, Ph.D , Department of Plant Pathology and Microbiology, National Taiwan University, Taiwan (Scopus ID: 55752594000), Taiwan, Province of China
12. Wani Hadi Utomo, Prof. , Land Management, Brawijaya University (Scopus ID: 6507144030), Indonesia
13. Kurniatun Hairiah, Prof. , Soil Biology, Brawijaya University (Scopus ID: 6602368888), Indonesia
14. Eko Handayanto, Prof. , Soil Organic Management, Brawijaya University, Indonesia
15. Jay Shankar Singh, Dr. , Department of Environmental Microbiology, Babasaheb Bhimrao Ambedkar University, Lucknow, India (Scopus ID: 16837178200), India

Editorial Board

1. Moch. Dawam Maghfoer, Prof. , Faculty of Agriculture Universty of Brawijaya (Scopus ID: 55440224300), Indonesia
2. Hagus Tarno, Dr. , Department of Plant Protection, University of Brawijaya (Scopus ID: 36163526900), Indonesia
3. Syahrul Kurniawan, Dr. , Faculty of Agriculture, University of Brawijaya (Scopus ID: 55876481800), Indonesia
4. Akhmad Rizali, Dr. , Department of Plant Pests and Diseases, Faculty of Agriculture, University of Brawijaya (Scopus ID: 6507320984), Indonesia
5. Darmawan Saptadi, Dr. , Faculty of Agriculture, University of Brawijaya (Scopus ID: 57193852160), Indonesia

Assistant Editor

1. Silvia Santi Wahyuni, Faculty of Agriculture University of Brawijaya, Indonesia
2. Endah Apriliani, Faculty of Agriculture University of Brawijaya, Indonesia
3. Aini Nur Laila, Faculty of Agriculture University of Brawijaya, Indonesia

About Agrivita

- Aim and Scope
- Editorial Team
- Publication Ethics
- Visitor Statistics
- Reviewer Acknowledgment

Issues

- Current Issue
- Back Issues
- Article in Press
- Accepted Papers

Information For Author

- Author Guidelines
- Template (docx)
- Template (pdf)

User

Username


Password

Remember me

Notifications

- View
- Subscribe

Indicators



Visitors

ID	29503	US	2924
IN	2114	PH	1933
MY	1581	IR	1162
TH	1127	CN	599
JP	434	RU	399
Newest:	KG You:	ID	
Today:		29	
Month:		1013	
Total:		47589	
Supercounters.com			

[HOME](#) | [ABOUT](#) | [LOGIN](#) | [REGISTER](#) | [SEARCH](#) | [CURRENT](#) | [ARCHIVES](#) | [ARTICLE IN PRESS](#) | [ANNOUNCEMENTS](#)

Home > About the Journal > People

People

Peer Reviewers

Shahabuddin Saleh, Prof. Dr. , Universits Tadulako, Palu (Scopus ID: 35335106500), Indonesia

Sun-Kee Hong, Prof. , Mokpo National University (Scopus ID: 34770558100), Korea, Republic of

Tetsuhiro Watanabe, Dr. , Kyoto University, Graduate School of Agriculture, Kyoto, Japan, Japan

Endang Arisoesilarningsih, Prof. , Brawijaya University, Indonesia

Nur Wakhid, M.Sc , Indonesian Swampland Agriculture Research Institute (Scopus ID: 57192961613), Indonesia

Edy Syahputra, Dr. , Faculty of Agriculture, Universitas Tanjungpura (Scopus ID: 55965649800), Indonesia

Rindang Dwiyani, Dr. , Udayana University, Indonesia

Neneng Laela Nurida, Dr. , Indonesian Soil Research Institute (Scopus ID : 56685751000), Indonesia

Antonius Kasno, M.Si , Indonesian Soil Research Institute, Indonesia

Fitri Rachmawati, Dr. , Indonesian Ornamental Crops Research Institute (Scopus ID: 39861797700), Indonesia

Puneet Singh Chauhan, Dr. , Division of Plant-Microbe interactions, CSIR-National Botanical Research Institute, Lucknow, 226001, India., India

Morio Tsukada, Dr. , Mie University, Graduate School of Bioresources, Tsu, Japan, Japan

Shashi Yellareddygar, Dr. , Department of Plant Pathology North Dakota State University, United States

Halil Akinci, Dr. , Department of Geomatics Engineering, Faculty of Engineering, Artvin Çoruh University (Scopus ID: ID: 26322246900), Turkey

Djajadi Djajadi, Dr. , Indonesia Research Institute for Tobacco and Fiber Crops (IToFCRI), Indonesia

Haviluddin Haviluddin, Ph.D , Fakultas Ilmu Komputer dan Teknologi Informasi, Universitas Mulawarman, Indonesia

Retno Mastuti, Dr. , Biology Department, Faculty of Mathematics and Natural Sciences, University of Brawijaya, Indonesia

Fernando B. Aurigue, Prof. , Philippine Nuclear Research Institute, Philippines

Makoto Takahashi, Ph.D , Department of Bioscience and Biotechnology, Faculty of Agriculture, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan (Scopus ID: 56287823400), Japan

Michael Henry Boehme, Prof. , Humboldt-University of Berlin Faculty of Life Sciences Dept. Horticultural Plant Systems, Germany

Robert J Holmer, Dr. , Urban Food International (UFI), Germany

Munir Ahmad, Ph.D , Department of Entomology, Faculty of Crop and Food Sciences, Pir Mehr Ali Shah, Arid Agriculture University, Pakistan

Naga Raju Maddela, Dr. , Department of Life Science, Universidad Estatal Amazónica, Santa Clara (Scopus ID: 56650913800), Ecuador

Gregory C. Luther, Ph.D , AVRDC - The World Vegetable Center, 60 Yi-min Liao, Shanhua, Tainan (Scopus ID: 8096082400), Taiwan, Province of China

Sudhanshu Shekar Behera, Dr. , Department of Biotechnology, National Institute of Technology, Raipur (Scopus ID: 56940110300), India

Anny Yanuriati, Dr. , Department of Agricultural Technology, Sriwijaya University (Scopus ID: 55749625200), Indonesia

About Agrivita

- [Aim and Scope](#)
- [Editorial Team](#)
- [Publication Ethics](#)
- [Visitor Statistics](#)
- [Reviewer Acknowledgment](#)

Issues

- [Current Issue](#)
- [Back Issues](#)
- [Article in Press](#)
- [Accepted Papers](#)

Information For Author

- [Author Guidelines](#)
- [Template \(docx\)](#)
- [Template \(pdf\)](#)

User

Username


Password

Remember me

Notifications

- [View](#)
- [Subscribe](#)

Indicators



Visitors

ID	29503	US	2924
IN	2114	PH	1933
MY	1581	IR	1162
TH	1127	CN	599
JP	434	RU	399
Newest:	KG	You:	ID
Today:			29
Month:			1013
Total:			47589
Supercounters.com			

Ragapadmi Purnamaningsih, Dr. , Indonesian Center for Agricultural Biotechnology & Genetic Resources, Indonesia

Dede J. Sudrajat, Dr. , Forest Tree Seed Technology Research Institute, Bogor, Indonesia

Eriyanto Yusnawan, Ph.D , Indonesian Legumes and Tuber Crops Research Institute, Indonesia

Bijaya Ketan Sarangi, Dr. , National Environmental Engineering Research Institute, India, India

Irawan Irawan, Dr. , Indonesian Soil Research Institute, Indonesia

Sri Wening, Dr. , Indonesian Oil Palm Research Institute (Scopus ID: 36524279800), Indonesia

Kusumawaty Kusumanegara, Dr. , Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (ICABIOGRAD), Indonesian Agency for Agricultural Research and Development (IAARD) (Scopus ID: 55331290000), Indonesia

Wan Fatma Zuharah, Dr. , School of Biological Sciences, Universiti Sains Malaysia, Malaysia

R. K. Gautam, Dr. , Central Island Agricultural Research Institute (CIARI), India

Ramadhani Eka Putra, Ph.D , School of Life Sciences and Technology, Institut Teknologi Bandung (Scopus ID: 26326479700), Indonesia

Zetty Norhana Balia Yusof, Dr. , Universiti Putra Malaysia (UPM), Malaysia

Chairani Martasari, Dr. , Indonesian Citrus and Subtropical Fruits Research Institute, Indonesia

Makruf Nurudin, Dr. , Department of Soil Science, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Reza Talebi, Dr. , Islamic Azad University, Sanandaj Branch (Scopus ID: 23010335000), Iran, Islamic Republic of

Fibrianty Minhal, M.Sc , Assessment Institute for Agricultural Technology of Yogyakarta, Indonesia

Rosyda Priyadarshini, Dr. , Faculty of Agriculture, UPN "Veteran" East Java, Indonesia

Anthon Monde, Dr. , Faculty of Agriculture, Universitas Tadulako, Indonesia

Sukartini Sukartini, Dr. , Indonesian Fruits Research Institute (IFTRI), Indonesia

Dewi Sukma, Dr. , Iant Breeding and Biotechnology Study Program, Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University (IPB) – Darmaga Campus, Bogor 16680, INDONESIA, Indonesia

Rinaldi Sjahril, Ph.D , Hasanuddin University, Department of Agronomy, Makassar, Indonesia, Indonesia

Sumaryono Sumaryono, M.Sc , Indonesian Research Institute for Biotechnology and Bioindustry, Indonesia

IGP Muliarta Aryana, Prof. , Faculty of Agriculture, University of Mataram Mataram, Lombok, NTB, Indonesia, Indonesia

Markus Anda, Ph.D , Indonesian Centre for Agricultural Land Resource Research and Development (Scopus ID: 23024287000), Indonesia

Nur Edy, Ph.D , Faculty of Agriculture, Tadulako University, Indonesia

Mohammad Eslami, Dr. , RWTH Aachen University: Aachen, DE, Germany

Astanto Kasno, Prof. , Indonesian Legume and Tuber Crops Research Institute, Indonesia

Kolluru Viswanatha Chaitanya, Dr. , Department of Biotechnology, Gandhi Institute of Technology and Management, Visakhapatnam, Visakhapatnam, India (Scopus ID: 6603655335), India

Kavitha Mary Jackson, Ph.D , Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore - 641 003, Tamil Nadu, India, India

P. Veeramani, Ph.D , Research Associate, Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore- 641 003, Tamil Nadu, India (Scopus ID: 54390279800), India

Lia Hapsari, M.Si , Purwodadi Botanic Garden - Indonesian Institution of Sciences (Scopus ID: 53873774600), Indonesia

Franciscus S. Rembon, Ph.D , Faculty of Agriculture, Universitas Haluoleo Jalan H.E.A Mokodompit No. 1 Anduonohu Kota Kendari, 93232, Indonesia (Scopus ID: 6504338021), Indonesia

Fadly Hairannoor Yusran, Prof. , Lambung Mangkurat University, Indonesia, Indonesia

Dita Agisimanto, Dr. , Balitjestro

Jay Shankar Singh, Dr. , Department of Environmental Microbiology, Babasaheb Bhimrao Ambedkar University, Lucknow, India (Scopus ID: 16837178200), India

Pham Van Du, Dr. , University of Science, Ho Chi Minh City Vietnam, Viet Nam

Keywords

2AP Agroforestry Biological control
Chrysanthemum DNA extraction
Morphology Oryza sativa REDD+ Rice
SSR markers corn **diversity** drought
genotype land productivity maize mungbean
population **rice** soil fertility soybean

Journal Content

Search

Search Scope **All** ▼

Browse

- By Issue
- By Author
- By Title

Information

- For Readers
- For Authors
- For Librarians

Journal Help

Yu Ya Aye, Dr. , Forest Research Institute, Forest Department Ministry of Environmental Conservation and Forestry Naypyitaw Myanmar, Myanmar

Ludji Pantja Astuti, Brawijaya University, Indonesia

Aji Prasetya Wibawa, Dr.

Roman Voliansky, Dr. , Dniprovsk State Technical University, Ukraine

Andri Pranolo, M.Cs, Universitas Ahmad Dahlan, Indonesia

Hakim Kurniawan, MP. , Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (Scopus ID: 35305429800), Indonesia

Dindin Hidayatul Mursyidin, Dr. , Faculty of Mathematics and Natural Sciences, Universitas Lambung Mangkurat (Scopus ID: 57190937708), Indonesia

Abdul Razak Purba, Dr. , Indonesian Oil Palm Research Institute (IOPRI), Medan, Indonesia (Scopus ID: 6507981002), Indonesia

Chindy Ulina Zanetta, MP. , School of Life Sciences and Technology - ITB, Indonesia

Dedi Ruswandi, Dr. , Department of Agrotechnology, Universitas Padjadjaran (Scopus ID: 56256211800), Indonesia

Abd El-Rahman Abd El-Raouf Ahmed, Prof. Dr. Agric. Eng. , Agricultural Engineering Research Institute, Egypt

Viet The Ho, Dr. , Ho Chi Minh City University of Food Industry.140 Le Trong Tan Street, Tan Phu district, Ho Chi Minh City,Vietnam (Scopus ID: 55318292200), Viet Nam

Vermando M. Aquino, Prof. Dr. , National Institute of Molecular Biology and Biotechnology National Science Complex College of Science University of the Philippines Diliman Quezon City, Philippines 1101, Philippines

Ravindra C. Joshi, Prof. Dr. , Pampanga State Agricultural University, Philippines (Scopus ID: 7202085232), Philippines

Supriadi Supriadi, Prof. Dr. , Indonesian Spice and Medicinal Crops Research Institute (ISMCR), Indonesia

Hardian Susilo Addy, Ph.D , Faculty of Agriculture, Universitas Jember, Indonesia

Joseph Adjei Adjetej, Dr , Botswana University of Agriculture & Natural Resources, Botswana

Eko Hanudin, Dr. , Faculty of Agriculture, Gadjah Mada University, Indonesia

Dedik Budianta, Prof. , Soil Science Department, Faculty of Agriculture, Sriwijaya University, Indonesia

Meredith P. Martin, Dr. , Yale School of Forestry & Environmental Studies, USA, United States

Puji Lestari, Ph.D , Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (Scopus ID: 6507413576), Indonesia

Arnoldus Klau Berek, Ph.D , Fakultas Pertanian Universitas Timor, Indonesia

Tri Puji Priyatno, Dr. , Balai Besar Penelitian dan Pengembangan Bioteknologi dan Sumber daya Genetik Pertanian

Hadiwiyono Hadiwiyono, Prof. , Faculty of Agriculture, Universitas Sebelas Maret, Indonesia

Cicik Udayana, M.Si , Department of Applied Ecology, Inland Norway University of Applied Sciences, Norway

Yogo - Setiawan, Mr. , University of Brawijaya, Indonesia

Joko Susilo Utomo, Dr. , Indonesian Legumes and Tuber Crops Research Institute (Scopus ID: 37069591500), Indonesia

Dilson Antonio Bisognin, Dr. , Universidade Federal de Santa Maria, Departamento de Fitotecnia, Santa Maria, Brazil, Brazil

Suntoro Suntoro, Prof. Dr. , Department of Soil Science, Faculty of Agriculture, Universitas Sebelas Maret, Surakarta (Scopus ID: 57199185618), Indonesia

Nono Carsono, Ph.D , Universitas Padjadjaran, Indonesia

Dono Wahyuno, Dr. , Indonesian Spice and Medicinal Crops Research Institute (Scopus ID: 6507969735), Indonesia

Darmawan Saptadi, Dr. , Faculty of Agriculture, University of Brawijaya (Scopus ID: 57193852160), Indonesia

Edi Santosa, Prof. Dr. , Department of Agronomy and Horticulture, Bogor Agricultural University (Scopus ID: 8790748700), Indonesia

Darunee Jothityangkoon, Ph.D , Department of Plant Science and Agricultural Resources, Faculty of Agriculture, Khon Kaen University, Khon Kaen 40002, Thailand (Scopus ID: 43461240900), Thailand

Kukuh Murtiaksono, Prof. Dr. , Department of Soil Sciences and Land Resources, Faculty of Agriculture, Bogor Agricultural University (Scopus ID: 56469817600), Indonesia

Shu-Dan Yeh, Dr. , Department of Life Science, National Central University (Scopus ID: 54997267200), Taiwan, Province of China

Tjandra Anggraeni, Dr. , School of Life Sciences and Technology, Institut Teknologi Bandung (Scopus ID: 24358584300), Indonesia

Heni Purnamawati, Dr. , Faculty of Agriculture, Bogor Agricultural University (Scopus ID: 57193848936), Indonesia

Kanokwan Teingtham, M.S. , Kasetsart University, Thailand

Mochtar Lutfi Rayes, Prof. , Soil Science Department, Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 56997327000), Indonesia

Saptowo Jumali Pardal, Dr. , Indonesian Center for Agricultural Biotechnology & Genetic Resources, Indonesia

Welhelmus I. I. Mella, Dr. , Faculty of Agriculture, University of Nusa Cendana (Scopus ID: 35305539200), Indonesia

Ika Roostika Tambunan, Dr. , Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (ICABIOGRAD), Indonesia

Syahrul Kurniawan, Dr. , Faculty of Agriculture, University of Brawijaya (Scopus ID: 55876481800), Indonesia

Reflinur Reflinur, Dr. , Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development, Bogor, Indonesia

Akhmad Rizali, Dr. , Department of Plant Pests and Diseases, Faculty of Agriculture, University of Brawijaya (Scopus ID: 6507320984), Indonesia

Luqman Qurata Aini, Ph.D , Department of Pest and Plant Disease, Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 35071053600), Indonesia

Laode Muhammad Harjoni Kilowasid, Dr. , Department of Agrotechnology, Faculty of Agriculture, Universitas Halu Oleo (Scopus ID: 56013958300), Indonesia

Marulak Simarmata, Ph.D. , Department of Agronomy, Bengkulu University (Scopus ID: 13403042100), Indonesia

Setyo Dwi Utomo, Prof. , Department of Agronomy and Horticulture, Faculty of Agriculture, University of Lampung (Scopus ID: 35249687600), Indonesia

Bambang Guritno, Prof. , Faculty of Agriculture, Universitas Brawijaya, Indonesia (Scopus ID: 55323950300), Indonesia

Riza Arief Putranto, Dr. , Indonesian Research Institute for Biotechnology and Bioindustry (Scopus ID: 54789964100), Indonesia

Himalaya Subedi, M. Sc. Agr , Resource Identification and Management Society, Dhadging, Nepal

Asmarhansyah Asmarhansyah, Ph.D , Agency for Agricultural Research and Development, Indonesia

Kevin Darras, Dr. , Department of Crop Sciences, University of Goettingen (Scopus ID: 53979686700), Germany

Jintana Unartngam, Ph.D , Department of Plant Pathology Faculty of Agriculture at Kamphaeng sane Kasertsat University (Scopus ID: 8528493500), Thailand

Hakan Sevik, Dr. , Landscape Architecture Department, Faculty of Engineering and Architecture, Kastamonu University (Scopus ID: 36633291300), Turkey

Nikardi Gunadi, Dr. , Indonesian Vegetable Research Institute (Scopus ID: 7801567795), Indonesia

Kyaw Min Htut, Ph.D , Central Agriculture Research and Training Center Myanmar (Scopus ID: 57192314425), Myanmar

Yayan Sanjaya, Dr. , Universitas Pendidikan Indonesia (Scopus ID: 55865083900), Indonesia

Khalid Abdallah Hussein, Ph.D , Botany and Microbiology Department, Faculty of Science, Assiut University (Scopus ID: 36938115000), Egypt

Md. Zaherul Islam, Ph.D , Bangladesh Agricultural Research Institute (Scopus ID: 56497388200), Bangladesh

K. V. Mohanan, Prof. , Genetics and Plant Breeding Division, Department of Botany, University of Calicut (Scopus ID: 56625173400), India

Yasushi Mitsuda, Ph.D , Faculty of Agriculture, University of Miyazaki (Scopus ID: 12798575600), Japan

Fumiaki Kitahara, Ph.D , Forestry and Forest Products Research Institute, Shikoku Research Center, Tsukuba (Scopus ID: 57193820237), Japan

Shamsuddin Jusop, Prof. , Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia (Scopus ID: 6601937415), Malaysia

Leo Mualim, Dr. , Bogor Agricultural University, Indonesia

Jianguo Wang, Prof. , Department of Plant Protection, Jiangxi Agricultural University (Scopus ID: 51666293100), China

Lita Soetopo, Prof. , Faculty of Agriculture, Universitas Brawijaya, Indonesia

Ellis Nihayati, Dr. , Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 55763928600), Indonesia

Slamet Minardi, Prof. , Faculty of Agriculture, Universitas Sebelas Maret, Indonesia

Catur Hermanto, Dr. , Agency for Agricultural Research and Development (Scopus ID: 51863728900), Indonesia

Tjintokohadi Tjintokohadi, M.Sc. , International Potato Centre (CIP) Regional Office for East South East Asia and The Pacific, Indonesia

Arifin Noor Sugiharto, Ph.D , Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 6507826962), Indonesia

Amran Muis, Dr. , Indonesian Cereals Research Institute, Indonesia

Suhartiningsih Dwi Nurcahyanti, Dr. , Faculty of Agriculture, Universitas Jember (Scopus ID: 55305711100), Indonesia

Dwi Guntoro, Dr. , Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University, Indonesia

Satoto Satoto, Dr. , Indonesian Center for Rice Research (Scopus ID: 56719665200), Indonesia

Heru Kuswantoro, Dr , Indonesian Legume and Tuber Crops Research Institute (Scopus ID: 53866614600), Indonesia

Sudir Sudir, Ir. , Indonesian Center for Rice Research, Indonesia

Hugo Volkaert, Dr. , Center for Agricultural Biotechnology, Kasetsart University (Scopus ID: 14827498100), Thailand

Widodo Widodo, Ph.D , Department of Biology, Universitas Brawijaya (Scopus ID: 57190247745), Indonesia

Thunya Taychasinpitak, Dr. , Department of Horticulture, Faculty Agriculture, Kasetsart University (Scopus ID: 55402494100), Thailand

Loekas Soesanto, Prof. , Department of Plant Pests and Diseases, Faculty of Agriculture, Jenderal Soedirman University (Scopus ID: 55763839700), Indonesia

Vrushali Pashte, Ph.D , Department of Rural Technology, Guru Ghasidas University, India

Zainuddin Basri, Prof. , Faculty of Agriculture, Tadulako University (Scopus ID: 15051504600), Indonesia

Masruri Masruri, Ph.D , Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya (Scopus ID: 55387384800), Indonesia

Bambang Pujiasmanto, Prof. , Faculty Agriculture, Sebelas Maret University (Scopus ID: 57195305824), Indonesia

Thomas Wijaya, Dr. , Indonesian Rubber Research Institute (Scopus ID: 56527010600), Indonesia

I Nengah Suwastika, Dr. Lif. Sc. , Biology Department, Faculty of Sciences, Universitas Tadulako (Scopus ID: 6507366174), Indonesia

Rehenuma Tabassum, Dr. , Department of Crop Botany and Tea Production Technology, Sylhet Agricultural University (Scopus ID: 56913124700), Bangladesh

Glen Lee Hartman, Prof. , Department of Crop Science, Illinois University (Scopus ID: 36622876800), United States

Ifar Subagiyo, Prof. , Faculty of Animal Husbandry, Universitas Brawijaya, Indonesia

Ridho Kurniati, Dr. , Indonesian Ornamental Crops Research Institute (Scopus ID: 57199653286), Indonesia

Budi Winarto, Dr. , Indonesian Ornamental Crops Research Institute (Scopus ID: 36509217200), Indonesia

Mahluga M. Yusifova, Dr. , Faculty of Ecology and Soil Sciences, Baku State University, Azerbaijan, Azerbaijan

Wolfgang Grüneberg, Ph.D , International Potato Center (CIP) Peru (Scopus ID: 6602602473), Peru

Meine van Noordwijk, Prof. , World Agroforestry Centre (ICRAF) (Scopus ID: 7003897764), Indonesia

Akira Kikuchi, Ph.D , Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 55516610400), Indonesia

Sholihin Halim, Dr. , Indonesian Legumes and Tuber Crops Research Institute (Scopus ID: 56103558900), Indonesia

Laurent Barbiero, Dr. , Institut de Recherche pour le Developpement (Institute of Research for Development/IRD) (Scopus ID: 6701542182), France

Tri Joko, Dr. , Faculty of Agriculture, Universitas Gadjah Mada (Scopus ID: 17345721200), Indonesia

Teo Chee How, Prof. , Taylor's University, Kuala Lumpur, Malaysia

A. Mujib, Ph.D , Department of Botany, Faculty of Science, Jamia Hamdard University (Scopus ID: 13407929400), India

Rina Devnita, Dr. , Department of Soil Science, Faculty of Agriculture University of Padjadjaran, Indonesia

Ahmad Faizal, Dr. , School of Life Sciences & Technology, Institut Teknologi Bandung (Scopus ID: 36727358600), Indonesia

Budi Waluyo, Dr. , Plant Breeding Lab., Department of Agronomy Faculty of Agriculture, Brawijaya University (Scopus ID: 56605006300), Indonesia

Slamet Rohadi Suparto, M.Agr. St , Faculty of Agriculture, University of Jenderal Soedirman (Scopus ID: 56052473800), Indonesia

Jiban Shrestha, M.Sc , Nepal Agricultural Research Council, Nepal, Nepal

Siti Zaharah Sakimin, Dr. , Department of Crop Sciences, Universiti Putra Malaysia (Scopus ID: 57190195119), Malaysia

Tatik Wardiyati, Prof. , Faculty of Agriculture, Universitas Brawijaya (Scopus ID: 55440256600), Indonesia

Yuuko Takeuchi, Dr. , Laboratory of Microbial Ecology, Faculty of Agriculture, Kyoto University (Scopus ID: 10044030300), Japan

Ferisman Tindaon, Prof. , Nommensen University, Medan (Scopus ID: 55323181300), Indonesia

Sudarsono Sudarsono, Prof. , Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University (Scopus ID: 16450098400), Indonesia

Agus Sutanto, Dr. , Indonesian Tropical Fruit Research Institute (Scopus ID: 51864702700), Indonesia

Mustaid Siregar, M.Si , Center for Plant Conservation Botanic Gardens (Bogor Botanic Gardens), Indonesian Institute of Sciences (Scopus ID: 6506442081), Indonesia

Murni Dwiati, Dr. , Faculty of Biology University of Jenderal Soedirman, Indonesia

Noladhi Wicaksana, Ph.D , Faculty of Agriculture, Universitas Padjadjaran (Scopus ID: 50362316600), Indonesia

Yosep Seran Mau, Ph.D , Faculty of Agriculture, University of Nusa Cendana (Scopus ID: 13105825100), Indonesia

Mesude Unal, Dr. , Kocaeli University (Scopus ID: 57197197928), Turkey

Farzane Porianejad, Dr. , Ministry of Agriculture (Scopus ID: 56527274200), Iran, Islamic Republic of

Sujin Patarapuwadol, Dr. , Faculty of Agriculture, University of Kasetsart (Scopus ID: 6504368181), Thailand

Magdi A. El Sayed, Ph.D , Department of Botany, Faculty of Science, Aswan University (Scopus ID: 7202416231), Egypt

Gun Mardiatmoko, Dr. , Forestry Department, Faculty of Agriculture, Pattimura University (Scopus ID: 57194205415), Indonesia

Catur Herison, Dr. , Department of Agronomy, Bengkulu University (Scopus ID: 56527196000), Indonesia

Untung Susanto, Dr. , Indonesian Center for Rice Research (ICRR) of Indonesian Agency for Agricultural Research and Development (IAARD) of the Ministry of Agriculture (Scopus ID: 55971074300), Indonesia

Sandy Loh, Prof. , School of Biosciences, University of Nottingham Malaysia Campus (Scopus ID: 18434637700), Malaysia

Totok Agung, Prof. , Faculty of Agriculture, Jenderal Soedirman University (Scopus ID: 6506751630), Indonesia

Muhamad Syukur, Dr. , Department of Agronomy and Horticulture, Bogor Agricultural University (Scopus ID: 56033993500), Indonesia

A Wihardjaka, M.Si , Indonesian Agricultural Environment Research Institute (Scopus ID: 6505981610), Indonesia

Eko Handayanto, Prof. , Faculty of Agriculture, Brawijaya University (Scopus ID: 6508116842), Indonesia

Jamsari Jamsari, Prof. , Faculty of Agriculture, University of Andalas (Scopus ID: 55790904900), Indonesia

Irawati Irawati, Dr. , Indonesian Institute of Science, Indonesia

Yuni Widyastuti, SP. , Indonesian Center for Rice Research (Scopus ID: 56719603200), Indonesia

B. Mohan Kumar, Dr. , Nalanda University (Scopus ID: 55435104500), India

Hidegori Tanaka, Dr. , Faculty of Agriculture, University of Miyazaki (Scopus ID: 55514958400), Japan

Koki Homma, Dr. , Crop Science Laboratory, University of Kyoto (Scopus ID: 7103360331), Japan

Wang Pu, Dr. , College of Agronomy and Biotechnology, China Agricultural University (Scopus ID: 24175198400), China

Dennis B. Egli, Prof. , Plant and Soil Sciences Department, University of Kentucky (Scopus ID: 7003399538), United States

Xinbin Feng, Prof. , Institute of Geochemistry Chinese Academy of Science (Scopus ID: 27868165200), China

Pablo Tittone, Prof. , Wageningen University (Scopus ID: 14520144600), Netherlands

Husnain Husnain, Ph.D. , Indonesian Soil Research Institute (Scopus ID: 25922548100), Indonesia

Denah Suswati, Dr. , Department of Soil Science, Tanjungpura University (Scopus ID: 56888802100), Indonesia

Katsumi Yamaguchi, Dr. , Department of Biotechnology and Life Science, Tokyo University of Agriculture and Technology (Scopus ID: 7405930314), Japan

Oteng Haridjaja, Dr. , Faculty of Agriculture, Bogor Agricultural University (Scopus ID: 56888681600), Indonesia

Masae Iwamoto Ishihara, Dr. , Graduate School for International Development and Cooperation, Hiroshima University (Scopus ID: 37015427300), Japan

Yusnita Yusnita, Prof. , Department of Agrotechnology, University of Lampung (Scopus ID: 55552150100), Indonesia

Ayman Esh, Ph.D. , Sugar Crops Research Institute, Agricultural Research Center (Scopus ID: 55970795600), Egypt

Triwidodo Arwiyanto, Prof. , Faculty of Agriculture, University of Gadjah Mada (Scopus ID: 54892106600), Indonesia

Tualar Simarmata, Prof. , Faculty of Agriculture, University of Padjajaran (Scopus ID: 6505785217), Indonesia

Patcharin Tanya, Ph.D. , Department of Agronomy, Faculty of Agriculture, Kasetsart University (Scopus ID: 11539498100), Thailand

Hagus Tarno, Dr. , Department of Plant Protection, University of Brawijaya (Scopus ID: 36163526900), Indonesia

Rikiya Endoh, Dr. , Microbe Division, Japan Collection of Microorganisms, RIKEN BioResource Center (Scopus ID: 24331184300), Japan

Mirza Hasanuzzaman, Prof. Dr. , Department of Agronomy, Shere-e Bagla Agricultural University (Scopus ID: 26030936600), Bangladesh

Syekhfani Syekhfani, Prof. , Department of Soil Science, Universitas Brawijaya (Scopus ID: 56272862600), Indonesia

Didi Ardi Suriadikarta, Prof. Dr. , Indonesian Soil Research Institute, Indonesia

Panca Jarot Santoso, Dr. , Indonesian Tropical Fruit Research Institute (Scopus ID: 57191268880), Indonesia

Affandi Affandi, Ph.D. , Indonesian Tropical Fruits Research Institute (Scopus ID: 55981822000), Indonesia

Nunun Barunawati, Dr. , Department of Agronomy, Universitas Brawijaya (Scopus ID: 56166205100), Indonesia

Moch. Dawam Maghfoer, Prof. , Faculty of Agriculture University of Brawijaya (Scopus ID: 55440224300), Indonesia

Budi Prasetya, Dr. , Faculty of Agriculture University of Brawijaya (Scopus ID: 57009851100), Indonesia

Kuswanto Kuswanto, Prof. , Faculty of Agriculture University of Brawijaya (Scopus ID: 57192702058), Indonesia

Kurniawan Budiarto, Ph.D. , Indonesian Ornamental Crops Research Institute (Scopus ID: 12243097600), Indonesia

Akhbar Akhbar, Dr. , University of Tadulako (Scopus ID: 56527212600), Indonesia

Widiatmaka Widiatmaka, Prof. , Department of Soil Science and Land Resources Bogor Agricultural University (Scopus ID: 56962708400), Indonesia

Mohammad Zahirul Islam, Ph.D. , Kangwon National University (Scopus ID: 56074653400), Korea, Republic of

[HOME](#) | [ABOUT](#) | [LOGIN](#) | [REGISTER](#) | [SEARCH](#) | [CURRENT](#) | [ARCHIVES](#) | [ARTICLE IN PRESS](#) | [ANNOUNCEMENTS](#)

Home > Archives > Vol 42, No 1 (2020)

Vol 42, No 1 (2020)

(IN PRESS)

DOI: <http://doi.org/10.17503/agrivita.v42i1>

Table of Contents

Articles

	Salinity-Induced Changes in the Nutritional Quality of Bread Wheat (<i>Triticum aestivum</i> L.) Genotypes <i>Muhammad Nadeem, Muhammad Nouman Tariq, Muhammad Amjad, Muhammad Sajjad, Muhammad Akram, Muhammad Imran, Mohammad Ali Shariati, Tanweer Aslam Gondal, Nadezhda Kenjiz, Dmitriy Kulikov</i>		PDF	Viewed: 70 times 
	Comparative Study of Nano-chitosan and Synthetic Bactericide Application on Chili Pepper (<i>Capsicum annuum</i> L.) Infected by <i>Xanthomonas campestris</i> <i>Rizkita R. Esyanti, Nadya Farah, Brahmani D. Bajra, Diah Nofitasari, Ronny Martien, Sunardi Sunardi, Ramadhani Safitri</i>		PDF	Viewed: 27 times 
	Tomato Growth Stage Monitoring for Smart Farm Using Deep Transfer Learning with Machine Learning-based Maturity Grading <i>Robert G. de Luna, Elmer P. Dadios, Argel A. Bandala, Ryan Rhay P. Vicerra</i>		PDF	Viewed: 56 times 
	Efficacy of Selected Plant Extracts to Control Leaf Miner (<i>Lyriomyza</i> spp.) in Chrysanthemum <i>Indijarto Budi Rahardjo, Budi Marwoto, Kurniawan Budiarto</i>		PDF	Viewed: 25 times 
	<i>Kaempferia galanga</i> L. A Medicinal Plant Used By Indonesian Ethnic Groups: Genetic Diversity Based on Inter-Simple Sequence Repeats (ISSR) <i>Dyah Subositi, Nina Kurnianingrum, Rohmat Mujahid, Yuli Widiyastuti</i>		PDF	Viewed: 28 times 
	Determination of Agronomical Characters as the Resistance Attributes of Twenty Soybean Varieties to Stink Bug (<i>Nezara viridula</i> L.) <i>Heru Kuswantoro, I Made Jana Mejaya, Yuliantoro Baliadi</i>		PDF	Viewed: 39 times 
	Direct Seed Planting System and Giving Liquid Organic Fertilizer as A New Method to Increase Rice Yield and Growth (<i>Oryza sativa</i> L.) <i>Mohamad Ikbal Bahua, Hayatiningsih Gubali</i>		PDF	Viewed: 26 times 
	Turning Volcanic Ash into Fertile Soil: Farmers' Options in Coffee Agroforestry After the 2014 Mount Kelud Eruption <i>Rizki Maulana Ishaq, Danny Dwi Saputra, Rika Ratna Sari, Didik Suprayogo, Widiyanto Widiyanto, Cahyo Prayogo, Kurniatun Hairiah</i>		PDF	Viewed: 11 times 
	Genetic Potential of Agronomic Traits and Seed Physicochemical Properties of Novel <i>Jatropha</i> Lines Developed from Complex Interspecific Hybridization <i>Narathid Muakrong, Peerasak Srinives</i>		PDF	Viewed: 9 times 
	Effect of Vermicompost and Nitrogen on N, K, Na Uptakes and Growth of Sugarcane in Saline Soil <i>Djajadi Djajadi, Roni Syaputra, Sulis Nur Hidayati, Yaumil Khairiyah</i>		PDF	Viewed: 16 times 

About Agrivita

- Aim and Scope
- Editorial Team
- Publication Ethics
- Visitor Statistics
- Reviewer Acknowledgment

Issues

- Current Issue
- Back Issues
- Article in Press
- Accepted Papers

Information For Author

- Author Guidelines
- Template (docx)
- Template (pdf)

User

Username


Password

Remember me

Notifications

- View
- Subscribe

Indicators



Visitors

ID	29503	US	2924
IN	2114	PH	1933
MY	1581	IR	1162
TH	1127	CN	599
JP	434	RU	399
Newest:	KG	You:	ID
Today:			29
Month:			1013
Total:			47589
Supercounters.com			

Direct Seed Planting System and Giving Liquid Organic Fertilizer as A New Method to Increase Rice Yield and Growth (*Oryza sativa* L.)

Mohamad Ikbal Bahua, Hayatiningsih Gubali

Abstract

The transplanting system is a common method in rice cultivation, whereas the direct seed planting system combined with the application of organic fertilizer is rarely applied. This study aimed to determine the impact both the direct seed planting system and the application of liquid organic fertilizer the growth and production of rice. The research used the factorial randomized block design, direct seed planting system tested by the transplanting system and application of liquid organic fertilizer doses 25, 35, and 45 l/ha. The observed variables were plant height, number of tillers, productive tillers, panicle length, weight of 1000 rice grains, and grain yield of plots. Data were analyzed by analysis of variance with 5% LSD test. The results revealed that direct seed planting system and application of liquid organic fertilizer impact the growth and yield of rice. The application of liquid organic fertilizer a dose of 45 l/ha gave the highest influence among all observed variables. There is a correlation between the treatment of the planting system and the application of liquid organic fertilizer on the growth of the number of tillers and productive tillers. The direct seed planting system increases the yield of rice more than the transplanting system.

Keywords

Direct seed planting; Liquid organic fertilizers; Transplanting; Rice








Full Text:

PDF

References

- Andres, A., Fogliatto, S., Ferrero, A., & Vidotto, F. (2015). Growth variability of Italian weedy rice populations grown with or without cultivated rice. *Crop Science*, 55(1), 394–402. <https://doi.org/10.2135/cropsci2013.07.0472>
- Birhane, A. (2013). Effect of planting methods on yield and yield components of rice (*Oryza sativa* L.) varieties in Tahtay Koraro Wereda, Northern Ethiopia. *International Journal of Technology Enhancements and Emerging Engineering Research*, 1(5), 1–5. Retrieved from <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.407.6481&rep=rep1&type=pdf>
- Colla, G., Roupael, Y., Canaguier, R., Svecova, E., & Cardarelli, M. (2014). Biostimulant action of a plant-derived protein hydrolysate produced through enzymatic hydrolysis. *Frontiers in Plant Science*, 5, 448. <https://doi.org/10.3389/fpls.2014.00448>
- Dimkpa, C. O., & Bindraban, P. S. (2016). Fortification of micronutrients for efficient agronomic production: a review. *Agronomy for Sustainable Development*, 36, 7. <https://doi.org/10.1007/s13593-015-0346-6>
- Douglas, J. A., & Dyson, C. B. (1985). Preliminary investigation of concentrations of minerals and nitrogen in wheat grain, and their relationship with baking quality and grain weight. *New Zealand Journal of Agricultural Research*, 28(1), 81–85. <https://doi.org/10.1080/00288233.1985.10426999>
- Fahrurrozi, Muktamar, Z., Setyowati, N., Sudjatmiko, S., & Chozin, M. (2019). Comparative effects of soil and foliar applications of Tithonia-enriched liquid organic fertilizer on yields of sweet corn in closed agriculture production system. *AGRIVITA Journal of Agricultural Science*, 41(2), 238–245. <https://doi.org/10.17503/agrivita.v41i2.1256>
- Gopal Ramdas, M., Manjunath, B. L., Narendra Pratap, S., Ramesh, R., Verma, R. R., Marutrao, L. A., ... Rahul, K. (2017). Effect of organic and inorganic sources of nutrients on soil microbial activity and soil organic carbon build-up under rice in west coast of India. *Archives of Agronomy and Soil Science*, 63(3), 414–426. <https://doi.org/10.1080/03650340.2016.1213813>
- Javaid, T., Awan, I. U., Baloch, M. S., Shah, I. H., Nadim, M. A., Khan, E. A., ... Abuzar, M. R. (2012). Effect of planting methods on the growth and yield of coarse rice. *Journal of Animal and Plant Sciences*, 22(2), 358–362. Retrieved from <http://www.thejaps.org.pk/docs/v-22-2/20.pdf>
- Jones, D. L., Cross, P., Withers, P. J. A., DeLuca, T. H., Robinson, D. A., Quilliam, R. S., ... Edwards-Jones, G. (2013). REVIEW: Nutrient stripping: the global disparity between food security and soil nutrient stocks. *Journal of Applied Ecology*, 50(4), 851–862. <https://doi.org/10.1111/1365-2664.12089>

Article Tools

-  Print this article
-  Indexing metadata
-  How to cite item
-  Supplementary files
-  Finding References
-  Email this article (Login required)
-  Email the author (Login required)

About Agrivita

- [Aim and Scope](#)
- [Editorial Team](#)
- [Publication Ethics](#)
- [Visitor Statistics](#)
- [Reviewer Acknowledgment](#)

Issues

- [Current Issue](#)
- [Back Issues](#)
- [Article in Press](#)
- [Accepted Papers](#)

Information For Author

- [Author Guidelines](#)
- [Template \(docx\)](#)
- [Template \(pdf\)](#)

User

Username


Password

Remember me

Notifications

- [View](#)
- [Subscribe](#)

Indicators



Kartika, Lakitan, B., Sanjaya, N., Wijaya, A., Kadir, S., Kurnianingsih, A., ... Meihana, M. (2018). Internal versus edge row comparison in jajar legowo 4:1 rice planting pattern at different frequency of fertilizer applications. *AGRIVITA Journal of Agricultural Science*, 40(2), 222–232. <https://doi.org/10.17503/agrivita.v40i0.1715>

Lal, B., Gautam, P., Nayak, A. K., Maharana, S., Tripathi, R., Shahid, M., ... Singh, S. (2020). Tolerant varieties and exogenous application of nutrients can effectively manage drought stress in rice. *Archives of Agronomy and Soil Science*, 66(1), 13–32. <https://doi.org/10.1080/03650340.2019.1587749>

Leogrande, R., Vitti, C., Stellacci, A. M., Coccozza, C., & Ventrella, D. (2016). Response of wheat crop during transition to organic system under Mediterranean conditions. *International Journal of Plant Production*, 10(4), 565–578. <https://doi.org/10.22069/ijpp.2016.3050>

Martínez-Alcántara, B., Martínez-Cuenca, M. R., Bermejo, A., Legaz, F., & Quiñones, A. (2016). Liquid organic fertilizers for sustainable agriculture: Nutrient uptake of organic versus mineral fertilizers in citrus trees. *PLoS ONE*, 11(10), e0161619. <https://doi.org/10.1371/journal.pone.0161619>

Moreno, J. L., Ondoño, S., Torres, I., & Bastida, F. (2017). Compost, Leonardite, and zeolite impacts on soil microbial community under barley crops. *Journal of Soil Science and Plant Nutrition*, 17(1), 214–230. <https://doi.org/10.4067/S0718-95162017005000017>

Qiao, J., Liu, Z., Deng, S., Ning, H., Yang, X., Lin, Z., ... Ding, Y. (2011). Occurrence of perfect and imperfect grains of six japonica rice cultivars as affected by nitrogen fertilization. *Plant and Soil*, 349(1–2), 191–202. <https://doi.org/10.1007/s11104-011-0861-4>

Richards, R. A. (2008). Genetic opportunities to improve cereal root systems for dryland agriculture. *Plant Production Science*, 11(1), 12–16. <https://doi.org/10.1626/pp.11.12>

Santosa, M., & Suryanto, A. (2015). The growth and yield of paddy Ciherang planted in dry and wet season and fertilized with organic and inorganic fertilizers. *AGRIVITA Journal of Agricultural Science*, 37(1), 24–29. <https://doi.org/10.17503/agrivita-2015-37-1-p024-029>

Shah-Al-Emran, Krupnik, T. J., Kumar, V., Ali, M. Y., & Pittelkow, C. M. (2019). Agronomic, economic, and environmental performance of nitrogen rates and source in Bangladesh's coastal rice agroecosystems. *Field Crops Research*, 241, 107567. <https://doi.org/10.1016/j.fcr.2019.107567>

Sharma, G., Giri, J., & Tyagi, A. K. (2016). Sub-functionalization in rice gene families with regulatory roles in abiotic stress responses. *Critical Reviews in Plant Sciences*, 35(4), 231–285. <https://doi.org/10.1080/07352689.2016.1265357>

Smith, S. F., Brye, K. R., Gbur, E. E., Chen, P., & Korth, K. (2014). Residue and water management effects on aggregate stability and aggregate-associated carbon and nitrogen in a wheat–soybean, double-crop system. *Soil Science Society of America Journal*, 78, 1378–1391. <https://doi.org/10.2136/sssaj2013.12.0534>

Thakur, A. K., & Uphoff, N. T. (2017). How the system of rice intensification can contribute to climate-smart agriculture. *Agronomy Journal*, 109(4), 1163–1182. <https://doi.org/10.2134/agronj2016.03.0162>

Tun, T. N., Guo, J., Fang, S., & Tian, Y. (2018). Planting spacing affects canopy structure, biomass production and stem roundness in poplar plantations. *Scandinavian Journal of Forest Research*, 33(5), 464–474. <https://doi.org/10.1080/02827581.2018.1457711>

Wang, W., Chen, Q., Hussain, S., Mei, J., Dong, H., Peng, S., ... Nie, L. (2016). Pre-sowing seed treatments in direct-seeded early rice: Consequences for emergence, seedling growth and associated metabolic events under chilling stress. *Scientific Reports*, 6, 19637. <https://doi.org/10.1038/srep19637>

Xiao, M., Zang, H., Ge, T., Chen, A., Zhu, Z., Zhou, P., ... Kuzyakov, Y. (2019). Effect of nitrogen fertilizer on rice photosynthate allocation and carbon input in paddy soil. *European Journal of Soil Science*, 70(4), 786–795. <https://doi.org/10.1111/ejss.12811>

Yoshinaga, S., Takai, T., Arai-Sanoh, Y., Ishimaru, T., & Kondo, M. (2013). Varietal differences in sink production and grain-filling ability in recently developed high-yielding rice (*Oryza sativa* L.) varieties in Japan. *Field Crops Research*, 150, 74–82. <https://doi.org/10.1016/j.fcr.2013.06.004>

Zandvakili, O. R., Barker, A. V., Hashemi, M., Etemadi, F., & Autio, W. R. (2019). Comparisons of commercial organic and chemical fertilizer solutions on growth and composition of lettuce. *Journal of Plant Nutrition*, 42(9), 990–1000. <https://doi.org/10.1080/01904167.2019.1589505>

Zeng, H. Y., Chen, L. H., Yang, Y., Deng, X., Zhou, X. H., & Zeng, Q. R. (2019). Basal and foliar treatment using an organic fertilizer amendment lowers cadmium availability in soil and cadmium uptake by rice on field micro-plot experiment planted in contaminated acidic paddy soil. *Soil and Sediment Contamination*, 28(1), 1–14. <https://doi.org/10.1080/15320383.2018.1525336>

DOI: <http://doi.org/10.17503/agrivita.v42i1.2324>



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

AGRIVITA

Journal of Agricultural Science

Universitas Brawijaya Online Journal - © 2016

Powered by [Open Journal System 2.4.7.1](https://www.openjournal.org/)

Visitors

ID	29503	US	2924
IN	2114	PH	1933
MY	1581	IR	1162
TH	1127	CN	599
JP	434	RU	399
Newest:	KG You:	ID	
Today:		29	
Month:		1013	
Total:		47589	
Supercounters.com			

Keywords

2AP Agroforestry Biological control
Chrysanthemum DNA extraction
Morphology *Oryza sativa* REDD+ Rice
SSR markers corn **diversity** drought
genotype land productivity maize mungbean
population **rice** soil fertility soybean

Journal Content

Search

Search Scope

Browse

- By Issue
- By Author
- By Title

Information

- For Readers
- For Authors
- For Librarians

Journal Help



This work is licensed under a

[Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)



Direct Seed Planting System and Giving Liquid Organic Fertilizer as A New Method to Increase Rice Yield and Growth (*Oryza sativa* L.)

Mohamad Ikbal Bahua*) and Hayatiningsih Gubali

Faculty of Agriculture, State University of Gorontalo, Indonesia

ARTICLE INFO

Keywords:

Direct seed planting
 Liquid organic fertilizers
 Transplanting
 Rice

Article History:

Received: August 8, 2019

Accepted: December 13, 2019

*) Corresponding author:

E-mail: mohamad.bahua@ung.ac.id

ABSTRACT

The transplanting system is a common method in rice cultivation, whereas the direct seed planting system combined with the application of organic fertilizer is rarely applied. This study aimed to determine the impact both the direct seed planting system and the application of liquid organic fertilizer the growth and production of rice. The research used the factorial randomized block design, direct seed planting system tested by the transplanting system and application of liquid organic fertilizer doses 25, 35, and 45 l/ha. The observed variables were plant height, number of tillers, productive tillers, panicle length, weight of 1000 rice grains, and grain yield of plots. Data were analyzed by analysis of variance with 5% LSD test. The results revealed that direct seed planting system and application of liquid organic fertilizer impact the growth and yield of rice. The application of liquid organic fertilizer a dose of 45 l/ha gave the highest influence among all observed variables. There is a correlation between the treatment of the planting system and the application of liquid organic fertilizer on the growth of the number of tillers and productive tillers. The direct seed planting system increases the yield of rice more than the transplanting system.

INTRODUCTION

The direct seed planting system applied in paddy rice is a cultivation technique that does not require an initial nursery phase. This has proven to be more efficient compared with the transplanting system, which requires the involvement of seedbed 3 weeks before the transfer into the paddy field. The direct seed planting system tends to increase output, mainly through time management production and land use.

The direct seed planting system is most widely used by Indonesian farmers in the cultivation of lowland rice. Direct seed planting systems and the application of liquid organic fertilizers can significantly increase rice production, because liquid organic fertilizers do not damage the environment and can improve soil fertility, so they can supply nutrients that can be absorbed by plants for a vegetative and generative growth (Kartika et al., 2018).

The use of organic fertilizer in cultivation aids

in maintaining fertility and physical, chemical, and biological soil properties. Liquid organic fertilizer is used in overcoming agricultural production constraints owing to their ability to provide the appropriate nutrients needed by plants (Moreno, Ondoño, Torres, & Bastida, 2017). Furthermore, the use of organic fertilizers serves to maintain the balance and sustainability of agricultural land ecosystems so that agricultural land can be sustainably used to produce food that is safe and healthy for public health (Martínez-Alcántara, Martínez-Cuenca, Bermejo, Legaz, & Quiñones, 2016).

The direct seed planting system is a good alternative to improve the production of lowland rice after the application of liquid organic fertilizer, as the nutrients that this fertilizer contains tend to be easily absorbed by plants. Moreover, there exist microorganisms that help reduce the incidence of nutrients getting washed away easily, subsequently providing a sufficient amount required for a quick

growth (Jones et al., 2013). The application of liquid organic fertilizers can help accelerate the absorption of nutrients from solid organic fertilizers, which are slowly absorbed by the plants in the rice cultivation system, as it takes a period of about 12 weeks to completely mineralize (Fahrurrozi, Mukhtamar, Setyowati, Sudjatmiko, & Chozin, 2019).

The results of a research by Leogrande, Vitti, Stellacci, Coccozza, & Ventrella (2016) revealed that the application of liquid organic fertilizer in rice cultivation is an alternative in increasing rice growth and production. Liquid organic fertilizer is more easily absorbed by plants because its elements contain decomposing bacteria that can provide nutrients needed by plants. The advantage of liquid fertilizer is that it contains nutrients, such as nitrogen (N), phosphorus (P), and potassium (K), which increase plant growth in the vegetative and generative phases (Colla, Roupheal, Canaguier, Svecova, & Cardarelli, 2014). This research will provide information on rice growth and production through the implementation of direct seed planting system and transplanting systems and the treatment with liquid organic fertilizer.

MATERIALS AND METHODS

The research was conducted in West Limboto District, Gorontalo Regency, from August 2017 to January 2018. Rice cultivation was done using the Legowo 4:1 row planting system. This research uses a complete randomized block design with two factors. The first factor is the planting system, which is the direct seed planting system and the transplanting system. The second factor is the dose of the liquid organic fertilizer: 25, 35, and 45 l/ha. So, six treatment combinations are repeated three times. The tools used in this study were hand tractors, hoes, sickles, hand sprayers, plastic buckets, meters, and stationery. The materials used were Mekongga variety of rice seeds, liquid organic fertilizer, pesticides, and water. The observed variables included plant height, number of tillers

and productive tillers, panicle length, weight of 1000 grains, and production per plot. The research data were analyzed statistically using analysis of variance, followed by the most significant difference test (LSD) at 5%. The six combinations of soil treatment and analysis at research sites are presented in Table 1 and Table 2.

RESULTS AND DISCUSSION

The direct seed planting system in rice cultivation has more advantages compared with the transplanting system, which include the following: the easy adaptation of seeds to the planting environment, faster harvest time of 10–15 days, less production and labor costs, and the use of seeds around 25–35 kg/ha.

Plant Growth

Direct seed planting system includes planting 2–3 rice seeds directly into the planting hole, without going through nurseries. To obtain optimal results, perfect tillage is needed to facilitate the growth of rice plants in the generative growth phase. Water management through drainage is carried out to prevent the seeds from being washed away during the plant growth process (Wang et al., 2016).

The most suitable location for the implementation of the direct seed planting system is the technical irrigated rice field agroecosystem. However, in dryland agroecosystems and rain-fed rice fields, the direct seed planting system can also be applied with water management requirements, and land preparation is carried out specifically by irrigating rice fields at the growth phase of rice using liquid organic fertilizers (Shah-Al-Emran, Krupnik, Kumar, Ali, & Pittelkow, 2019). The results revealed that the direct seed planting system, transplanting system, and treatment with liquid organic fertilizer had an impact on the height of rice plants, particularly the tall ones. The average height growth of rice plants is presented in Table 3.

Table 1. Combination treatment of planting systems and liquid organic fertilizers

No	Planting system	Dose of liquid organic fertilizer (l/ha)
1	Transplanting	25 l/ha ~ 3.48 kg N/ha ~ 4.33 kg P ₂ O ₅ /ha ~ 2.60 kg K ₂ O/ha
2	Direct Seed	25 l/ha ~ 3.48 kg N/ha ~ 4.33 kg P ₂ O ₅ /ha ~ 2.60 kg K ₂ O/ha
3	Transplanting	35 l/ha ~ 4.86 kg N/ha ~ 6.08 kg P ₂ O ₅ /ha ~ 3.64 kg K ₂ O/ha
4	Direct Seed	35 l/ha ~ 4.86 kg N/ha ~ 6.08 kg P ₂ O ₅ /ha ~ 3.64 kg K ₂ O/ha
5	Transplanting	45 l/ha ~ 6.25 kg N/ha ~ 7.81 kg P ₂ O ₅ /ha ~ 4.69 kg K ₂ O/ha
6	Direct Seed	45 l/ha ~ 6.25 kg N/ha ~ 7.81 kg P ₂ O ₅ /ha ~ 4.69 kg K ₂ O/ha

M. Ikbal Bahua et al.: *New method to increase rice yield*

Table 2. Soil analysis at the research site

Type of analysis	Unit	Soil	
		Results*	Criteria**
pH H ₂ O	-	3.72	Very Sour
pH KCl	-	3.52	Sour
C-Organic	%	1.83	Low
N-Total	%	0.16	Low
P-Bray I	µg/g	5.38	Very Low
K-dd	cmol (+) / kg	0.52	Medium
Na	cmol (+) / kg	0.45	Medium
Ca	cmol (+) / kg	3.02	Low
Mg	cmol (+) / kg	0.52	Low
Cation exchange capacity	cmol (+) / kg	17.41	Medium
Al-dd	cmol (+) / kg	1.45	-
H-dd	cmol (+) / kg	0.42	-
Texture			Clay
Sand	%	21.35	
Dust	%	34.95	
Clay	%	46.67	

Remarks: *) The results of data analysis from the soil fertility laboratory, Faculty of Agriculture, Sam Ratulangi University, Manado, 2018; **) The criteria based on land research center 1983.

Table 3. Average plant height growth in the planting system and liquid organic fertilizer

Treatment	Plant height (cm)			
	3 wap	5 wap	7 wap	9 wap
Planting system				
Transplanting	35.23 b	55.46 b	83.57 b	99.58 b
Direct seed	30.45 a	61.67 a	93.56 a	107.57 a
LSD 5%	1.70	3.72	2.49	2.71
Liquid organic fertilizer (l/ha)				
25	31.03	53.63 a	83.34 a	96.48 a
35	35.89	60.21 b	95.54 b	109.09 b
45	43.46	63.36 b	103.31 c	115.49 c
LSD 5%	ns	4.56	3.05	3.32

Remarks: The numbers followed by different letters in the same column are significantly different from the 5% LSD test; wap = week after planting

The results of the research presented in Table 3 reveal that at the direct seed planting system, the average height of rice plants was higher compared with that of rice plants at the transplanting system. The average increase in the growth of rice plants occurs at 5, 7, and 9 wap. At 9 wap, the height of rice plants at the direct seed planting system reached 107.57 cm, whereas in the transplanting system, it only reached 99.58 cm, which means that at the

direct seed planting system, the rice plant height increased by 7.99 cm. At the direct seed planting system, proper spacing can facilitate the entry of sunlight which is useful to accelerate the process of photosynthesis (Andres, Fogliatto, Ferrero, & Vidotto, 2015). The lengthening of a segment or increase in plant height can be caused by auxin activity or natural growth regulators in plants contained in liquid organic fertilizer. Auxin will carry out its function in cell

elongation in the direction of the entry light, causing the plants to grow taller (Richards, 2008).

The results revealed that the direct seed planting system, transplantation system, and treatment with liquid organic fertilizer have an impact on the growth of the number of tillers at 3, 5, and 7 wap. The average growth of the number of tillers in the planting system treatment and the application of liquid organic fertilizer is presented in Table 4.

Table 4. Average number of tillers in the planting system treatment and the application of liquid organic fertilizer

Treatment	Number of tillers (stem)		
	3 wap	5 wap	7 wap
Planting system			
Transplanting	6.29 b	14.57 a	25.94 a
Direct seed	7.65 a	17.37 b	29.06 b
LSD 5%	0.59	0.73	0.98
Liquid organic fertilizer (l/ha)			
25	6.46	14.26 a	25.32 a
35	7.51	16.67 b	27.72 b
45	6.93	16.99 b	29.44 c
LSD 5%	ns	0.89	1.20

Remarks: The numbers followed by different letters in the same column are significantly different from the 5% LSD test; wap = week after planting

Table 4 demonstrates that the direct seed planting system and the treatment with liquid organic fertilizer affect the growth of the number of tillers compared with the transplanting system. The growth in the number of tillers in the direct seed planting system at 7 wap reached 29.06 number of tillers, while in the transplanting system, it only reached to 25.94 number of tillers. This means that in the direct seed planting system, there is an increase of 3.12 in the number of tillers.

The results of the research by Javaid et al. (2012) revealed that in the direct seed planting system, seed growth is not tight because only 1–2 seeds are planted on each hole, so plants can easily absorb nutrients and get sunlight which help in the optimal metabolic process for growth in the number of tillers. However, in the transplanting system, the number of clumps of plants moved from the nursery will affect the density of plant growth; therefore, it will reduce the quality of plant clumps in increasing the growth of the number of tillers (Thakur & Uphoff,

2017). The correlation between the direct seed planting system and transplanting system with the liquid organic fertilizer on the growth of the number of tillers is explained in Fig. 1.

Fig. 1 explains that the direct seed planting system, transplanting system, and treatment with liquid organic fertilizer affect the growth of the number of tillers. The direct seed planting system will reach the generative stage faster, thus shortening the production period and increasing stem growth and rice plant population, whereas the transplanting system has a long production period, because plants need time to adapt to the planting environment after being removed from the nursery (Birhane, 2013).

The results revealed that the direct seed planting system and treatment with liquid organic fertilizer significantly affected the growth of the number of productive tillers, as a marked level of interaction is identified as explained in Fig. 2.

Fig. 2 explains that the interaction between the treatment of cropping systems and liquid organic fertilizer affects the growth of the number of productive tillers. The direct seed planting system and the application of liquid organic fertilizer at a dose of 45 l/ha had a higher influence on the growth of the number of productive tillers compared with that on the growth of the number of productive tillers in the transplanting system and application of liquid organic fertilizer at a dose of 45 l/ha. The application of liquid organic fertilizer can increase the growth of the number of productive tillers, especially in the 5 to 7 wap age phase of the rice, because in this phase the growth of leaves and stems of rice plants increases so that it can support the growth of a number of productive tillers, so liquid organic fertilizer is needed to provide organic material that can supply nutrients such as N, P, and K during the generative growth phase (Dimkpa & Bindraban, 2016).

The availability of nutrients in the soil is a factor that can increase crop production. If the land cannot provide sufficient nutrients for the plants, then the application of fertilizer is necessary to overcome nutrient deficiency (Qiao et al., 2011). Therefore, the application of liquid organic fertilizer is very helpful for plant growth, especially in the supply of nutrients such as N, P, and K, because liquid organic fertilizer contains essential nutrients and bacteria that can increase soil fertility (Zeng et al., 2019). The relationship between nutrient absorption of N, P, K, and plant growth is explained in Fig. 3.




Fig. 1. The effect of the average growth of the number of tillers on the correlation between the direct seed planting system and the planting system as well as between the treatment of liquid organic fertilizer.




Fig. 2. The average number of productive tillers in the interaction of direct seed planting system and transplanting with the liquid organic fertilizer treatment.




Fig. 3. Relationship between nutrient absorption and plant growth

Fig. 3 shows that the results of the regression analysis of the relationship between nutrient absorption of N, P, and K (kg/ha) and the growth of rice plants reveal a linear pattern, so that the increased absorption of nutrients N, P, and K contained in higher liquid organic fertilizer can increase higher plant growth. During the growth of rice plants, the

absorption of nutrients N, P, and K of 9.505 kg/ha will increase plant height growth by 88.01 cm. The absorption of nutrients N, P, K of 10.72 kg/ha on rice growth will increase the growth of 24,443 number of tillers. Furthermore, the absorption of N, P, K nutrients of 2.06 kg/ha on rice growth will increase growth by 23,373 number of productive tillers.

The growth of height plants in a cumulative manner (R^2) absorbs 96.57% of nutrients N, P, and K, meaning that 3.43% of rice plant height growth is influenced by other factors. The growth of the number of tillers in a cumulative manner (R^2) absorbs 99.99% of nutrients N, P, and K, meaning that 0.01% of the growth of the number of tillers is influenced by other factors. The growth of the number of productive tillers in a cumulative manner (R^2) absorbs 99.1% of nutrients N, P, and K, meaning that 0.09% of the growth in the number of productive tillers is influenced by other factors.

Nitrogen nutrients are part of the chlorophyll molecule because the provision of nitrogen nutrients in sufficient quantities will increase the vegetative growth of rice plants, particularly increase in plant height and number of tillers. Nitrogen is also a functionally important constituent of enzymes that plays a vital role in plant metabolism, because the enzymes are composed of proteins (Gopal Ramdas et al., 2017). Nitrogen is a very mobile element in plants, which means that functional proteins that

contain N can decompose in older plant parts. It will then be transported through the process of photosynthesis to young plant tissues that grow actively in the vegetative growth phase (Smith, Brye, Gbur, Chen, & Korth, 2014). Phosphorus plays a role in strengthening plant root systems in order to increase the number of tillers and strengthen plant stems, whereas K helps in forming proteins and carbohydrates that can accelerate the growth of the number of productive tillers (Xiao et al., 2019).

Plant Yield

Direct seed planting systems that use superior seeds, balanced fertilization, and good water management can achieve yields of around 5–6.2 t/ha of harvested unhusked rice when compared with the results on the transplanting system which only reaches 3–4.77 t/ha of the dried grain harvest. Rice yields can be seen from the following indicators: panicle length, 1000 grain weight, and yield per plot. The rice yields at the direct seed planting, transplanting system, and the liquid organic fertilizer treatment are presented in Fig. 4.




Fig. 4. Average crop yields at direct seed planting system and transplanting system




Fig. 5. The average yield of rice at the treatment with liquid organic fertilizer

Fig. 4 shows that panicle length at the direct seed planting system reached 35.18 cm, whereas in the transplanting system, it only reached 27.92 cm. The weight of 1000 grains at the direct seed planting system reached 40.25 g, whereas that at the transplanting system only reached 30.78 g. The yield per plot at the direct seed planting system reached 6.93 kg/ha, and at the transplanting system, the yield per plot only reached 5.49 kg/ha.

The increase in panicle length is usually caused by genetic and environmental factors, such as plant age, number of productive tillers, and water availability in the generative phase. The increase in panicle length will affect the number of results, because as each panicle length increases, the branches of grain stalk will grow to produce more grain (Sharma, Giri, & Tyagi, 2016). The increase in the weight of 1000 grains is influenced by genetic factors related to the shape and size of the grain. The average grain weight is a characteristic of each rice variegation that plays a role in rice production. Dry matter in the grains is determined based on filled rice grains produced by rice panicles whose growth increases by photosynthesis occurring in plant parts (Lal et al., 2020). The determining component of rice production is the percentage of the number of productive tillers, the weight of 1000 grains, and panicle length. The more the number of productive tillers and the grains of filled panicle during the generative growth phase, the higher the productivity of rice per hectare (Tun, Guo, Fang, & Tian, 2018).

The results in Fig. 4 reveal that rice production is higher in the direct seed planting system compared with that in the transplanting system. Rice production in the direct seed planting system reached 6.93 kg/ha, whereas that in the transplant system reached 5.49 kg/ha. Based on the distance

of the 4:1 Legowo system with a spacing of 25 × 25 cm, with a planting area of 625 cm (6.25 m²), the rice production in each planting system is converted to tons/ha, so that the production of rice in the direct seed planting system reached 11.1 t/ha, and that in the transplanting system reached 8.8 t/ha. Liquid organic fertilizer is a technology that can increase rice yield, because the amount of nutrients in the soil, especially N, P, and K, is very limited, so it requires additional nutrients from liquid organic fertilizer. The impact of liquid organic fertilizer on rice yield is explained in Fig. 5.

Fig. 5 explains that the treatment with liquid organic fertilizer with a dose of 45 l/ha has an effect on panicle length, weight of 1000 grains, and yield per plot. Applying liquid organic fertilizer to plants can increase crop production through the activation of microorganisms contained therein and those in the environment. The use of liquid organic fertilizer will provide better quality of rice yields, because it is more evenly distributed. Moreover, there is no accumulation of fertilizer concentration in one place, because organic fertilizer is not problematic with regard to nutrient leaching, and it is able to overcome deficiency in nutrients needed by plants for increased rice yields (Yoshinaga, Takai, Arai-Sanoh, Ishimaru, & Kondo, 2013). The relationship between nutrient absorption of N, P, and K and plant yield is explained in Fig. 6.

Fig. 6 shows that the results of the regression analysis of the relationship between nutrient absorption of N, P, and K (kg/ha) and plant yields revealed a linear pattern, so that the increased absorption of nutrients N, P, and K contained in higher liquid organic fertilizer can increase higher plant yields. Each rice plant yield absorbs 3.275 kg/ha of N, P, and K nutrients so that the panicle length will increase by 22.473 cm.




Fig. 6. Relationship between nutrient absorption and plant yield

Then every absorption of 2,385 kg/ha of nutrients N, P, and K will increase the weight of 1000 grains by 25.31 g, and each absorption of 0.34 kg/ha of nutrients N, P, and K will increase rice yield per plot by 2.8467 kg.

The growth of panicle length cumulatively (R^2) absorbs N, P, and K nutrients by 97.35%, meaning that 2.65% of panicle length growth is influenced by other factors. The absorption of the nutrients N, P, and K to cumulatively produce the weight of 1000 grains (R^2) is 79.23%, meaning that 20.77% of the resulting weight of 1000 grains of rice is influenced by other factors. The absorption of N, P, and K nutrients to produce rice production per plot cumulatively (R^2) is 99.54%, meaning 0.46% rice production yield per plot is influenced by other factors.

Increasing the doses of organic fertilizer will increase the availability of nutrients N, P, and K in the soil. Nutrients N, P, and K play a role in photosynthesis, which can affect the increase in rice yields, such as panicle length, grain number, and grain weight. Nutrient N increases the chlorophyll content of plant leaves so that plant photosynthesis increases. The application of liquid organic fertilizer in the rice planting process can increase the availability of N in the final stages of rice growth, which can affect leaf metabolism during seed filling (Santosa & Suryanto, 2015). The nutrient element P plays a role in the supply and transfer of energy throughout the biochemical process of rice, one of which is the acceleration of the cooking and development of grain so that the grain weight increases (Douglas & Dyson, 1985). Nutrient K plays a role in the formation of sugars, starches, and various enzymes to increase the amount of grain per panicle and grain weight percentage (Zandvakili, Barker, Hashemi, Etemadi, & Autio, 2019).

CONCLUSION

The treatment of direct seed planting system confers a direct effect on the growth and yield of paddy rice, in contrast with the transplanting system, and the best liquid organic fertilizer treatment was at a dose of 45 l/ha. Based on the growth of plants, there was also a better influence on the number of tillers and productive tillers, whereas the enhanced panicle length, production, and weight of 1000 grain are indicative of a superior yield. This suggests that the direct seed planting system combined with the application of liquid organic fertilizers by farmers as

an improvement effort promotes plant yield, as well as vegetative and generative growth.

ACKNOWLEDGEMENT

The results of this study are research funded by the Faculty of Agriculture, the State University of Gorontalo as a research-based study program and science.

REFERENCES

- Andres, A., Fogliatto, S., Ferrero, A., & Vidotto, F. (2015). Growth variability of Italian weedy rice populations grown with or without cultivated rice. *Crop Science*, 55(1), 394–402. <https://doi.org/10.2135/cropsci2013.07.0472>
- Birhane, A. (2013). Effect of planting methods on yield and yield components of rice (*Oryza sativa* L.) varieties in Tahtay Koraro Wereda, Northern Ethiopia. *International Journal of Technology Enhancements and Emerging Engineering Research*, 1(5), 1–5. Retrieved from <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.407.6481&rep=rep1&type=pdf>
- Colla, G., Roupael, Y., Canaguier, R., Svecova, E., & Cardarelli, M. (2014). Biostimulant action of a plant-derived protein hydrolysate produced through enzymatic hydrolysis. *Frontiers in Plant Science*, 5, 448. <https://doi.org/10.3389/fpls.2014.00448>
- Dimkpa, C. O., & Bindraban, P. S. (2016). Fortification of micronutrients for efficient agronomic production: a review. *Agronomy for Sustainable Development*, 36, 7. <https://doi.org/10.1007/s13593-015-0346-6>
- Douglas, J. A., & Dyson, C. B. (1985). Preliminary investigation of concentrations of minerals and nitrogen in wheat grain, and their relationship with baking quality and grain weight. *New Zealand Journal of Agricultural Research*, 28(1), 81–85. <https://doi.org/10.1080/00288233.1985.10426999>
- Fahrurrozi, Muktamar, Z., Setyowati, N., Sudjatkiko, S., & Chozin, M. (2019). Comparative effects of soil and foliar applications of Tithonia-enriched liquid organic fertilizer on yields of sweet corn in closed agriculture production system. *AGRIVITA Journal of Agricultural Science*, 41(2), 238–245. <https://doi.org/10.17503/agrivita.v41i2.1256>
- Gopal Ramdas, M., Manjunath, B. L., Narendra Pratap, S., Ramesh, R., Verma, R. R., Marutao, L. A., ... Rahul, K. (2017). Effect of organic and inorganic sources of nutrients on soil microbial activity and

M. Ikbal Bahua et al.: *New method to increase rice yield*

- soil organic carbon build-up under rice in west coast of India. *Archives of Agronomy and Soil Science*, 63(3), 414–426. <https://doi.org/10.1080/003650340.2016.1213813>
- Javaid, T., Awan, I. U., Baloch, M. S., Shah, I. H., Nadim, M. A., Khan, E. A., ... Abuzar, M. R. (2012). Effect of planting methods on the growth and yield of coarse rice. *Journal of Animal and Plant Sciences*, 22(2), 358–362. Retrieved from <http://www.thejaps.org.pk/docs/v-22-2/20.pdf>
- Jones, D. L., Cross, P., Withers, P. J. A., DeLuca, T. H., Robinson, D. A., Quilliam, R. S., ... Edwards-Jones, G. (2013). REVIEW: Nutrient stripping: the global disparity between food security and soil nutrient stocks. *Journal of Applied Ecology*, 50(4), 851–862. <https://doi.org/10.1111/1365-2664.12089>
- Kartika, Lakitan, B., Sanjaya, N., Wijaya, A., Kadir, S., Kurnianingsih, A., ... Meihana, M. (2018). Internal versus edge row comparison in jajar legowo 4:1 rice planting pattern at different frequency of fertilizer applications. *AGRIVITA Journal of Agricultural Science*, 40(2), 222–232. <https://doi.org/10.17503/agrivita.v40i0.1715>
- Lal, B., Gautam, P., Nayak, A. K., Maharana, S., Tripathi, R., Shahid, M., ... Singh, S. (2020). Tolerant varieties and exogenous application of nutrients can effectively manage drought stress in rice. *Archives of Agronomy and Soil Science*, 66(1), 13–32. <https://doi.org/10.1080/03650340.2019.1587749>
- Leogrande, R., Vitti, C., Stellacci, A. M., Cocozza, C., & Ventrella, D. (2016). Response of wheat crop during transition to organic system under Mediterranean conditions. *International Journal of Plant Production*, 10(4), 565–578. <https://doi.org/10.22069/ijpp.2016.3050>
- Martínez-Alcántara, B., Martínez-Cuenca, M. R., Bermejo, A., Legaz, F., & Quiñones, A. (2016). Liquid organic fertilizers for sustainable agriculture: Nutrient uptake of organic versus mineral fertilizers in citrus trees. *PLoS ONE*, 11(10), e0161619. <https://doi.org/10.1371/journal.pone.0161619>
- Moreno, J. L., Ondoño, S., Torres, I., & Bastida, F. (2017). Compost, leonardite, and zeolite impacts on soil microbial community under barley crops. *Journal of Soil Science and Plant Nutrition*, 17(1), 214–230. <https://doi.org/10.4067/S0718-95162017005000017>
- Qiao, J., Liu, Z., Deng, S., Ning, H., Yang, X., Lin, Z., ... Ding, Y. (2011). Occurrence of perfect and imperfect grains of six japonica rice cultivars as affected by nitrogen fertilization. *Plant and Soil*, 349(1–2), 191–202. <https://doi.org/10.1007/s11104-011-0861-4>
- Richards, R. A. (2008). Genetic opportunities to improve cereal root systems for dryland agriculture. *Plant Production Science*, 11(1), 12–16. <https://doi.org/10.1626/pps.11.12>
- Santosa, M., & Suryanto, A. (2015). The growth and yield of paddy Ciherang planted in dry and wet season and fertilized with organic and inorganic fertilizers. *AGRIVITA Journal of Agricultural Science*, 37(1), 24–29. <https://doi.org/10.17503/agrivita-2015-37-1-p024-029>
- Shah-Al-Emran, Krupnik, T. J., Kumar, V., Ali, M. Y., & Pittelkow, C. M. (2019). Agronomic, economic, and environmental performance of nitrogen rates and source in Bangladesh's coastal rice agroecosystems. *Field Crops Research*, 241, 107567. <https://doi.org/10.1016/j.fcr.2019.107567>
- Sharma, G., Giri, J., & Tyagi, A. K. (2016). Sub-functionalization in rice gene families with regulatory roles in abiotic stress responses. *Critical Reviews in Plant Sciences*, 35(4), 231–285. <https://doi.org/10.1080/07352689.2016.1265357>
- Smith, S. F., Brye, K. R., Gbur, E. E., Chen, P., & Korth, K. (2014). Residue and water management effects on aggregate stability and aggregate-associated carbon and nitrogen in a wheat–soybean, double-crop system. *Soil Science Society of America Journal*, 78, 1378–1391. <https://doi.org/10.2136/sssaj2013.12.0534>
- Thakur, A. K., & Uphoff, N. T. (2017). How the system of rice intensification can contribute to climate-smart agriculture. *Agronomy Journal*, 109(4), 1163–1182. <https://doi.org/10.2134/agronj2016.03.0162>
- Tun, T. N., Guo, J., Fang, S., & Tian, Y. (2018). Planting spacing affects canopy structure, biomass production and stem roundness in poplar plantations. *Scandinavian Journal of Forest Research*, 33(5), 464–474. <https://doi.org/10.1080/02827581.2018.1457711>
- Wang, W., Chen, Q., Hussain, S., Mei, J., Dong, H., Peng, S., ... Nie, L. (2016). Pre-sowing seed treatments in direct-seeded early rice: Consequences for emergence, seedling growth and associated metabolic events under chilling stress. *Scientific Reports*, 6, 19637. <https://doi.org/10.1038/srep19637>

M. Ikbal Bahua et al.: *New method to increase rice yield*

- Xiao, M., Zang, H., Ge, T., Chen, A., Zhu, Z., Zhou, P., ... Kuzyakov, Y. (2019). Effect of nitrogen fertilizer on rice photosynthate allocation and carbon input in paddy soil. *European Journal of Soil Science*, *70*(4), 786–795. <https://doi.org/10.1111/ejss.12811>
- Yoshinaga, S., Takai, T., Arai-Sanoh, Y., Ishimaru, T., & Kondo, M. (2013). Varietal differences in sink production and grain-filling ability in recently developed high-yielding rice (*Oryza sativa* L.) varieties in Japan. *Field Crops Research*, *150*, 74–82. <https://doi.org/10.1016/j.fcr.2013.06.004>
- Zandvakili, O. R., Barker, A. V., Hashemi, M., Etemadi, F., & Autio, W. R. (2019). Comparisons of commercial organic and chemical fertilizer solutions on growth and composition of lettuce. *Journal of Plant Nutrition*, *42*(9), 990–1000. <https://doi.org/10.1080/01904167.2019.1589505>
- Zeng, H. Y., Chen, L. H., Yang, Y., Deng, X., Zhou, X. H., & Zeng, Q. R. (2019). Basal and foliar treatment using an organic fertilizer amendment lowers cadmium availability in soil and cadmium uptake by rice on field micro-plot experiment planted in contaminated acidic paddy soil. *Soil and Sediment Contamination*, *28*(1), 1–14. <https://doi.org/10.1080/15320383.2018.1525336>