

Evaluation of the Antiplasmodial Properties of Andrographis paniculata (Burm.f.) and Peperomia pellucida (L.) Kunth

by Weny Musa

Submission date: 09-Apr-2023 02:41AM (UTC-0400)

Submission ID: 2059404755

File name: 74481-278396-3-PB.pdf (658.24K)

Word count: 5963

Character count: 30501

Evaluation of the Antiplasmodial Properties of *Andrographis paniculata* (Burm.f.) and *Peperomia pellucida* (L.) Kunth

Nurhayati Bialangi¹, Mohamad Adam Mustapa², Yuszda Salimi¹, Weny Musa¹, Ari Widiyantoro³, Agus Malik Ibrahim⁴, Boima Situmeang⁴, and Julinton Sianturi⁵

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

²Department of Pharmacy, Faculty of Sport and Health Sciences, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Tanjungpura, Pontianak 78115, Indonesia

⁴Department of Chemistry, Sekolah Tinggi Analisis Kimia Cilegon, Banten, 42411, Indonesia

⁵Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, 14424 Potsdam, Germany

* Corresponding author:

tel: +62-85262897312

email: boimatumeang@gmail.com

Received: May 4, 2022

Accepted: October 18, 2022

DOI: 10.22146/ijc.74481

Abstract: Plasmodium species are the infectious agents that are responsible for malaria, a disease that claims the lives of approximately 400,000 people annually. The fact that drug resistance against malaria is on the rise suggests that new antimalarial compounds need to be discovered. It is well known that medicinal plants present the best opportunity for the identification of novel antimalarial chemicals. Both the *Andrographis paniculata* (Burm.f.) and *Peperomia pellucida* (L. Kunth) species have been tested for their antiplasmodial ability against the *Plasmodium falciparum* strain. The *P. pellucida* (L. Kunth) species has also been subjected to in vitro and in vivo biological research. *P. pellucida* was used to isolate the steroid known as 3-hydroxy-24-ethyl-5,22-cholestadiene (1) and the triterpenoid known as 3-hydroxy-9-lanosta-7,24E-dien-26-oic acid (2). Both compounds were then tested for their activity in vitro. In the mice model, triterpenoid 2 had a substantial chemo-suppressive impact.

Keywords: *A. paniculata* (Burm.f.); *P. pellucida* L. Kunth; *Plasmodium falciparum*; inhibition; 3 β -hydroxy-9-lanosta-7,24E-dien-26-oic acid

■ INTRODUCTION

Malaria is a significant threat to global public health, as it affects individuals in every region of the world. The disease is brought on by the parasite *Plasmodium*, which is transmitted from female *Anopheles* mosquitoes to human hosts [1]. Since there are not enough preventative measures in place, and because the malaria parasite is becoming more resistant to many antimalarial drugs that are already on the market, such as quinine, chloroquine, and piperazine; this disease continues to be a concern for the health of people all over the world [2-4]. In spite of the devoted research that has been conducted over the

course of several decades, there is still no malaria vaccine that can be bought in stores at this moment. As a result, chemotherapies that make use of combination medications appear to be the best alternative, and novel antimalarial chemicals are an absolute necessity. Regular mining is done on plants that are used for therapeutic purposes with the purpose of determining whether they have the potential to provide novel active chemicals [5]. For instance, Tu [6] created Artemisinin in China in the early 1970s. This medication was derived from *Artemisia annua* L. and was utilized in the treatment of malaria. It is widely known that tropical plants contain a rich reservoir of bioactive compounds. Because of this, there

is the possibility that these plants could serve as sources for the development of novel antimalarial drugs [7].

One of the most common plant families that are farmed in tropical and subtropical regions is Acanthaceae family [8]. Traditional uses for this herb include the treatment of acute diarrhea, coughing, the common cold, inflammation, boils, skin eruptions, and seasonal fever [9]. Other ailments that have been alleviated by using this herb include boils, skin eruptions, and seasonal fever. The investigation of phytochemicals has resulted in the identification of a wide range of secondary metabolites that have significant pharmacological effects. The *Andrographis paniculata* species, more often known as "Sambiloto," which could be found all over Indonesia, is regularly exploited as an element of traditional medical practices [10]. Traditional cultures held the belief that combining this plant with *Peperomia pellucida* could boost the efficacy of antimalarial treatment [11-13]. In spite of the fact that this plant possesses anti-plasmodial activity against many strains of *Plasmodium falciparum*, this is still the case. We investigated the antiplasmodial activity of the *A. paniculata*, and *P. pellucida* isolates against a malarial parasite strain in order to give data in favor of this theory. In addition, we tested the efficacy of these compounds on infected mice and isolated the active compound(s) in an effort to gain a deeper comprehension of the biological role that these compounds play. On the other hand, the investigations regarding the isolated compounds of PP have not been documented in detail. Only one of the studies that we looked at in the literature had isolated chemicals that came from the *P. pellucida* plant.

■ EXPERIMENTAL SECTION

Materials

In June 2016, fresh leaves and stems of *A. paniculata* and *P. pellucida* were harvested in north Gorontalo, Gorontalo Province, Indonesia. The plant was identified at the Herbarium Biology Laboratory, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo, and voucher specimens (No. 130/H47.B4.Bio.Lab Bio/LL/2016; for *P. pellucida* and No.

131/H48.B4.Bio.Lab Bio/LL/2016; for *A. paniculata*) were deposited to the Herbarium.

Instrumentation

The instruments used in the experiment for molecular determination were Japan Electron Optics Laboratory (JEOL) 125 MHz ^1H -NMR and JEOL 500 MHz ^{13}C -NMR, for the photoelectron analysis the instrument used was JEOL UV-Vis Spectrophotometer, the cell culture was done in ESCO Scientific Laminar Cabinet, and the infrared instrument used was ThermoFisher Scientific Infrared Spectrometer.

Procedure

Plant extraction and isolation

Dried leaves and stems of *P. pellucida* (PP) (2 kg) and *A. paniculata* (AP) (2 kg) were macerated with methanol (12 L) at room temperature for 2 d. After filtrating, the filtrate was evaporated under reduced pressure to yield the PP extract (211 g) and AP extract (278 g). The PP extract was suspended with water (500 mL) and partitioned successively with *n*-hexane and ethyl acetate. Evaporation gave the crude *n*-hexane (15.3 g), ethyl acetate (13.2 g) and aqueous (18.5 g) extracts. The isolation procedures are shown in detail in Scheme S1. The ethyl acetate extract was purified by column chromatography using silica gel G60 (132 g), eluted by *n*-hexane/EtOAc (200 mL) with a linear gradient from 10% to 100% of EtOAc to yield eleven fractions (A1–A11). Fraction A6 (0.27 g) was subjected to silica column chromatography using the isocratic elution mixture of *n*-hexane/EtOAc (90:10-0:100) to yield 40 fractions (B1–B40). Fractions B9 were recrystallized using *n*-hexane and chloroform to afford compound **1** (18 mg) as a needle-shaped crystal. Fraction A7 (0.97 g) was subjected to silica column chromatography (10 g of silica G60) using an isocratic system, eluted by *n*-hexane/EtOAc/MeOH (7:2:1) to yield 25 fractions (B1–B25). Fractions C13–C15 (90 mg) were combined and purified by silica column chromatography on Octa Decyl Silane (ODS), eluted by MeOH/H₂O (4:1) to yield compound **2** (12 mg) as a white powder.

Synchronization and maintenance of parasite strain

The University of Tokyo's chloroquine-sensitive malaria parasite Pf D-10. Parasite Pf D-10 was maintained in fresh O+ human erythrocytes with a hematocrit of 4% in RPMI 1640 culture medium (Gibco) supplemented with 10% (v/v) human serum, 25 mM HEPES (N-(2-hydroxyethyl)piperazine-N'-2-ethanesulfonic acid, Sigma Aldrich), 45 µg/mL hypoxanthine, 32 mM NaHCO₃, and 50 µg/mL gentamicin, incubated at 37 °C under a gas mixture of 5% O₂ and 5% CO. After 96 h, parasites were microscopically synchronized with 5% sorbitol.

In vivo antiparasmodial activity

The Animal Experimental Development Unit at Universitas Gadjah Mada, Yogyakarta, Indonesia, donated 30 male BALB/c albino mice aged 6–8 weeks weighing 25–28 g for this investigation. At the Universitas Airlangga Institute of Tropical Disease Animal Laboratory, the mice were fed and watered ad libitum. Eijkman Institute for Molecular Biology in Jakarta provided the parasite Pb ANKA strain. At the Institute of Tropical Disease, Universitas Airlangga, male BALB/c mice and cryogenic storage maintained the parasite.

As described by Peter [14], with slight modifications, the sample extract AP/PP (20/80 mg) and isolated chemical (10 mg) were tested in mice infected with Pb ANKA for antiparasmodial activity. Thirty BALB/c albino mice were intraperitoneally infected with Pb ANKA (106 parasitized erythrocytes). The volume of Inoculum used was 200 µL. Five six-mouse groups were formed (three experimental and two control groups). Three groups received the sample at 1, 10, and 100 mg/kg/body in 0.5% CMC-Na (200 µL). A negative control group received 0.5% CMC-Na while a positive control group received chloroquine diphosphate (25 mg/kg/body) intraperitoneally once a day for 4 d. After 4 d, the animals ceased treatment. On the 6th day, thin mouse tail blood smears were made on slides. The slide was stained with 15% Giemsa-dyed solution for 10 min [15-16]. Rinsing and drying at room temperature completed the slide. Giemsa-stained blood smears and microscopy determined parasitemia and growth inhibition. Day 21 post-infection survival was monitored.

Antimalarial assay in vitro

The antimalarial activity of the compounds was tested using the Desjardins method [17]. The test was performed using *P. falciparum* 3D7 strain. The antimalarial examination used 96 wells, each well filled by parasitemia culture 1%. The RPHS media was replaced by a sample containing RPHS media with different sample concentrations. All wells were then stored in the Laminar Airflow cabinets until all solvents were evaporated. Further, we added 50 µL of parasitemia red blood cells into the well and then incubated at 37 °C. After 48 h, slide preparation was stained using Giemsa 20%. The parasite was then swapped with Giemsa-colored blood. The parasitemia percentage was calculated by comparing the number of infected erythrocytes to 500 erythrocytes.

Statistical analysis

The experiments were performed independently in triplicate, and the average data are presented. Statistical analysis was performed by unpaired two-tailed *t*-test using GraphPad Prism. The statistics were significant when **p* < 0.05, ***p* < 0.01 and ****p* < 0.001.

RESULTS AND DISCUSSION

Fresh leaves and stems of species AP and PP were extracted with crude methanolic extracts. A thin blood smear stained with Giemsa was obtained after incubating synchronized *P. falciparum* D-10 strain (Pf D-10) with the AP-PP extract at varied ratios (80:20 to 20:80) in dimethyl sulfoxide (DMSO), chloroquine (CQ) (1 M), and 0.2% DMSO. Microscopically; parasitemia, parasite growth, and inhibition rates were assessed. The chemoprevention of parasitemia against Pf D-10 was investigated. The activity of the AP-PP extract was limited (2.8 to 3.0%) at a ratio of 80:20 to 40:60. However, when the PP extract ratio was increased, the parasitemia rate reduced considerably (Fig. 1(a)). Low suppressions of Pf D-10 were seen at ratios ranging from 80:20 to 40:60, whereas a ratio of 20:80 indicated a 50% inhibitory effect (*p* 0.05). The AP-PP extract's antiparasmodial activity was boosted in a ratio-dependent way (Fig. 1(b)). Furthermore, the IC₅₀ value of AP-PP




Fig 1. The AP-PP extract was tested *in vitro* against *Pf D-10* at a variety of ratios, ranging from 80:20 to 20:80, as well as the PP extract on its own. (a) The level of parasitemia, and (b) the percentage of parasites that were inhibited by treatment with the AP-PP extract, (c) A result of the IC_{50} of the AP-PP extract when it was mixed in a ratio of 20:80. (d) The prevalence of parasitemia, and (e) the percentage of parasites that were inhibited by the PP extract. (f) A value for the IC_{50} of the PP extract. With the use of the GraphPad Prism program, an IC_{50} value was determined. As a negative control, we used parasites that had been treated with 0.2% DMSO, and as a positive control, we employed CQ at a dose of 1 μ M. Every experiment was carried out three times, so the total number of participants was three. The standard deviation is represented by the error bars. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, not significant (ns)

extract at a 20:80 ratio was investigated (Fig. 1(c)). The extract was tested at concentrations of 0.01, 0.1, 1, 10, and 100 g/mL and the inhibition rates were determined after 48 h of incubation. The IC_{50} value was calculated through an examination of inhibition-response curves. Based on

an *in vitro* study with an IC_{50} value of 62.01 g/mL, the AP-PP extract at a 20:80 ratio was rated mildly potent (Table S1) [17].

The PP extract alone must be assessed because its larger ratio had a stronger antiparasmodial effect than the

AP extract. *In vitro* concentrations were 0.01 to 100 g/mL. Fig. 1(d) shows that PP extract at 0.1 to 100 g/mL had antiparasmodial potential. The greatest dose of *P. pellucida* extract reduced parasitemia by 1.2% (p 0.01). PP extract was investigated for parasite PfD-10 inhibition (Fig. 1(e)). At 10 and 100 g/mL, 51% (p 0.01) and 92% (p 0.001) inhibition were observed. The PP extract demonstrated promising antiparasmodial activity with an IC_{50} of 4.0 g/mL (Table S1). According to Kwansa-Bentum et al. [17], PP L. Kunth crude methanolic extract had higher antiparasmodial activity than AP (Burm.f.) Nees (IC_{50} , 7.2 g/mL). These results indicate that the PP extract ratio increased antiparasmodial activity.

In addition, we tested the PP extract's efficacy in BALB/c albino mice infected with *Plasmodium berghei* ANKA (Pb ANKA) (Fig. S1). The extract had antiparasmodial action, with a 50% effective dose (ED_{50}) of 12.86 mg/kg/body weight (Fig. S1(a)). A strong chemosuppression of parasitemia (Fig. S1(b)), inhibitory effect (Fig. S1(d)), and chemosuppression of parasitemia from day 0 to day 4 were clearly demonstrated after the fourth day of treatment (Fig. S1(c)). The treatment of the *P. pellucida* extracts intraperitoneally improved the survival of infected mice (Fig. S1(e)).

The crude methanolic extract of PP L. Kunth demonstrated promising antiparasmodial action (Fig. 1(f)),




Fig 2. The IC_{50} values for each of the extracts compounds 1-2, and chloroquine, which served as a positive control. To determine the IC_{50} values of the separated compounds, concentrations ranging from 0.78 to 100 g/mL were used in the experiment

indicating the existence of chemicals responsible for this activity [17]. To further understand the mechanism of antiparasmodial activity of the PP extract, the compound(s) must be isolated and identified to assist in finding their biological function and mode of action. The antiparasmodial activity of crude *n*-hexane (E1), ethyl acetate (E2), and aqueous (E3) extracts were first determined. The antiparasmodial activity test shows the EtOAc (Fig. 2(b)) fraction has the most potent antiparasmodial activity due to having the lowest IC_{50} value of 1.533 $\mu\text{g/mL}$ among the other fraction of *n*-hexane (Fig. 2(a)) with IC_{50} of 3.892 $\mu\text{g/mL}$ and aqueous extract (Fig. 2(c)) with IC_{50} value of 5.016 $\mu\text{g/mL}$.

Because E2 was the most active extract in our investigation, we isolated and identified the compound(s) from it. The E2 was treated to standard, reverse-phase silica column chromatography and recrystallizations, yielding compounds **1**–**2** (Fig. 3).

The IR spectrum of compound **1** revealed the existence of O-H stretching, C-O bond, C-C, CH_2 vibrations, hydroxyl, and cycloalkane groups at 3373.6, 1247.0, 1641.0, 1457.3, 1381.6, 1038.0, and 881.6 cm^{-1} , respectively. Two methyl singlets, three methyl doublets, and a methyl signal as a triplet was observed in the proton NMR of **1**. Characteristic signals were also detected at 4.91, 5.18, and 5.31 ppm, indicating the presence of trisubstituted and disubstituted olefinic groups. The ^{13}C -NMR spectrum exhibited the presence of six sp^3 methyl carbons resonated at δ_{C} 12.4 (C-29), 18.4 (C-28), 19.3 (C-27), 20.4 (C-26), 12.9 (C-24) and 40.5 (C-18) and four sp^2 carbons resonated at δ_{C} 140.5, 121.9, 138.7 and 129.4 ppm indicated to a steroid compound related with stigmasterol

[18]. An oxygenated carbon peak at δ_{C} 71.9 ppm suggested the presence of a hydroxyl group and was usually located at C-3. Furthermore, the correlation of proton and carbon signals of **1** was confirmed by ^1H - ^1H COSY and ^1H - ^{13}C HMBC experiments. Therefore, isolated compound **1** was identified as 3 β -Hydroxy-24-ethyl-5,22-cholestadiene.

Compound **2** was obtained as a white crystal (12 mg), and the IR spectrum displayed absorption peaks for hydroxyl, carboxyl, carbonyl and olefinic groups. The ^1H -NMR spectrum of **2** showed the presence of seven signals of methyl groups as a singlet, resonating at δ_{H} 0.89–1.70 ppm. An oxymethine protons resonated at δ_{H} 3.49 (1H, dd, $J = 4.0$ and 12.0 Hz, H-3) indicated as a β isomer [19]. Additional functionalities included the signals of two olefinic protons resonated at δ_{H} 5.26 (1H, dd, $J = 6.5$ and 3.0 Hz (H-7) and 5.98 (1H, t, $J = 7.8$ Hz, H-24). A total of 30 carbons resonances were observed in the ^{13}C -NMR and DEPT 135° spectra. Four sp^2 carbons include two methine carbons and two quaternary carbons resonated at δ_{C} 119.6 (C-7), 144.3 (C-24), 147.4 (C-8), and 128.6 (C-25), respectively. Seven sp^3 methyl carbons resonated at δ_{C} 24.7 (C-18), 17.1 (C-19), 18.4 (C-21), 12.6 (C-27), 29.5 (C-28), 23.7 (C-29), and 30.8 (C-30). Nine methylene carbons resonated at δ_{C} 35.5 (C-1), 28.1 (C-2), 23.1 (C-6), 22.8 (C-11), 33.4 (C-12), 34.7 (C-15), 28.5 (C-16), 35.0 (C-22), and 25.5 (C-23). Five sp^3 methines resonated at δ_{C} 77.1 (C-3), 48.5 (C-5), 48.4 (C-9), 53.5 (C-17), and 35.9 (C-20). Five sp^3 quaternary carbon signals resonated at δ_{C} 38.8 (C-4), 35.8 (C-10), 43.5 (C-13), and 52.7 (C-14) while one carbonyl group resonated at δ_{C} 171.8 (C-26)




Fig 3. Chemical structure of 3 β -hydroxy-24-ethyl-5,22-cholestadiene (**1**) and 3 β -hydroxy-9-lanosta-7,24 E -dien-26-oic acid (**2**)

was assigned as a carboxylic acid group. The presence of seven sp^3 methyls, one carboxylic acid group, and nine sp^3 methylene carbons were assigned as a triterpenoid tetracyclic similar to a lanostane derivative skeleton [20-21]. The position of the tetracyclic systems (A, B, C and D) were assigned by the correlation between proton and carbon using HMBC and ^1H - ^1H COSY spectra. The HMBC correlations between six sp^3 methyl carbons (C-18, C-19, C-21, C-27, C-28, C-29, and C-30) and six sp^3 quaternary carbon C-13, C-10, C-20, C-4, C-4, and C-14, respectively, confirmed the tetracyclic systems. The position of the hydroxy group at C-3 was confirmed by the correlation between sp^3 oxygenated methine carbon at δ_c 77.1 and sp^3 methyl carbons at δ_c 29.5 (C-28) and 23.7 (C-29). The position of an olefin group at C-7/C-8 was confirmed by HMBC correlations between H-5, H-6, H-9 to C-7 and H-6, H-9, H-11 to C-8. Additional correlation between H-24 to C-25 and C-27; H-27 to C-24, C-25, and C-26 (δ_c 171.8), suggested that a carboxylic acid group is not located in the tetracyclic systems. Another olefin group was confirmed at C-24/C-25 by HMBC correlations between H-23 to C-24 and H-27 to C-25. The side chain of **2** was further confirmed by a continuous sequence from C-15 to C-24 as deduced from HMBC and ^1H - ^1H COSY spectra. In addition, the HMBC correlation of H-21 to C-20 indicated the side chain is connected to C-20. Therefore, isolated compound **2** was

identified as a 3β -hydroxy-9-lanosta-7,24E-dien-26-oic acid. The isolated compound **2** was first reported from this plant. All compounds were identified, as reported previously [22-23].

Compounds **1-2** were evaluated for their ability to suppress parasitemia and inhibit the parasite growth. The *in vitro* antiparasmodial activity of **1-2** and chloroquine as a positive control are reported in Fig. 2(d-e). Steroid **1** exhibited antiparasmodial activity against Pf D-10 with an IC_{50} value of 7.35 $\mu\text{g/mL}$. Interestingly, triterpenoid **2** displayed a more pronounced antiparasmodial activity (IC_{50} , 5.30 $\mu\text{g/mL}$) than chloroquine (IC_{50} , 7.84 $\mu\text{g/mL}$). The IC_{50} value of chloroquine is in good agreement with Perumal et al. [23].

Next, we take a look at the parasitemia as well as the inhibition rates of **1-2**. In the same way, as was the case with the PP extract, the antiparasmodial activity of compounds **1** and **2** increased in a manner that was dependent on the concentration. For **1**, the prevalence of parasitemia ranged from roughly 4.6 to 1.1%, and for **2**, the prevalence ranged from approximately 3.7 to 0.7% within a range of 0.78 to 100 $\mu\text{g/mL}$, respectively (Fig. 4(a)). Fig. 4(b) demonstrates that the growth of the parasite Pf D-10 was stifled as a result of exposure to **1-2**. Under the same conditions, steroid **1** showed a moderate suppression of Pf D-10 at a concentration of




Fig 4. An *in vitro* antiparasmodial activity of steroid **1** and triterpenoid **2**. (a) Parasitemia and (b) inhibition rates of untreated parasites (0.2% DMSO) and parasites treated with compounds **1-2** at a concentration of 0.78 to 100 $\mu\text{g/mL}$ and chloroquine (1 μM) are shown. All experiments were performed in triplicate ($n = 3$). The standard Deviation (SD) is indicated by the error bars. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and not significant (ns)

0.78 g/mL (1.7%) and 1.56 g/mL (4.0%), but triterpenoid **2** showed a considerable suppression impact of 14% and 25% ($p < 0.05$). In addition, the ability of compounds **1** and **2** to preserve the life of parasitized red blood cells was investigated. Steroid **1** had a moderate effect (about 96%) when tested at a diluted quantity, whereas triterpenoid **2** demonstrated superior parasite killing at a rate of 86% when tested at the same doses. The survival rates dropped in a manner that was dependent on the concentration of the organisms (Fig. S2).

The substantial response of triterpenoid **2** against Pf D-10 was mediated by a carboxyl functional group, which promoted carboxyl-carboxylate association for the

purpose of providing a stabilizing force in protein interaction [24]. Contrary to compound **2**, a low response of **1** is most likely caused by the compound's cytotoxicity rather than particular action against the parasite itself or being negatively influenced by poor pharmacokinetics [25-27]. This is because compound **1** has poor pharmacokinetics. The antiplasmodial activity of compounds **1-2** supported the hypothesis that the species PP L. Kunth could serve as a possible source of novel antimalarial metabolites. This allowed for the identification of potential antiplasmodial leads derived from PP L. Kunth, which could lead to the discovery of a new anti-malarial candidate.




Fig 5. *In vivo* test of triterpenoid **2** in BALB/c albino mice infected with *Pb ANKA*. (a) *In vivo* four-day suppressive test. Infected mice were treated with 0.5% CMC-Na as a negative control (NC), CQ as a positive control (25 mg/kg/body) and triterpenoid **2** at a daily dose of 1, 10 and 100 mg/kg/body. Mice received treatment from day 1 to day 4. On the 5th day, Giemsa-stained blood smears were prepared and analyzed microscopically. (b) ED₅₀ value of triterpenoid **2** in mice. (c) Parasitemia after the fourth day of treatment. (d) Chemo suppression of parasitemia from day 0 to day 4. (e) Suppression rates after the fourth day of treatment. (f) Survival rates of *Pb ANKA* in BALB/c albino mice. SD is indicated by the error bars. *** $p < 0.001$, **** $p < 0.0001$

Triterpenoid **2** proved to be the most powerful antimalarial agent in this investigation, so the next thing we did was test it on mice infected with the malaria parasite Pb ANKA. The method described by Peters et al. [26] was implemented for the purpose of analyzing the *in vivo* test of **2**. Mice of the BALB/c albino strain were given Pb ANKA by the intraperitoneal route. After 4 d of treatment with 0.5% CMC-Na as a negative control, chloroquine diphosphate as a positive control, and triterpenoid **2** at daily doses of 1, 10, and 100 mg/kg/body, thin blood smears stained with Giemsa were obtained for each mouse on the fifth day. This was followed by 4 d of treatment with chloroquine diphosphate as a positive control. On a microscopic level, parasitemia and the reduction of parasite development were observed and measured (Fig. 5(a)). Up until day 21, the survival of mice was carefully monitored and tracked. It was believed that triterpenoid **2** had some biological activity, with an ED₅₀ value of 18.51 mg/kg/body weight (Fig. 5(b)). The drug quinine, which is available for clinical use, has an elimination half-life (ED₅₀) of 34 mg/kg/day and a slow clearance [28]. Triterpenoid **2** was successful in inhibiting the development of parasitemia in mice infected with Pb ANKA, as shown in Fig. 5(c). Even at the lowest dose (1 mg/kg/body), intraperitoneal administration of triterpenoid **2** resulted in a 78% reduction in parasitemia. To be more specific, **2** was successful in killing 22% of the parasites. The incidence of parasitemia fell to lower levels as higher doses of **2** were administered. The rates of parasitemia were reduced to approximately 54 and 34%, respectively when triterpenoid **2** was administered at doses of 10 and 100 mg/kg/body. In addition, the effectiveness of chemotherapy in reducing parasitemia was evaluated from day 0 to day 4 of treatment (Fig. 5(d)). The levels of parasitemia in mice that had not been given any treatment rose consistently after infection, but the number **2** therapy had a significant chemo-suppressive effect that was dose-dependent. Fig. 5(e) shows the suppression of parasite growth that was caused by triterpenoid **2**. This compound had a substantial inhibitory effect against Pb ANKA in mice. Triterpenoid **2** strongly suppressed Pb ANKA, as can be seen in Fig. 5(e), which suggests that it also inhibited the proliferation of parasites

in mice. The suppression rates varied from 66 (at 100 mg/kg/body) to 46 (10 mg/kg/body) to 22% (1 mg/kg/body). These findings made it abundantly evident that factor **2** was responsible for eliminating the parasites. The treatment of **2** also increased the number of ill mice that survived longer. Based on the *in vivo* experiment (Fig. 5(f)) mice that were not given any treatment succumbed to the Pb ANKA infection after 10 d, whereas mice that were given 1, 10, or 100 mg/kg/body of triterpenoid **2** lived until 12, 17, and 21 d, respectively. Positive control groups were given chloroquine diphosphate at a daily dose of 25 mg/kg/body for as long as 21 d, which led researchers to conclude that the Pb ANKA infection was completely eradicated in the mice who participated in the study. These findings are in agreement with those found by Fang et al. [29].

■ CONCLUSION

In this study, the PP extract displayed a more pronounced antiplasmodial activity than the AP extract. Steroid **1** and triterpenoid **2** exhibited promising antiplasmodial activity. This is the first report of 3 β -hydroxy-9-lanosta-7,24*E*-dien-26-oic acid (**2**) as an antiplasmodial agent, and we have shown its promising efficacy in an *in vivo* mouse model. Our results suggest a novel triterpenoid that has the potential for use as an antimalarial agent. Although 3 β -hydroxy-9-lanosta-7,24*E*-dien-26-oic acid may not yet be an ideal drug candidate for treating malaria disease in humans, it may provide a useful lead structure to produce more effective antiplasmodial agents. We plan to do an *in vivo* test with the combination of triterpenoid **2** and antimalarial drug artesunate to improve the efficacy of drug action [30].

■ ACKNOWLEDGMENTS

We thank Dr. Sofa Fajriah in Research Center for Chemistry, Indonesian Science Institute for NMR measurements and Mrs. Hilkatul Ilmi, Institute of Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia, for the technical support provided during the *in vivo* studies. We are grateful to Dr. Esther M. Yoo, Mr. Andrew Miller and Dr. Chandradhish

Ghosh at the Max Planck Institute for their helpful discussions.

AUTHOR CONTRIBUTIONS

The experiment was conducted by NB with the analysis being conducted by MAM and YS. The NMR and *in vivo* analysis were done by WM and AW. The manuscript was arranged and revised by AMI, BS, and JS. All the authors mentioned have agreed to the final version of this manuscript.

REFERENCES

- [1] World Health Organization, 2019, *World malaria report 2019*, World Health Organization, Geneva, <https://apps.who.int/iris/handle/10665/330011>, License: CC BY-NC-SA 3.0 IGO
- [2] Menard, D., and Dondorp, A., 2017, Antimalarial drug resistance: A threat to malaria elimination, *Cold Spring Harbor Perspect. Med.*, 7, a025619.
- [3] Diagana, T.T., 2015, Supporting malaria elimination with 21st century antimalarial agent drug discovery, *Drug Discovery Today*, 20 (10), 1265–1270.
- [4] Cowell, A.N., and Winzeler, E.A., 2019, The genomic architecture of antimalarial drug resistance, *Briefings Funct. Genomics*, 18 (5), 314–328.
- [5] Babatunde, K.A., and Adenuga, O.F., 2022, Neutrophils in malaria: A double edged sword role, *Front. Immunol.*, 13, 922377.
- [6] Tu, Y., 2011, The discovery of artemisinin (*qinghaosu*) and gifts from Chinese medicine, *Nat. Med.*, 17 (10), 1217–1220.
- [7] Gómez-García, A., and Medina-Franco, J.L., 2022, Progress and impact of Latin American natural product databases, *Biomolecules*, 12 (9), 1202.
- [8] Hedrén, M., Chase, M.W., and Olmstead, R.G., 1995, Relationships in the *Acanthaceae* and related families as suggested by cladistic analysis of *rbcl* nucleotide sequences, *Plant Syst. Evol.*, 194 (1), 93–109.
- [9] Dirar, A.I., Adhikari-Devkota, A., Kunwar, R.M., Paudel, K.R., Belwal, T., Gupta, G., Chellapan, D.K., Hansbro, P.M., Dua, K., and Devkota, H.P., 2021, Genus *Blepharis* (Acanthaceae): A review of ethnomedicinally used species, and their phytochemistry and pharmacological activities, *J. Ethnopharmacol.*, 265, 113255.
- [10] Wahidah, B.F., and Husain, F., 2018, Etnobotani tumbuhan obat yang dimanfaatkan oleh masyarakat Desa Samata Kecamatan Somba Opu Kabupaten Gowa Sulawesi Selatan, *Life Sci.*, 7 (2), 39–88.
- [11] Widyawaruyanti, A., Asrory, M., Ekasari, W., Setiawan, D., Radjaram, A., Tumewu, L., and Hafid, A.F., 2014, *In vivo* antimalarial activity of *Andrographis paniculata* tablets, *Procedia Chem.*, 13, 101–104.
- [12] Sari, N., Wahidah, B.F., and Gaffar, N.A., 2017, Etnobotani tumbuhan yang digunakan dalam pengobatan tradisional di Kecamatan Sinjai Selatan Kabupaten Sinjai Sulawesi Selatan, *Pros. Sem. Nas. Biol. Life*, 3 (1), 6–13.
- [13] Dalimartha, S., and Adrian, F., 2013, *Ramuan Herbal Tumpas Penyakit*, Penebar Swadaya, Jakarta, Indonesia.
- [14] Ekasari, W., Fatmawati, D., Khoiriah, S.M., Baquiddin, W.A., Nisa, H.Q., Maharupini, A., Wahyuni, T.S., Oktarina, R.D., Suhartono, E., and Sahu, R.K., 2022, Antimalarial activity of extract and fractions of *Sauropus androgynus* (L.) Merr, *Scientifica*, 2022, 3552491.
- [15] Giemsa, G., 1902, Färbemethoden für malariaparasiten, *Zentralbl. Bakteriol.*, 31, 429–430.
- [16] McLean, R.C., 1943, Microscopic technique in biology and medicine, *Nature*, 152 (3868), 709.
- [17] Kwansa-Bentum, B., Agyeman, K., Larbi-Akor, J., Anyigba, C., and Appiah-Opong, R., 2019, *In vitro* assessment of antiplasmodial activity and cytotoxicity of *Polyalthia longifolia* leaf extracts on *Plasmodium falciparum* strain NF54, *Malar. Res. Treat.*, 2019, 6976298.
- [18] Kamaraj, C., Kaushik, N.K., Mohanakrishnan, D., Elango, G., Bagavan, A., Zahir, A.A., Rahuman, A.A., and Sahal, D., 2012, Antiplasmodial potential of medicinal plant extracts from Malaiyur and Javadhu hills of South India, *Parasitol. Res.*, 111 (2), 703–715.
- [19] Greca, M.D., Monaco, P., and Previtera, L., 1990, Stigmasterols from *Typha latifolia*, *J. Nat. Prod.*, 53 (6), 1430–1435.

- [20] Wang, G.W., Lv, C., Yuan, X., Ye, J., Jin, H.Z., Shan, L., Xu, X.K., Shen, Y.H., and Zhang, W.D., 2015, Lanostane-type triterpenoids from *Abies faxoniana* and their DNA topoisomerase inhibitory activities, *Phytochemistry*, 116, 221–229.
- [21] Shen, J., Kai, J., Tang, Y., Zhang, L., Su, S., and Duan, J.A., 2016, The chemical and biological properties of *Euphorbia kansui*, *Am. J. Chin. Med.*, 44 (2), 253–273.
- [22] Isaka, M., Chinthanom, P., Sappan, M., Supothina, S., Vichai, V., Danwisetkanjana, K., Boonpratuang, T., Hyde, K.D., and Choeyklin, R., 2017, Antitubercular activity of mycelium-associated *Ganoderma* lanostanoids, *J. Nat. Prod.*, 80 (5), 1361–1369.
- [23] Perumal, P., Sowmiya, R., Kumar, S.P., Ravikumar, S., Deepak, P., and Balasubramani, G., 2018, Isolation, structural elucidation and antiplasmodial activity of fucosterol compound from brown seaweed, *Sargassum linearifolium* against malarial parasite *Plasmodium falciparum*, *Nat. Prod. Res.*, 32 (11), 1316–1319.
- [24] Workman, S.D., Worrall, L.J., and Strynadka, N.C.J., 2018, Crystal structure of an intramembranal phosphatase central to bacterial cell-wall peptidoglycan biosynthesis and lipid recycling, *Nat. Commun.*, 9 (1), 1159.
- [25] Fu, Y., Ding, Y., Wang, Q., Zhu, F., Tan, Y., Lu, X., Guo, B., Zhang, Q., Cao, Y., Liu, T., Cui, L., and Xu, W., 2020, Blood-stage malaria parasites manipulate host innate immune responses through the induction of sFGL2, *Sci. Adv.*, 6 (9), eaay9269.
- [26] Peters, W., Portus, J.H., and Robinson, B.L., 1975, The chemotherapy of rodent malaria, XXII. The value of drug-resistant strain of *P. berghei* in screening for blood schizontocidal activity, *Ann. Trop. Med. Parasitol.*, 69 (2), 155–171.
- [27] Sianturi, J., Manabe, Y., Li, H.S., Chiu, L.T., Chang, T.C., Tokunaga, K., Kabayama, K., Tanemura, M., Takamatsu, S., Miyoshi, E., Hung, S.C., and Fukase, K., 2019, Development of α -gal-antibody conjugates to increase immune response by recruiting natural antibodies, *Angew. Chem., Int. Ed.*, 58 (14), 4526–4530.
- [28] Jiménez-Díaz, M.B., Viera, S., Ibáñez, J., Mulet, T., Magán-Marchal, N., Garuti, H., Gomez, V., Cortés-Gil, L., Martínez, A., Ferrer, S., Fraile, M.T., Calderón, F., Fernández, E., Shultz, L.D., Leroy, D., Wilson, D.M., García-Bustos, J.F., Gamo, F.J., and Angulo-Barturen, I., 2013, A new *in vivo* screening paradigm to accelerate antimalarial drug discovery, *PLoS One*, 8 (6), e66967.
- [29] Fang, Y., He, X., Zhang, P., Shen, C., Mwangi, J., Xu, C., Mo, G., Lai, R., and Zhang, Z., 2019, *In vitro* and *in vivo* antimalarial activity of LZ1, a peptide derived from snake cathelicidin, *Toxins*, 11 (7), 379.
- [30] Corral, M.G., Leroux, J., Stubbs, K.A., and Mylne, J.S., 2017, Herbicidal properties of antimalarial drugs, *Sci. Rep.*, 7 (1), 45871.

Evaluation of the Antiplasmodial Properties of *Andrographis paniculata* (Burm.f.) and *Peperomia pellucida* (L.) Kunth

ORIGINALITY REPORT

14%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Lukman Abdul Rauf Laliyo, Bambang Sumintono, Citra Panigoro. "Measuring Changes in Hydrolysis Concept of Students Taught by Inquiry Model: Stacking and Racking Analysis Techniques in Rasch Model", *Heliyon*, 2022
Publication

1 %
- 2

www.ncbi.nlm.nih.gov
Internet Source

1 %
- 3

Ali Ramazani. "In vitro and in vivo anti-malarial activity of *Boerhavia elegans* and *Solanum surattense*", *Malaria Journal*, 2010
Publication

<1 %
- 4

Dewa Gede Katja, Kindi Farabi, Desi Harneti, Tri Mayanti, Unang Supratman. "Cytotoxic Triterpenoid from the Stembark of *Chisocheton celebicus* (Meliaceae)", *Makara Journal of Science*, 2017
Publication

<1 %

5

Masahiko Isaka, Panida Chinthanom, Malipan Sappan, Sumalee Supothina et al. "Antitubercular Activity of Mycelium-Associated Lanostanoids ", Journal of Natural Products, 2017

Publication

<1 %

6

www.jurnal.ugm.ac.id

Internet Source

<1 %

7

ndl.ethernet.edu.et

Internet Source

<1 %

8

Panida Chinthanom, Kitlada Srichomthong, Pranee Rachtawee, Thitiya Boonpratuang et al. "Lanostane triterpenoids from cultivated fruiting bodies of Ganoderma sichuanense: Determination of the C-25 absolute configuration of ganoderic acid A and its derivatives using the phenylglycine methyl ester (PGME) method", Phytochemistry, 2021

Publication

<1 %

9

Cedric Derick Jiatsa Mbouna, Brice Mariscal Tchatat Tali, Patrick Valere Tsouh Fokou, Eugenie Aimee Madiesse Kemgne et al. "Specific sub fractions from Terminalia mantaly (H. Perrier) extracts potently inhibit Plasmodium falciparum rings, merozoite egress and invasion", Journal of Ethnopharmacology, 2021

Publication

<1 %

10 Desi Harneti, Asep Supriadin, Mariam Ulfah, Agus Safari, Unang Supratman, Khalijah Awang, Hideo Hayashi. "Cytotoxic constituents from the bark of *Aglaia eximia* (Meliaceae)", *Phytochemistry Letters*, 2014
Publication

11 www.scribd.com
Internet Source

12 Ahmed M. Abdou, Abdel-latif S. Seddek, Noha Abdelmageed, Mohamed O. Badry, Yoshifumi Nishikawa. "Wild Egyptian medicinal plants show in vitro and in vivo cytotoxicity and antimalarial activities", *BMC Complementary Medicine and Therapies*, 2022
Publication

13 He-Ping Chen, Zhen-Zhu Zhao, Zheng-Hui Li, Ying Huang et al. " Anti-Proliferative and Anti-Inflammatory Lanostane Triterpenoids from the Polish Edible Mushroom ", *Journal of Agricultural and Food Chemistry*, 2018
Publication

14 Masahiko Isaka, Panida Chinthanom, Vanicha Vichai, Sujinda Sommai, Rattaket Choeyklin. " Ganoweberianones A and B, Antimalarial Lanostane Dimers from Cultivated Fruiting Bodies of the Basidiomycete ", *Journal of Natural Products*, 2020
Publication

15

Ping-Chung Kuo. "Eurycomalin A, a new dimeric dihydrobenzofuran from *Eurycoma longifolia*", *Heterocycles*, 20040901

Publication

<1 %

16

Guo-Wei Wang, Chao Lv, Xing Yuan, Ji Ye, Hui-Zi Jin, Lei Shan, Xi-Ke Xu, Yun-Heng Shen, Wei-Dong Zhang. "Lanostane-type triterpenoids from *Abies faxoniana* and their DNA topoisomerase inhibitory activities", *Phytochemistry*, 2015

Publication

<1 %

17

Shen, De-Yang, Yu-Yi Chan, Tsong-Long Hwang, Shin-Hun Juang, Shioh-Chyn Huang, Ping-Chung Kuo, Tran Dinh Thang, E-Jian Lee, Amooru G. Damu, and Tian-Shung Wu. "Constituents of the Roots of *Clausena lansium* and Their Potential Anti-inflammatory Activity", *Journal of Natural Products*, 2014.

Publication

<1 %

18

www.ntno.org

Internet Source

<1 %

19

Junping Chen, Yichen Li, Jianguo Cao, Jinwen Huang, Can Jiang, Xiling Dai, Guozheng Huang. "Adiantic acid, a new unsaturated fatty acid with a cyclopropane moiety from *L.* ", *Natural Product Research*, 2020

Publication

<1 %

20	acikbilim.yok.gov.tr Internet Source	<1 %
21	www.thieme-connect.com Internet Source	<1 %
22	Nguyen Xuan Nhiem, Phan Van Kiem, Chau Van Minh, Jung-Jin Lee, Ja Hyeong Ku, Chang-Seon Myung, Young Ho Kim. "A potential inhibitor of rat aortic vascular smooth muscle cell proliferation from the pollen of <i>Typha angustata</i> ", Archives of Pharmacal Research, 2010 Publication	<1 %
23	dadun.unav.edu Internet Source	<1 %
24	ejnmmites.springeropen.com Internet Source	<1 %
25	kimia.unjani.ac.id Internet Source	<1 %
26	smujo.id Internet Source	<1 %
27	www.asianpubs.org Internet Source	<1 %
28	www.biorxiv.org Internet Source	<1 %

29	Submitted to Ho Chi Minh University of Technology and Education	<1 %
Student Paper		

30	Submitted to The Robert Gordon University	<1 %
Student Paper		

31	nepis.epa.gov	<1 %
Internet Source		

32	Prajakta P. Dandekar. "Curcumin-loaded hydrogel nanoparticles: Application in anti-malarial therapy and toxicological evaluation", Journal of Pharmaceutical Sciences, 2010	<1 %
Publication		

33	Purwani, Ni Nyoman, Handoko Darmokoesoemo, and Ni Nyoman Tri Puspaningsih. "Hydrolysis of Corncob Xylan using β -xylosidase GbtXyl43B from Geobacillus Thermoleovorans IT-08 Containing Carbohydrate Binding Module (CBM)", Procedia Chemistry, 2016.	<1 %
Publication		

34	patents.google.com	<1 %
Internet Source		

35	pt.scribd.com	<1 %
Internet Source		

36	fenix.tecnico.ulisboa.pt	<1 %
Internet Source		

Desi Harneti, Roekmiati Tjokronegoro, Agus Safari, Unang Supratman et al. "Cytotoxic triterpenoids from the bark of *Aglaia smithii* (Meliaceae)", *Phytochemistry Letters*, 2012

Publication

Hai-Guo Su, Qin-Mei Zhou, Li Guo, Yuan-Jin Huang, Cheng Peng, Liang Xiong. "Lanostane triterpenoids from *Ganoderma luteomarginatum* and their cytotoxicity against four human cancer cell lines", *Phytochemistry*, 2018

Publication

Hiroki Kumamoto, Sayoko Onuma, Kumiko Tsuchiya, Yuko Egusa, Hiromichi Tanaka, Tsuyoshi Satoh. "SULFOXIDE-METAL EXCHANGE FOR THE SYNTHESIS OF THE 2'-TRIBUTYLSTANNYL DERIVATIVE OF 2',3'-DIDEHYDRO-2',3'-DIDEOXYURIDINE (d4U): A GENERAL ENTRY TO 2'-CARBON-SUBSTITUTED ANALOGUES OF d4U", *Nucleosides, Nucleotides and Nucleic Acids*, 2002

Publication

Widyawaruyanti, Aty, Muhammad Asrory, Wiwied Ekasari, Dwi Setiawan, Achmad Radjaram, Lidya Tumewu, and Achmad Fuad Hafid. "In vivo Antimalarial Activity of

Andrographis Paniculata Tablets", Procedia Chemistry, 2014.

Publication

42

par.nsf.gov

Internet Source

<1 %

43

www.researchgate.net

Internet Source

<1 %

44

"Ethnobotany of the Mountain Regions of Southeast Asia", Springer Science and Business Media LLC, 2021

Publication

<1 %

45

"Treated Plasmodium berghei infected pregnant mice by Andrographis paniculata tablet (AS201-01) decreasing the TLR-4 expression and apoptosis index of placental tissue", Journal of Applied Pharmaceutical Science, 2018

Publication

<1 %

46

Anke Harupa, Laura De Las Heras, Gonzalo Colmenarejo, Sally Lyons-Abbott et al. " Identification of Selective Inhibitors of N-Myristoyltransferase by High-Throughput Screening ", Journal of Medicinal Chemistry, 2019

Publication

<1 %

47

Chinnaperumal Kamaraj, Abdul Abdul Rahuman, Selvaraj Mohana Roopan, Asokan

<1 %

Bagavan et al. "Bioassay-guided isolation and characterization of active antiplasmodial compounds from *Murraya koenigii* extracts against *Plasmodium falciparum* and *Plasmodium berghei*", Parasitology Research, 2014

Publication

48

Gaurav Sharma, Devika Rana, Sandeep Sundriyal, Ankusha Sharma, Pankaj Panwar, Neeraj Mahindroo. "Plants from Annonaceae family as antimalarials: An ethnopharmacology and phytochemistry review to identify potential lead molecules", South African Journal of Botany, 2023

Publication

<1 %

49

Muthaura, C.N.. "Antimalarial activity of some plants traditionally used in treatment of malaria in Kwale district of Kenya", Journal of Ethnopharmacology, 20070725

Publication

<1 %

50

repositorio.ufmg.br

Internet Source

<1 %

51

researchspace.ukzn.ac.za

Internet Source

<1 %

52

scholarworks.utep.edu

Internet Source

<1 %

53

www.degruyter.com

Internet Source

<1 %

54

www.jimmunol.org

Internet Source

<1 %

55

www.science.gov

Internet Source

<1 %

56

S BoseDasgupta, B B Das, S Sengupta, A Ganguly, A Roy, S Dey, G Tripathi, B Dinda, H K Majumder. "The caspase-independent algorithm of programmed cell death in Leishmania induced by baicalein: the role of LdEndoG, LdFEN-1 and LdTatD as a DNA 'degradesome'", Cell Death & Differentiation, 2008

Publication

<1 %

57

Wijayanti, Mahardika Agus, Eti Nurwening Sholikhah, Iqmal Tahir, Ruslin Hadanu, Jumina, Supargiyono, and Mustofa. "Antiplasmodial Activity and Acute Toxicity of N-alkyl and N-benzyl-1,10-Phenanthroline Derivatives in Mouse Malaria Model", JOURNAL OF HEALTH SCIENCE, 2006.

Publication

<1 %

58

Bastiana, Aty Widyawaruyanti, Hilkatul Ilmi, Lidya Tumewu, Budi Prasetyo, Achmad Fuad Hafid, Aryati. " A tablet derived from complements dihydroartemisinin-piperaquine

<1 %

treatment of malaria in pregnant mice ",
Journal of Basic and Clinical Physiology and
Pharmacology, 2021

Publication

59

Guo-Wei Wang, Chao Lv, Yun-Heng Shen, Wei-
Dong Zhang. " Triterpenoids from and their
cytotoxic activities ", Natural Product
Research, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Evaluation of the Antiplasmodial Properties of Andrographis paniculata (Burm.f.) and Peperomia pellucida (L.) Kunth

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11