



Search this journal...



Submit manuscript

My tasks

About

Articles

Issues

Author Guidelines

Editorial Team

Contacts

My tasks

Pending

My manuscripts

Active	0
Published	1
Rejected	3
Incomplete	1
Uploaded	0

Submission #96817

Current status

Metadata

Suppl. files (0)

Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds

Weny J. A. Musa ✉, Nurhayati Bialangi ✉, Ahmad Kadir Kilo ✉, Boima Situmeang ✉, Ninik Triayu Susparini

Article type: *Research Article* Type of review: Conventional peer review

Subject e

Review round 1



Review version

Manuscript file

Show more files



Manuscript files

WORD file

Reviewer(s)

Reviewer 1

Minor revision

4 Nov 2022

Review:

it is advisable to indicate which biologically active compounds are determined in fractions, indicate their chemical and qualitative composition, indicate the possible mechanism of antioxidant action

[show more](#)

Reviewer 2

Minor revision

27 Nov 2022

Review:

Dear Editor, The manuscript entitled "Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract" could be published in Pharmacia. The manuscript...

[show more](#)

Editorial decision

This website uses cookies in order to improve your web experience. [Read our Cookies](#)

Rumiana Simeonova

[Policy](#) [revision](#)

Review:

OK

29 Nov 2022

DEAR AUTHORS, PLEASE CONSIDER REVIEWERS' COMMENTS AND SUGGESTIONS!

[show more](#)

Review round 2



Author's revised files

[Revised figure 4, revised chemical name, revised discussion and bibliography](#)

Author comments:

Reviewer 1: It is advisable to indicate which biologically active compounds are determined in fractions, indicate their chemical composition, indicate the possible mechanism of antioxidant action Answer: Thank you for the advice and input, the biologist refers...

[show more](#)

Editorial decision

Rumiana Simeonova

Accept

1 Dec 2022

Review:

Dear Authors, Thank you for considering the reviewers' comments

[show more](#)

Copy editing 1



Author's revised files

Edit

[Revised figure 4, revised chemical name, revised discussion and bibliography](#)

Copy editor

Georgi Momekov

17 Jan 2023



Please find enclosed the copy edited version

[show more](#)

Layout editor 1



Author's revised files

Edit

[text](#)

Layout editor

Polina Petrakieva

20 Jan 2023



[show more](#)

Layout editor 2

**Author's revised files**[Edit](#)[Revised figure citation in-text](#)

Author comments:

Dear Editor, Thanks you for the corrections. We have done the necessary changes according to what suggested and for the error. The in-text citation has been added to the proofing but we apologize for the messy layout. The co-authors information is complete...

[show more](#)

Layout editor


Polina Petrakieva

23 Jan 2023





Dear Authors, I've placed the citations of figures in the text, but if you have more corrections send them in the usual way, using the comment function in the pdf file (or as an alternative, as a...)

[show more](#)

Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extractWeny J. A. Musa , Nurhayati Bialangi , Ahmad Kadir Kilo , **Boima Situmeang** , Ninik Triayu Susparini , Ilham Dwi Rusydi Article type: *Research Article* Type of review:  Conventional peer reviewSubject editor: Rumiana Simeonova **Review round 1****Review version**[Manuscript file](#)**Manuscript files**[WORD file](#)[Cover letter](#)[Download all files
as zip archive](#)[Show fewer files](#) 

Reviewer(s)

Reviewer 1	Minor revision 4 Nov 2022	Review:  it is advisable to indicate which biologically active compounds are determined in fractions, indicate their chemical and qualitative composition, indicate the possible mechanism of antioxidant action show less	No files uploaded
Reviewer 2	Minor revision 27 Nov 2022	Review:  Dear Editor, The manuscript entitled "Antioxidant, Cholesterol Lowering Activity, and Analysis of <i>Caesalpinia bonducella</i> Seeds Extract" could be published in <i>Pharmacia</i> . The manuscript seems suitable for publication according to the reviewer, only few points may require slight clarification. Comments: Please use H ₂ SO ₄ instead of H2SO4, LC/MS instead of LCMS. In depth discussion, concerning comparison to previous <i>Caesalpinia</i> research as well as the contribution of the secondary metabolites to the biological activity of the extracts is needed. show less	No files uploaded

Editorial decision

Rumiana Simeonova	Minor revision 29 Nov 2022	Review:  DEAR AUTHORS, PLEASE CONSIDER REVIEWERS' COMMENTS AND SUGGESTIONS! show less	No files uploaded
-------------------	-------------------------------	--	-------------------

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Chairunnisa Lamanganjo¹, Suleman Duengo², Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini²

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

Corresponding author: Boima Situmeang (boimatumeang@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction were made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction were analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al, 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al, 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, SriLanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili seed showed that tombili seed contains saponins, phytosterols,

flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals, thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant

department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for extraction and fractionation. The 2,2- diphenyl-1-

picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g). Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LCMS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (*n*= 3). The inhibition percentage (%) was computed as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influenced by several factors, including the chemical screening of phytochemicals, the partition method used,

the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extraction yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also been seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LCMS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088		
	25	9.227	10.041	9.803	100.798	
	30	12.409	12.008	11.764	99.505	100.933±1.4995
	35	14.994	15.010	14.860	102.495	
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832		
	25	8.299	10.972	9.420	88.499	
	30	12.348	13.474	14.078	81.286	86.153±4.2161
	35	14.372	16.554	17.184	88.675	
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617		
	25	12.938	9.227	10.084	93.551	
	30	16.248	12.409	11.974	95.890	94.053±1.6445
	35	18.115	14.994	15.336	92.718	
	40	22.066	17.786	18.907		


**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpan-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the invitro DPPH radical scavenging ability of tomboli seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical

scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction.

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (n = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600 ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen bonding. Unlike ethyl acetate, the

intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	t _R (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C17H16N2O2	Identified	281.1275	280.12118	9.89
5	Candidate Mass C26H38O7	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C17H16N2O2	Identified	515704		+H
5	Candidate Mass C26H38O7	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C17H16N2O2	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C26H38O7	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 4. The profiles of the identified five major compounds measured with LCMS apparatus.




Figure 5. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3 β - glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LCMS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LCMS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed

was first reported. The highest antioxidant activity was shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LCMS/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety.* 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/artic le/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbrn.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties.* 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia.* 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitarty N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Editor-in-Chief

Pharmacia Journal

Dear Editorial Team,

Enclosed please find the manuscript entitled “**Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seed extract**” to be considered for publication in Pharmacia Journal.

The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extracts were fractionated using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical contents of the ethyl acetate fraction were analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. Adding cholesterol solution absorbance was employed to determine cholesterol levels of extract and fractions. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol salt. The highest anti-cholesterol shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % cholesterol salt. The LC-MS/MS determination showed that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

We would like to declare that our work is original. The results presented in this manuscript, in whole or in any part, have not been published for publication elsewhere. All authors listed have approved the enclosed manuscript. The authors declare no conflicts of interest associated with this manuscript.

Thank you for your consideration of this manuscript.

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Chairunnisa Lamanganjo¹, Suleman Duengo², Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini²

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

Corresponding author: Boima Situmeang (boimatumeang@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction were made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction were analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al, 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al, 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, SriLanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili seed showed that tombili seed contains saponins, phytosterols,

flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals, thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant

department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for extraction and fractionation. The 2,2- diphenyl-1-

picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g). Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LCMS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (*n*= 3). The inhibition percentage (%) was computed as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influenced by several factors, including the chemical screening of phytochemicals, the partition method used,

the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration. This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LCMS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088		
	25	9.227	10.041	9.803	100.798	
	30	12.409	12.008	11.764	99.505	100.933±1.4995
	35	14.994	15.010	14.860	102.495	
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832		
	25	8.299	10.972	9.420	88.499	
	30	12.348	13.474	14.078	81.286	86.153±4.2161
	35	14.372	16.554	17.184	88.675	
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617		
	25	12.938	9.227	10.084	93.551	
	30	16.248	12.409	11.974	95.890	94.053±1.6445
	35	18.115	14.994	15.336	92.718	
	40	22.066	17.786	18.907		


**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpan-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the invitro DPPH radical scavenging ability of tomboli seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical

scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction.

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (n = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600 ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen bonding. Unlike ethyl acetate, the

intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	t_R (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C17H16N2O2	Identified	281.1275	280.12118	9.89
5	Candidate Mass C26H38O7	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C17H16N2O2	Identified	515704		+H
5	Candidate Mass C26H38O7	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C17H16N2O2	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C26H38O7	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 4. The profiles of the identified five major compounds measured with LCMS apparatus.




Figure 5. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3 β - glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LCMS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LCMS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed

was first reported. The highest antioxidant activity was shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LCMS/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety.* 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/artic le/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbrn.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties.* 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia.* 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitarty N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Review round 2



Author's revised files

Revised figure 4, revised chemical name, revised discussion and bibliography



Download all files
as zip archive

Author comments:

Reviewer 1: It is advisable to indicate which biologically active compounds are determined in fractions, indicate their chemical and qualitative composition, indicate the possible mechanism of antioxidant action

Answer: Thank you for the advice and input, the biologically active compound refers to the secondary metabolites, especially homoplantagin in the chemical composition of homoplantagin obtained from the fraction consists of hydrazyl and phenyl with the abundance of hydroxyl group located outside the cyclic group thus allowing it to react with the DPPH to exchange "radicals".

Reviewer 2:

Please use H₂SO₄ instead of H2SO4 and LC/MS instead of LCMS.

An in-depth discussion, concerning comparison to previous *Caesalpinia* research as well as the contribution of the secondary metabolites to the biological activity of the extracts, is needed.

Response: Thank you for the review, the necessary changes for the writing have been done and I really appreciate the reviewer pointing them out. The discussion regarding the comparison with the existing or previously done research of *Caesalpinia* has been done using the same genus of *Caesalpinia* but different species due to the plant being endemic to the region itself. The comparison results in the similarity of the most active compound extracted and tested using the same DPPH method

[show less](#)

Editorial decision

Rumiana Simeonova

Accept

1 Dec 2022

Review:

Dear Authors,

Thank you for considering the reviewers' comments

[show less](#)

No files uploaded

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis that the ethyl acetate fraction contained glucoside, homoplantagin, and verolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili

seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidant activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo

State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for

extraction and fractionation. The 2,2- diphenyl-1-picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (*n* = 3). The inhibition percentage (%) was computed

as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influ-

enced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LC/MS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088	100.798 99.505 102.495	100,933±1.4995
	25	9.227	10.041	9.803		
	30	12.409	12.008	11.764		
	35	14.994	15.010	14.860		
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832	88.499 81.286 88.675	86.153±4.2161
	25	8.299	10.972	9.420		
	30	12.348	13.474	14.078		
	35	14.372	16.554	17.184		
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617	93.551 95.890 92.718	94.053±1.6445
	25	12.938	9.227	10.084		
	30	16.248	12.409	11.974		
	35	18.115	14.994	15.336		
	40	22.066	17.786	18.907		



**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpin-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the invitro DPPH radical scavenging ability of tombili seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the

previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice.

**Figure 4.** Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one)

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times ($n = 3$) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen

bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpin-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	tr (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpin-3- β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpin-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	281.1275	280.12118	9.89
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpin-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	515704		+H
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpin-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3 β - glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LC/MS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was

shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- Badami, S., Moorkoth, S., Rai, S. R., Kannan, E., & Bhojraj, S. (2003). Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin*, 26(11), 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety*. 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/article/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbs.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties*. 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia*. 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitay N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Copy editing 1



Author's revised files



Edit

Revised figure 4, revised chemical name, revised discussion and bibliography



Download
all files as
zip archive

Copy editor

Georgi Momekov

17 Jan 2023



Please find enclosed the copy edited version

[show less](#)



DOWNLOAD
PROOF

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with an IC₅₀ value of 86.153 µg/mL. The second was the methanol fraction with an IC₅₀ value of 94.053 µg/mL, and the third was the *n*-hexane fraction with an IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis showed that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili

seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo

State University, Indonesia with the specimen being identified by Wenry JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for

extraction and fractionation. 2,2-diphenyl-1-picrylhydrazyl (DPPH) was purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (*n* = 3). The inhibition percentage (%) was computed

as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influ-

enced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LC/MS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088		
	25	9.227	10.041	9.803	100.798	
	30	12.409	12.008	11.764	99.505	100.933±1.4995
	35	14.994	15.010	14.860	102.495	
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832		
	25	8.299	10.972	9.420	88.499	
	30	12.348	13.474	14.078	81.286	86.153±4.2161
	35	14.372	16.554	17.184	88.675	
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617		
	25	12.938	9.227	10.084	93.551	
	30	16.248	12.409	11.974	95.890	94.053±1.6445
	35	18.115	14.994	15.336	92.718	
	40	22.066	17.786	18.907		



**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpan-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the in vitro DPPH radical scavenging ability of tomboli seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tomboli seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the

previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice.

**Figure 4.** Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one)

Cholesterol lowering activity of tomboli seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (n = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen

bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	tr (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	281.1275	280.12118	9.89
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	515704		+H
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3β- glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LC/MS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was

shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- Badami, S., Moorkoth, S., Rai, S. R., Kannan, E., & Bhojraj, S. (2003). Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin*, 26(11), 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety*. 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/article/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbs.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties*. 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia*. 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitay N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili

seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidant activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo

State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for

extraction and fractionation. The 2,2- diphenyl-1-picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (n= 3). The inhibition percentage (%) was computed

as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influ-

enced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LC/MS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088	100.798 99.505 102.495	100.933±1.4995
	25	9.227	10.041	9.803		
	30	12.409	12.008	11.764		
	35	14.994	15.010	14.860		
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832	88.499 81.286 88.675	86.153±4.2161
	25	8.299	10.972	9.420		
	30	12.348	13.474	14.078		
	35	14.372	16.554	17.184		
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617	93.551 95.890 92.718	94.053±1.6445
	25	12.938	9.227	10.084		
	30	16.248	12.409	11.974		
	35	18.115	14.994	15.336		
	40	22.066	17.786	18.907		



**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarp-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the invitro DPPH radical scavenging ability of tomboli seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tomboli seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the

previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice.

**Figure 4.** Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one)

Cholesterol lowering activity of tomboli seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times ($n = 3$) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen

bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	tr (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	281.1275	280.12118	9.89
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	515704		+H
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3β- glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LC/MS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was

shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- Badami, S., Moorkoth, S., Rai, S. R., Kannan, E., & Bhojraj, S. (2003). Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin*, 26(11), 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety*. 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/article/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbs.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties*. 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia*. 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitay N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Layout editor 1



Author's revised files

 Edit

text



Download
all files as
zip archive

Layout editor

Polina Petrakieva

20 Jan 2023



Dear Authors, Four of the co-authors - Nurhayati Bialangi, Ahmad Kadir Kilo, Ninik Triayu Susparini, and Ilham Dwi Rusydi - are missing from the site, but are present in the manuscript file. Can you...

[show more](#)



DOWNLOAD
PROOF

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with IC₅₀ value of 86.153 µg/mL. The second is the methanol fraction with IC₅₀ value of 94.053 µg/mL, and the third is the *n*-hexane fraction with IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis that the ethyl acetate fraction contained glucoside, homoplantagin, and verolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili

seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidant activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo

State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for

extraction and fractionation. The 2,2- diphenyl-1-picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (n= 3). The inhibition percentage (%) was computed

as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influ-

enced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.


Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LC/MS profiles of (1) Blank, (2) Ethyl acetate fraction.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088	100.798 99.505 102.495	100.933±1.4995
	25	9.227	10.041	9.803		
	30	12.409	12.008	11.764		
	35	14.994	15.010	14.860		
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832	88.499 81.286 88.675	86.153±4.2161
	25	8.299	10.972	9.420		
	30	12.348	13.474	14.078		
	35	14.372	16.554	17.184		
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617	93.551 95.890 92.718	94.053±1.6445
	25	12.938	9.227	10.084		
	30	16.248	12.409	11.974		
	35	18.115	14.994	15.336		
	40	22.066	17.786	18.907		



**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpin-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the invitro DPPH radical scavenging ability of tombili seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the

previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice.

**Figure 4.** Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one)

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times ($n = 3$) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen

bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3.

Table 3. The profiles of the identified five major compounds.

Peak	tr (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	281.1275	280.12118	9.89
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	515704		+H
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C ₁₇ H ₁₆ N ₂ O ₂	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C ₂₆ H ₃₈ O ₇	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3β- glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min. Based on the LC/MS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was

shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- Badami, S., Moorkoth, S., Rai, S. R., Kannan, E., & Bhojraj, S. (2003). Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin*, 26(11), 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety*. 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/artic le/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbs.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties*. 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia*. 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitay N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Antioxidant, Cholesterol Lowering Activity, and Analysis of *Caesalpinia bonducella* Seeds Extract

Weny J.A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang², Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with an IC₅₀ value of 86.153 µg/mL. The second was the methanol fraction with an IC₅₀ value of 94.053 µg/mL, and the third was the *n*-hexane fraction with an IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5 % of the cholesterol activity. The LC-MS/MS analysis showed that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

Antioxidant, Cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS Analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili

seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo

State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for

extraction and fractionation. 2,2-diphenyl-1-picrylhydrazyl (DPPH) was purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H₂SO₄), and acetic anhydride were purchased from Merck (Darmstadt, Germany).




Figure 1. Tombili plant (1), Tombili fruit (2), Tombili powder (3), Extraction of tombili (4), Extract of tombili seed (5), Tombili extract evaporation, (6) Tombili thick extract (7).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis. The external appearance and necessary procedure for the extraction are shown on Fig 1.

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al., 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in tripli-

cate (*n* = 3). The inhibition percentage (%) was computed as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H₂SO₄ (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H₂SO₄ (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion


Extraction and partition

The extraction efficiency from plant materials is influenced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35 %), *n*-hexane (14.7 %) while ethyl acetate (31.7 %). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.

Antioxidant activity of tombili seed extracts

Item name: Blanko

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)



Item name: 19074-1-ToE

Channel name: 1: TOF MS^E (50-1200) 6eV ESI+ - Low CE (BPI)




Figure 2. LC/MS profiles of (1) Blank, (2) Ethyl acetate fraction.

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration. This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088	100.798 99.505 102.495	100,933±1.4995
	25	9.227	10.041	9.803		
	30	12.409	12.008	11.764		
	35	14.994	15.010	14.860		
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832	88.499 81.286 88.675	86.153±4.2161
	25	8.299	10.972	9.420		
	30	12.348	13.474	14.078		
	35	14.372	16.554	17.184		
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617	93.551 95.890 92.718	94.053±1.6445
	25	12.938	9.227	10.084		
	30	16.248	12.409	11.974		
	35	18.115	14.994	15.336		
	40	22.066	17.786	18.907		



**Figure 3.** Chemical structures of compounds: (1) 9,10-Dimethoxy-pterocarpin-3β-glucoside, (2) homoplantagin, (3) vernolic acid.

Table 2 shows the in vitro DPPH radical scavenging ability of tomboli seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tomboli seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the

previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice as illustrated in figure 4 which illustrate the reaction of the molecule.

**Figure 4.** Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one)

Cholesterol lowering activity of tomboli seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (n = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Figure 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in figure 3. The profiles for the compounds are shown on figure 5.

Table 3. The profiles of the identified five major compounds.


Peak	t _R (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3β-glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C17H16N2O2	Identified	281.1275	280.12118	9.89
5	Candidate Mass C26H38O7	Identified	463.2671	462.26175	9.35

	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C17H16N2O2	Identified	515704		+H
5	Candidate Mass C26H38O7	Identified	69954		+H

	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	C ₂₃ H ₂₆ O ₁₀	-1.4
2	Homoplantagin	Identified	C ₂₂ H ₂₂ O ₁₁	-1.2
3	Vernolic acid	Identified	C ₁₈ H ₃₂ O ₃	0.8
4	Candidate Mass C17H16N2O2	Identified	C ₁₇ H ₁₆ N ₂ O ₂	-0.9
5	Candidate Mass C26H38O7	Identified	C ₂₆ H ₃₈ O ₇	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.**Figure 6.** The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3β- glucoside.

role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min as shown on figure 6. Based on the LC/MS/MS spectra database finding, the compounds 9,10-dimethoxy-pterocarpan-3 β glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed

was first reported. The highest antioxidant activity was shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian J. Plant Res.* 1: 91-102.
- Badami, S., Moorkoth, S., Rai, S. R., Kannan, E., & Bhojraj, S. (2003). Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin*, 26(11), 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T. (2018). Use of Natural Antioxidant in the Inhibition of Oxidation: Review. *Comprehensive Reviews in Food Science and Food Safety*. 17: 1465-1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover), *Biomed Res. Int.* <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*, *Int. Res. J. Pharm.* 7: 12-17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, JK WP (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *Ejournal.Unsri.Ac.Id.* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/artic le/view/5159>.
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *J. Clin. Biochem. Nutr.* 40: 163-173. <https://doi.org/10.3164/jcbrn.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties*. 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia*. 12(3): 156-163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of *Caesalpinia bonduc*(L.) Roxb. *Pharmacogn J.* 10(1): 123-127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitany N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indones. J. Pharm.* 31: 171-180. <https://doi.org/10.22146/ijp.598>

Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extract

Weny J. A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang²,
Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Received 28 October 2022 ♦ Accepted 1 December 2022 ♦ Published @@ ##### 2023

Citation: Musa WJA, Bialangi N, Kilo AK, Situmeang B, Susparini NT, Rusydi ID (2023) Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extract. Pharmacia @@(@): @-@: <https://doi.org/10.3897/pharmacia.@@.e96817>

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with an IC₅₀ value of 86.153 µg/mL. The second was the methanol fraction with an IC₅₀ value of 94.053 µg/mL, and the third was the *n*-hexane fraction with an IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5% of the cholesterol activity. The LC-MS/MS analysis showed that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

antioxidant, cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb

et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for extraction and fractionation. 2,2- diphenyl-1-picrylhydrazyl (DPPH) was purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H_2SO_4), and acetic anhydride were purchased from Merck (Darmstadt, Germany).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis. The external appearance and necessary procedure for the extraction are shown on Fig. 1.




Figure 1. Tombili plant (a), Tombili fruit (b), Tombili powder (c), Extraction of tombili (d), Extract of tombili seed (e), Tombili extract evaporation, (f) Tombili thick extract (g).

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al. 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate (n=3). The inhibition percentage (%) was computed as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H_2SO_4 (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cho-

lesterol solution in a cuvette (BRAN UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H_2SO_4 (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influenced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35%), *n*-hexane (14.7%) while ethyl acetate (31.7%). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.




Figure 2. LC/MS profiles of (a) Blank, (b) Ethyl acetate fraction.

Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration. This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Table 2 shows the in vitro DPPH radical scavenging ability of tombili seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH

radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice as illustrated in Fig. 4 which illustrate the reaction of the molecule.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088		
	25	9.227	10.041	9.803	100.798	
	30	12.409	12.008	11.764	99.505	100,933±1.4995
	35	14.994	15.010	14.860	102.495	
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832		
	25	8.299	10.972	9.420	88.499	
	30	12.348	13.474	14.078	81.286	86.153±4.2161
	35	14.372	16.554	17.184	88.675	
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617		
	25	12.938	9.227	10.084	93.551	
	30	16.248	12.409	11.974	95.890	94.053±1.6445
	35	18.115	14.994	15.336	92.718	
	40	22.066	17.786	18.907		

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6




Figure 3. Chemical structures of compounds: (a) 9,10-Dimethoxy-pterocarpan-3β-glucoside, (b) homoplantagin, (c) vernolic acid.

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (*n* = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.




Figure 4. Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one).

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

.	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C17H16N2O2	Identified	281.1275	280.12118	9.89
5	Candidate Mass C26H38O7	Identified	463.2671	462.26175	9.35

.	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C17H16N2O2	Identified	515704		+H
5	Candidate Mass C26H38O7	Identified	69954		+H

.	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	C23H26O10	-1.4
2	Homoplantagin	Identified	C22H22O11	-1.2
3	Vernolic acid	Identified	C18H32O3	0.8
4	Candidate Mass C17H16N2O2	Identified	C17H16N2O2	-0.9
5	Candidate Mass C26H38O7	Identified	C26H38O7	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for n-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the

highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than n-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen bonding.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3β- glucoside.

Table 3. The profiles of the identified five major compounds.

Peak	tR (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3β-glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	–
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	–

Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Fig. 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside, homoplantagin, and vernolic acid are shown in Fig. 3. The profiles for the compounds are shown on Fig. 5.

Role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min as shown on Fig. 6. Based on the LC/MS/MS spectra database finding; the compounds 9,10-dimethoxy-pterocarpan-3β glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (m/z) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3β-glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.


References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian Journal of Plant Science and Research* 1: 91–102.
- Badami S, Moorkoth S, Rai SR, Kannan E, Bhojraj S (2003) Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin* 26(11): 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T (2018) Use of natural antioxidant in the inhibition of oxidation: review. *Comprehensive Reviews in Food Science and Food Safety* 17: 1465–1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover). *BioMed Research International* 2015: 643285. <https://doi.org/10.1155/2015/643285>

- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. Indonesian Journal of Pharmacy 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*. International Research Journal of Pharmacy 7: 12–17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, Pangkahila W, Budhiarta AAG (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. eJournal UNSR 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/article/view/5159>
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. Journal of Clinical Biochemistry and Nutrition 40: 163–173. <https://doi.org/10.3164/jcbl.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). International Journal of Food Properties 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. Jurnal Pendidikan Kimia 12(3): 156–163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical screening, total flavonoid and total phenolic content and antioxidant activity of different parts of *Caesalpinia bonduc*(L.) Roxb. Pharmacognosy Journal 10(1): 123–127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitary N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. Indonesian Journal of Pharmacy 31: 171–180. <https://doi.org/10.22146/ijp.598>

Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extract


Weny J. A. Musa ✉, Nurhayati Bialangi ✉, Ahmad Kadir Kilo ✉, **Boima Situmeang** ✉, Ninik Triayu Susparini ✉, Ilham Dwi Rusydi ✉

Article type: *Research Article* Type of review:  Conventional peer review

Subject editor: Rumiana Simeonova ✉

Date submitted: 28 October 2022

Authors

Name	Affiliation	City	Country	Corresponding author
Weny J. A. Musa ✉ ✎	Universitas Negeri Gorontalo	Gorontalo	Indonesia	✓
Nurhayati Bialangi ✉ ✎	Universitas Negeri Gorontalo	Gorontalo	Indonesia	
Ahmad Kadir Kilo ✉ ✎	Universitas Negeri Gorontalo	Gorontalo	Indonesia	
Boima Situmeang *  ✉ ✎	Department of Chemistry, Sekolah Tinggi Analisis Kimia Cilegon	Banten	Indonesia	
Ninik Triayu Susparini ✉ ✎	Department of Chemistry, Sekolah Tinggi Analisis Kimia Cilegon	Banten	Indonesia	
Ilham Dwi Rusydi ✉ ✎	Universitas Gadjah Mada	Sleman	Indonesia	

* Only the submitting author can handle this manuscript in the journal system.

Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extract

Weny J. A. Musa¹, Nurhayati Bialangi¹, Ahmad Kadir Kilo¹, Boima Situmeang²,
 Ninik Triayu Susparini², Ilham Dwi Rusydi³

¹ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Negeri Gorontalo, Gorontalo 96128, Indonesia

² Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten 043259, Indonesia

³ Department of Chemistry, Faculty of Mathematic and Natural Science, Universitas Gadjah Mada, Sleman 55281, Indonesia

Corresponding author: Weny J.A. Musa (wenymusa977@gmail.com)

Received 28 October 2022 ♦ Accepted 1 December 2022 ♦ Published 27 December 2023

Citation: Musa WJA, Bialangi N, Kilo AK, Situmeang B, Susparini NT, Rusydi ID (2023) Antioxidant, cholesterol lowering activity, and analysis of *Caesalpinia bonducella* seeds extract. Pharmacia 70(1): 97–103. <https://doi.org/10.3897/pharmacia.70.e96817>

Abstract

Tombili plant (*Caesalpinia bonducella*) belongs to the family of *Fabaceae*. The seed extract of tombili has been empirically used as a traditional medicine. The purpose of this research was to fractionate tombili seed extract and test their antioxidant and cholesterol lowering activities. Extraction was made into fractions using *n*-hexane, ethyl acetate, and methanol as a solvent. The chemical compound of the ethyl acetate fraction was analyzed using liquid chromatography-mass spectrometry (LC-MS/MS). Antioxidant activity was tested using the DPPH method. The highest antioxidant activity was shown in ethyl acetate fraction with an IC₅₀ value of 86.153 µg/mL. The second was the methanol fraction with an IC₅₀ value of 94.053 µg/mL, and the third was the *n*-hexane fraction with an IC₅₀ value of 100.933 µg/mL. The cholesterol lowering activity analysis showed that all fractions could inhibit cholesterol. The highest anti-cholesterol activity shown in ethyl acetate fraction with the concentration of 600 µg/mL can inhibit 81.5% of the cholesterol activity. The LC-MS/MS analysis showed that the ethyl acetate fraction contained glucoside, homoplantagin, and vernolic acid compounds.

Keywords

antioxidant, cholesterol lowering activity, *Caesalpinia bonducella*, LC-MS/MS analysis

Introduction

Cardiovascular diseases (CVDs) are the diseases of heart and blood vessels which considered one of the leading causes of death worldwide. Hypercholesterolemia is a key risk factor for cardiovascular disease and monitors the primary mortality of developing heart disease. The release of toxic free radicals by endothelial cells and vascular smooth muscle cells is the primary pathogenic factor for CVDs (Taleb

et al. 2018). The cholesterol present in these matrices also can oxidize in favorable conditions such as the presence of water, temperature, pH, and the form of substrate (de Oliveira et al. 2018). Cholesterol oxidation susceptible to attack by reactive oxygen species (ROS) which are formed of molecules hydroxy (OH), peroxy (OOH) groups, ¹O₂, ³O₂ and hydrogen peroxide (H₂O₂). The use of natural antioxidants from herbs represents an efficient alternative to control the formation of cholesterol oxidation product.

One common plant that has been used in traditional medicine is *Caesalpinia bonducella*, belongs to the family of *Fabaceae* (Aswar and Bhanudas 2011). This plant originated in Gorontalo, Indonesia (Sembiring et al. 2018). In Indonesia, *C. bonducella* is known as Tombili, Bagore, Kalici, Tinglur, and Areuy (Kakade et al. 2017). Tombili plants are also distributed in other tropical and subtropical parts of Asia, such as India, Sri Lanka, Vietnam, China, Myanmar, and Bangladesh (Musa et al. 2020).

In an earlier study, the phytochemical screening of Tombili seed showed that tombili seed contains saponins, phytosterols, flavonoids, and phenolics group compounds (Singh and Raghav 2012). This plant has proven pharmacological activity including the reduction of glucose levels (Esmaeili et al. 2015), reduce postprandial blood glucose levels and prevent the reduction of degraded plasma insulin levels in diabetic male albino rats (Larasati et al. 2016). Tombili seed methanol extract was also reported to have antihyperglycemic and hypolipidemic activity in induced diabetic rats (Modak et al. 2007), but limited report that studied its advantages to lowering cholesterol activity. Since the process of cholesterol oxidation is similar to the process of other unsaturated lipids, we try to prove the effectiveness antioxidants activity of tombili to disrupt the radical chain reactions by donating hydrogen atoms to these molecules. The active hydrogen atom of the antioxidant abstracted by the reactive radicals; thus, inactive species are formed. In this research, antioxidant, cholesterol lowering activity, and chemical content analyses from fractions of tombili were evaluated. Antioxidant activity *in vitro* method was done by DPPH scavenging (Rafi et al. 2020).

Materials and methods

Materials

The tombili seeds collected from Bobohu villages, Gorontalo province, Indonesia. Tombili was identified at the Laboratory of Biology Department (Plant Taxonomy), Faculty of Mathematics and Natural Science, Gorontalo State University, Indonesia with the specimen being identified by Weny JA Musa and verified by the relevant department under specimen number of 021/UN47.B4./LL/2019 as *Caesalpinia Bonducella* (L.) Fleming. The chemicals used were methanol, ethyl acetate, and *n*-hexane solvent (Pro Analysis Grade) used for extraction and fractionation. 2,2- diphenyl-1-picrylhydrazyl (DPPH) was purchased from Sigma-Aldrich (St. Louis, MO, USA). The cholesterol solution Sigma-Aldrich (St. Louis, MO, USA), sulfate acid (H_2SO_4), and acetic anhydride were purchased from Merck (Darmstadt, Germany).

Extraction and partition

The extract was made with 1.2 kg of Tombili seed powder and macerated for three days. To obtain *n*-hexane and ethyl acetate fractions, 200 of the extract was partitioned by liquid-liquid extraction with *n*-hexane and ethyl acetate at a 1:1 ratio. The rotary evaporator was used to concentrate all fractions. The yield of each solvent was then determined. The fractions were created by combining *n*-hexane (29.5 g) and ethyl acetate (63.4 g) Antioxidant and cholesterol-lowering activity was assessed in all fractions. The fraction with the highest antioxidant activity and the lowest cholesterol activity was injected into an LC-MS/MS for chemical compound analysis. The external appearance and necessary procedure for the extraction are shown on Fig. 1.




Figure 1. Tombili plant (a), Tombili fruit (b), Tombili powder (c), Extraction of tombili (d), Extract of tombili seed (e), Tombili extract evaporation, (f) Tombili thick extract (g).

Antioxidant activity test procedure

The antioxidant activity of each fraction was tested using the DPPH technique. According to Fareza et al. 2021, the test process was followed. A total of 2400 L stock solution of fractions was made in series concentration (200, 250, 300, 350, and 400 ppm). 600 L of DPPH solution with a concentration of 160 g/mL was added to each tube. The tube was incubated at room temperature for 30 minutes. The absorbance was measured at 517 nm using UV-Vis spectrophotometry (Rafi et al. 2020). As a control, methanol and DPPH were employed. The experiments were carried out in triplicate ($n=3$). The inhibition percentage (%) was computed as follows:

$$\% \text{inhibition} = \left[\frac{(\text{absorbance control} - \text{absorbance sample})}{\text{absorbance control}} \right] \times 100\%$$

Determination of cholesterol-lowering potential

The fractions' reduced cholesterol levels were determined by comparing them to the blank absorbance (cholesterol solution). The cholesterol solution stock was created by combining 100 mg of cholesterol in 100 mL of ethanol (1000 ppm). H_2SO_4 (0.1 mL) and acetic anhydride (2 mL) were added to 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cho-

lesterol solution in a cuvette (BRAN UV cuvette) and diluted with ethanol solvent until 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions, 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution (1000 ppm) to concentrations of 100, 200, 400, and 600 ppm sample, H_2SO_4 (0.1 mL), acetic anhydride (2 mL), and ethanol until 5 mL were added, respectively. After being incubated for 30 minutes at room temperature, the absorbance of the solutions was measured at λ 423 nm (Musa et al. 2019).

Results and discussion

Extraction and partition

The extraction efficiency from plant materials is influenced by several factors, including the chemical screening of phytochemicals, the partition method used, the size of sample particles, and the presence of interfering substances (Stalikas 2007). The extractions yield depends on the temperature, extraction time, solvent polarity, and sample content. In this work, the extracts of tombili seed were partitioned by using various range polarity of solvent from nonpolar (*n*-hexane) to polar (methanol). The final methanol fraction obtained 107.1 g (55.35%), *n*-hexane (14.7%) while ethyl acetate (31.7%). It can also be seen that the fraction yield of methanol is higher than ethyl acetate. These results show that the extraction yield.




Figure 2. LC/MS profiles of (a) Blank, (b) Ethyl acetate fraction.

Antioxidant activity of tombili seed extracts

DPPH was used to determine the antioxidant activity of tombili seed extracts. With the highest absorption band at 517 nm, DPPH was used as a stable organic free radical reagent. The absorption of this radical species was lost when the polarity of the solvent used in the extraction process of tombili seed was increased. Primary metabolites (proteins and carbohydrates) have been extracted in water and ethanol, resulting in a higher yield than ethyl acetate and *n*-hexane.

Accepted, resulting in a purple-to-yellow visual discoloration. This radical reacts quickly with a wide range of samples and is sensitive enough to distinguish active ingredients at low concentrations (Esmaeili et al. 2015). The results of the antioxidant activity test of fractions shown in Table 1.

Table 2 shows the in vitro DPPH radical scavenging ability of tombili seed extract in different solvents. The *n*-hexane extract had the lowest DPPH radical inhibition, with an IC₅₀ value of 100.933, while the ethyl acetate extract had the highest activity (86.153 g/mL). Although the DPPH

radical scavenging activities of different solvent extracts were lower than those of ascorbic acid as a standard reference compound, this study shows that tombili seed extract has free radical scavenging properties as a primary antioxidant. Some free radical reactions are inhibited by phenolic and flavonoid compounds. Some phenolic compounds in ethyl acetate extracts, such as flavonoid and phenolic acids, demonstrated strong antioxidant activity via their reductive capacity (Rafi et al. 2020). The hydroxyl group of homoplantagin and vernolic acid compounds was thought to be involved in free radical reduction. Considering the abundance of hydroxyl group of homoplantagin it can be considered the biologically active from the fraction of ethyl acetate, thus allowing the antioxidant activity of the fraction toward the DPPH. Homoplantagin contains several aromatic rings and hydroxyl group to implement the antioxidant mechanism. In comparison with the previous research done by Badami et al. (2003) using *Caesalpinia Sappan* yield a very strong antioxidant activity from the ethyl acetate extract while no further metabolites were explained from the literature. The secondary metabolites consisted of glucoside, homoplantagin, and vernolic acid contributed to the fraction ability as antioxidant for the DPPH method analysis due to the hydroxyl group from the homoplantagin to prevent free radicals from DPPH to oxidize the supposed protected molecule in practice as illustrated in Fig. 4 which illustrate the reaction of the molecule.

Table 1. Antioxidant activity test of fractions of *Caesalpinia bonducella* seed.

Fractions	Concentration (ppm)	% inhibition ¹	% inhibition ²	% inhibition ³	IC ₅₀ µg/mL	Average (µg/mL)
<i>n</i> -Hexane	20	6.308	5.900	6.088		
	25	9.227	10.041	9.803	100.798	
	30	12.409	12.008	11.764	99.505	100,933±1.4995
	35	14.994	15.010	14.860	102.495	
	40	16.856	17.080	16.718		
Ethyl acetate	20	4.251	5.678	6.832		
	25	8.299	10.972	9.420	88.499	
	30	12.348	13.474	14.078	81.286	86.153±4.2161
	35	14.372	16.554	17.184	88.675	
	40	17.712	20.692	18.840		
Methanol	20	11.334	6.308	6.617		
	25	12.938	9.227	10.084	93.551	
	30	16.248	12.409	11.974	95.890	94.053±1.6445
	35	18.115	14.994	15.336	92.718	
	40	22.066	17.786	18.907		

Table 2. Cholesterol lowering activity test of fractions *Caesalpinia bonducella* seed.

Sample	Concentration (ppm)	% exhibited ¹	% exhibited ²	% exhibited ³	Average (%)
<i>n</i> -hexane	100	7.9	6.1	7	7
	200	2	6.1	4.05	4.05
	400	22.8	21.2	22	22
	600	25.7	31.3	28.5	28.5
Positive Control	100	57.4	49.5	53.5	53.5
Ethyl acetate	100	15.9	13.8	14.9	14.9
	200	43	40.6	41.8	41.8
	400	67.9	67.5	67.7	67.7
	600	80.9	82	81.5	81.5
Positive Control	100	90.9	91.6	91.4	91.3
Methanol	100	15.5	7.7	11.6	11.6
	200	47.1	39.5	43.3	43.3
	400	68.2	65.5	66.9	66.9
	600	76.2	77.3	76.8	76.8
Positive Control	100	83.7	83.4	83.6	83.6




Figure 3. Chemical structures of compounds: (a) 9,10-Dimethoxy-pterocarpan-3β-glucoside, (b) homoplantagin, (c) vernolic acid.

Cholesterol lowering activity of tombili seed extracts

The amount of cholesterol-lowering salt was reduced dose-dependently. The cholesterol level in the untreated fractions, also known as the negative control, was calculated to be 100%. All experiments were repeated three times (*n* = 3) and compared to negative and positive controls. Table 2 shows the outcome of the cholesterol-lowering activity.




Figure 4. Antioxidant chemical reaction of 2,2-diphenyl-1-picrylhydrazyl and homoplantagin (5-hydroxy-2-(4-hydroxyphenyl)-6-methoxy-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one).

Item name: 19074-1-ToE, Sample position: 2:A,5, Replicate number: 1

.	Component name	Identification status	Observed m/z	Neutral mass (Da)	Observed RT (min)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	485.1405	462.15260	8.63
2	Homoplantagin	Identified	463.1223	462.11621	7.16
3	Vernolic acid	Identified	319.2252	296.23514	9.35
4	Candidate Mass C17H16N2O2	Identified	281.1275	280.12118	9.89
5	Candidate Mass C26H38O7	Identified	463.2671	462.26175	9.35

.	Component name	Identification status	Detector counts	Response	Adducts
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	34844	24981	+Na
2	Homoplantagin	Identified	72640	49657	+H
3	Vernolic acid	Identified	3466951	2607711	+Na
4	Candidate Mass C17H16N2O2	Identified	515704		+H
5	Candidate Mass C26H38O7	Identified	69954		+H

.	Component name	Identification status	Formula	Mass error (mDa)
1	9,10-Dimethoxy-pterocarpan-3-O-β-D-glucoside	Identified	C23H26O10	-1.4
2	Homoplantagin	Identified	C22H22O11	-1.2
3	Vernolic acid	Identified	C18H32O3	0.8
4	Candidate Mass C17H16N2O2	Identified	C17H16N2O2	-0.9
5	Candidate Mass C26H38O7	Identified	C26H38O7	-2.0

Figure 5. The profiles of the identified five major compounds measured with LC/MS apparatus.

It was first reported that tombili fractions have cholesterol-lowering properties. All fractions were carried out at concentrations of 100, 200, 400, and 600 ppm, respectively. The colorimetric experiment revealed an anti-cholesterol effect for *n*-hexane, ethyl acetate, and methanol fractions. The ethyl acetate component has the highest percentage. More than 50% of cholesterol can be detected in a 600ppm ethyl acetate fraction. Ethyl acetate has a greater effect than *n*-hexane and methanol.

Caused by the concentration of the chemical compound in fraction with the hydroxyl group that interacted with another functional group through hydrogen bonding. Unlike ethyl acetate, the intramolecular reaction of *n*-hexane and methanol fractions does not occur.

Identification of chemical content

The ethyl acetate fraction TEA3 was analyzed for its chemical composition using LC-MS/MS (Fig. 2). Further MS analysis of fraction was conducted on three major compounds 9,10-Dimethoxy-pterocarpan-3-O- β -D-glucoside, homoplantagin, and vernolic acid are shown in Fig. 3. The profiles for the compounds are shown on Fig. 5.

Role in its higher activity as an anticancer therapy. These outcomes Liquid chromatography-mass spectrophotometry (LC/MS) analysis of ethyl acetate fraction showed five obvious peaks at retention times of 7.34, 8.63, 9.31, 9.38, and 9.89 min as shown on Fig. 6. Based on the LC/MS/MS spectra database finding; the compounds 9,10-dimethoxy-

pterocarpan-3 β glucoside, homoplantagin, and vernolic acid were identified at retention times 7.34, 8.63, and 9.31 min, respectively. The fragmentation resulted in five candidate masses with spectra (*m/z*) of 462.11621, 485.1406, 39.2252, 463.2671, and 281.127 are shown in Table 3.

Conclusion

The tested cholesterol lowering property of *n*-hexane, ethyl acetate, and methanol fraction from tombili seed was first reported. The highest antioxidant activity was shown in the ethyl acetate fraction. All fractions showed cholesterol lowering activity. Most importantly, the raised concentration of fractions exhibited a dose-dependent manner. The analysis of chemical content showed that the ethyl acetate fraction contains 9,10-Dimethoxy-pterocarpan-3 β -glucoside, homoplantagin, and vernolic acid compounds as lowering cholesterol agent.

Acknowledgements

The authors would like to thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which has provided research funding through PTUPT schemes in 2021. Thank you, Mrs. Sofa Fajriah, in the Indonesian Institute of Sciences for LC/MS measurements.




Figure 6. The mass spectrometer analysis on 9,10-dimethoxy- pterocarpan-3 β - glucoside.

Table 3. The profiles of the identified five major compounds.

Peak	tR (min)	Formula	Observed actual mass	Neutral mass (Da)	Adduct	Mass error (mDa)	Identification
1	8.63	C ₂₃ H ₂₆ O ₁₀	485.1405	462.15260	+Na	-1.4	9,10-dimethoxy-pterocarpan-3 β -glucoside
2	7.34	C ₂₂ H ₂₂ O ₁₁	463.1223	462.11621	+H	-1.2	Homoplantagin
3	9.31	C ₁₈ H ₃₂ O ₃	319.2252	296.23514	+Na	0.8	Vernolic acid
4	9.89	C ₁₇ H ₁₆ N ₂ O ₂	281.127	280.12118	+H	-0.9	-
5	9.38	C ₂₆ H ₃₈ O ₇	463.2671	462.26175	+H	-2.0	-

References

- Aswar BR, Bhanudas SK (2011) Assessment of hypoglycemic and antidiabetic effects of *Caesalpinia bonduc* (L.) Roxb. seeds in alloxan induced diabetic rat and its phytochemical, microscopic, biochemical, and histopathological evaluation. *Asian Journal of Plant Science and Research* 1: 91–102.
- Badami S, Moorkoth S, Rai SR, Kannan E, Bhojraj S (2003) Antioxidant activity of *Caesalpinia sappan* heartwood. *Biological and Pharmaceutical Bulletin* 26(11): 1534–1537. <https://doi.org/10.1248/bpb.26.1534>
- de Oliveira VS, Ferreira FS, Cople MCR, Labre TS, Augusta IM, Gamallo OD, Saldanha T (2018) Use of natural antioxidant in the inhibition of oxidation: review. *Comprehensive Reviews in Food Science and Food Safety* 17: 1465–1483. <https://doi.org/10.1111/1541-4337.12386>
- Esmaili AK, Taha AK, Mohajer S, Banisalam B (2015) Antioxidant activity and total phenolic and flavonoid content of various solvent extracts from *in-vivo* and *in-vitro* grown *Trifolium pratense* L. (Red clover). *BioMed Research International* 2015: 643285. <https://doi.org/10.1155/2015/643285>
- Fareza MS, Choironi NA, Susilowati SS, Rini MP (2021) LC-MS / MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* Stembark. *Indonesian Journal of Pharmacy* 32: 356–364.
- Kakade NR, Pingale SS, Chaskar MG (2017) Phytochemical and pharmacological review of *Caesalpinia bonducella*. *International Research Journal of Pharmacy* 7: 12–17. <https://doi.org/10.7897/2230-8407.0712139>
- Larasati V, Pangkahila W, Budhiarta AAG (2016) Oral administration of *Caesalpinia bonducella* seeds ethanolic extract decreased post prandial blood glucose level and prevented the reduction of fasting insulin. *eJournal UNSR* 3: 87–95. <https://ejournal.unsri.ac.id/index.php/jkk/article/view/5159>
- Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA (2007) Indian herbs and herbal drugs used for the treatment of diabetes. *Journal of Clinical Biochemistry and Nutrition* 40: 163–173. <https://doi.org/10.3164/jcbn.40.163>
- Musa WJA, Situmeang B, Sianturi J (2019) Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae). *International Journal of Food Properties* 22(1): 1439–1444. <https://doi.org/10.1080/10942912.2019.1650762>
- Musa WJA, Duengo S, Kilo AK, Situmeang B (2020) Alkaloid compound from Tombili (*Caesalpinia bonduc*) as biopesticide agent on rice plants. *Jurnal Pendidikan Kimia* 12(3): 156–163. <https://doi.org/10.24114/jpkim.v12i3.21165>
- Sembiring EN, Elya B, Sauriasari R (2017) Phytochemical screening, total flavonoid and total phenolic content and antioxidant activity of different parts of *Caesalpinia bonduc* (L.) Roxb. *Pharmacognosy Journal* 10(1): 123–127. <https://doi.org/10.5530/pj.2018.1.22>
- Rafi M, Meitary N, Septaningsih DA, Bintang M (2020) Phytochemical profile and antioxidant activity of *Guazuma ulmifolia* leaves extracts using different solvent extraction. *Indonesian Journal of Pharmacy* 31: 171–180. <https://doi.org/10.22146/ijp.598>