

BUKTI KORESPONDENSI ARTIKEL JURNAL INTERNASIONAL BEREPUTASI

Judul artikel : Nanomineral Fortification on Traditional Saddah Gohoge Grits
Nama Jurnal : Food Science and Technology (Brazil), Volume 43, 27 April 2023
Penulis : Rahim Husain, Fernandy Djailani, Rita Marsuci Harmain

No	PERIHAL	TANGGAL
1	Account Created	23 September 2022
2	Manuscript Submission	23 September 2022
3	Manuscript Decision (Tahap 1)	8 Oktober 2022
4	Pengiriman Revisi	5 November 2022
5	Manuscript Decision (Tahap 2)	7 November 2022
6	Letter of Acceptance	3 Maret 2023
7	Artikel Proofreading (Tahap 1)	9 Maret 2023
8	Pengiriman Revisi Proofreading (Tahap 1)	16 Maret 2023
9	Artikel Proofreading (Tahap 2)	16 Maret 2023
10	Pengiriman Revisi Proofreading (Tahap 2)	27 Maret 2023
11	Final proofreading completed	27 Maret 2023
12	Layout Proof	30 Maret 2023
13	Approval Layout Proof oleh penulis	1 Maret 2023
14	Artikel terbit April 2023, Volume 43	

Account Created
(23 September 2022)



Rahim Husain <rahim@ung.ac.id>

Food Science and Technology - Account Created in ScholarOne Manuscripts

Marcos Revista <onbehalf@manuscriptcentral.com>

23 September 2022 pukul 15.07

Balas Ke: revista@sbcta.org.br

Kepada: rahim@ung.ac.id

23-Sep-2022

Dear Dr. Husain:

Welcome to the Food Science and Technology - ScholarOne Manuscripts site for online submission and review.

Your USER ID for your account at <https://mc04.manuscriptcentral.com/cta-scielo> is as follows:USER ID: rahim@ung.ac.id

If you are unsure of the password that you set when you created your account you should click the link below which will take you directly to the option for setting a new password.

https://mc04.manuscriptcentral.com/cta-scielo?URL_MASK=1682377735fb43e4ac5730e6017ff7f2 Please note that the single use link will expire on 26-Sep-2022 8:07:27 AM GMT / 26-Sep-2022 5:07:27 AM BRT. If the single use link has expired, you can generate a single use password by entering your email address into the Password Help function on your site log in page: <https://mc04.manuscriptcentral.com/cta-scielo>

Thank you for your participation.

Sincerely,
Food Science and Technology Editorial Office

To create a new ORCID iD record or to link your user account to an existing ORCID iD, simply click this link:

https://mc04.manuscriptcentral.com/cta-scielo?URL_MASK=b9563ac5a3a94780b196b01c6ebc50f3[Log in to Remove This Account](#)

Manuscript Submission

(23 September 2022)



Rahim Husain <rahim@ung.ac.id>

Food Science and Technology - Manuscript ID CTA-2022-0991

Marcos Revista <onbehalf@manuscriptcentral.com>

23 September 2022 pukul 15.25

Balas Ke: revista@sbcta.org.br

Kepada: rahim@ung.ac.id

Cc: rahim@ung.ac.id, fernandydjailani@ung.ac.id, ritaharmain@ung.ac.id

23-Sep-2022

Dear Dr. Husain:

Your manuscript entitled "NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS" has been successfully submitted online and is presently being given full consideration for publication in the Food Science and Technology.

Your manuscript ID is CTA-2022-0991.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at <https://mc04.manuscriptcentral.com/cta-scielo> and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to <https://mc04.manuscriptcentral.com/cta-scielo>.

Thank you for submitting your manuscript to the Food Science and Technology.

Sincerely,
Food Science and Technology Editorial Office

NANOMINERAL FORTIFICATION OF FISH BONE FAMILY *Lutjanus* sp. ON TRADITIONAL SADDAH *GOHOGE* PORRIDGE

Rahim Husain¹, Fernandy Djailani², Rita Marsuci Harmain³

1,2,3). Lecturers of fishery processing technology, Faculty of fisheries and marine sciences, Universitas Negeri Gorontalo

Email¹ : rahim@ung.ac.id

ABSTRACT

Saddah gohoge porridge is a traditional food that is favored by the people of Gorontalo in Indonesia, especially the elderly and when children are growing. The aim of the study was to obtain the physical, chemical and organoleptic characteristics of the saddah *gohoge* porridge fortification of nanomineral fish bones from the family *Lutjanus* sp. at 3 treatment levels fortification concentration of 0%, 15% and 20%. The results of the physical analysis showed that the viscosity ranged from 8309.47 – 14069.73 cP and was significant, the coloring ranged from 47.1 – 51.33(L), the water content ranged from 80.33 – 86.82%, the ash content was 0.92 – 1.53%, the fat content was 0.47 – 1.22%, the protein content was 7.16 – 10.51%. , fiber content 1.77 – 3.8% and carbohydrate content by difference 92.48 – 95.43%. The results of fortification of bone waste from the family of *Lutjanus* sp. showed that the higher the fortification concentration, the higher the nanomineral content in the saddah gohoge slurry. The results of the hedonic quality organoleptic test based on the response of the panelists, namely the clean appearance is not dull, the texture is slightly smooth, slightly thick, the color is slightly yellowish, the aroma is slightly pungent, the aroma of corn, the taste is rather good, the taste of corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma and taste.

Keywords: saddah *gohoge* porridge, fortification, characteristics, *Lutjanus* sp. nanomineral

Introduction

Saddah *gohoge* porridge is a local name for food from the Gorontalo region in Indonesia which is generally popular, especially for the elderly and growing children because it has a soft texture and corn flavor. This porridge contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (RI No. 43 20Permenkes19 2019) .

Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63g/kg), phosphorus (2.38g/kg), and carbonates.

In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite $[Ca_{10}(OH)(PO_4)_6]$ in the form of crystals attached to fibrillar collagen.(Malde et al. 2010)(Agronomy Research 2003)(Litopenaeus et al. 2012). (Miller, Jarvis, and McBean 2001).

Fish bones with calcium mineral content are found in extracellular fluid and in cell tissue. Calcium is found in the parathyroid glands, extracellular calcium is in intestinal absorption, renal reabsorption and bone formation(Agronomy Research 2003).

Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients(Greiner 2009)(Lekahena et al. 2014).

This study carried out fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the saddah gohoge porridge.

Method

The material used is ground corn starch of the *Bisi 2* variety in the Gorontalo area. Fish bone waste of the *Lutjanus* sp. obtained from industrial waste processed fish fillet family *Lutjanus* sp. The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission *type*), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5.

Fishbone preparation method *by* (Harmain, R.M. 2018) *modification*

extraction of nanominerals precipitation method, immersion time 48 hours (Litopenaeus et al. 2012) .

Making corn grits using a hammer mill and a 60 mesh disc mill(Ahmad, Une, and Bait 2019)

Moisture content, ash content, fiber content (Al-mentafji 2005).

Carbohydrates by different crude protein content, fat content) (Al-mentafji 2005).

Calcium Atomic Absorption Spectrometry (AAS) analysis by wet ashing method at = 422.7 nm particle size containing nanominerals and morphology of fish bones using the Scanning Electron Microscope (SEM) Phenom ProX G5.

The nanocalcium particle size was analyzed using Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan and Kim, 2010),(Thunnus 2010).

Theory

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Xliii 2018).

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometre so, they can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size(Singh 2016).

Findings And Discussion

Physical Test Results

The results of the physical test of the thickness of the saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals, the value of the viscosity level increased (Table 1). This was because the volume of boiled water for saddah *gohoge* porridge was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if you want to achieve the same thickness as without nanomineral fortification.

Table 1. Viscosity levels of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp.

Concentration	Saddah <i>Gohoge</i> Viscosity (cP)			Average
	1	2	3	
0 %	8353.2	8268.4	8306.8	8309.47
15 %	10369.3	10127.9	10296.9	10264.7
20 %	13961.9	14167.8	14079.5	14069.73

The results of the staining test of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 45.5 – 50.8 (L) and the notations a and b depended on the fortification of nanominerals in the saddah *gohoge* porridge which affected the basic color of the saddah *gohoge* porridge, which was yellowish (Table 2).

Table 2. Color of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp.

Concentration	Saddah Porridge Color								
	1			2			3		
	L	a	b	L	a	b	L	a	b
0 %	50.1	0.7	16	45.5	3.9	13.5	45.7	6.7	18.4
15 %	53.6	-125	17.8	46.5	3.4	17.4	50	3.9	13.7
20 %	50.7	0.1	13.5	50.8	2.5	17.8	52.5	3.5	18.2

Chemical Test Results

Chemical test results (proximate) include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

Table 3. Proximate levels of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp.

Concentration	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Karbohydrates Content
0 %	86.82 ^b	0.92 ^b	7.16 ^b	0.47 ^b	1.77 ^b	95.37 ^b
15 %	85.03 ^a	1.13 ^a	8.53 ^b	0.74 ^b	3.3 ^a	95.43 ^b
20 %	80.33 ^a	1.31 ^a	9.63 ^b	1.21 ^a	3.60 ^a	92.48 ^b
25 %	79.70 ^b	1.53 ^a	10.51 ^b	1.22 ^a	3.8 ^a	92.96 ^b

Note: the same letter notation shows no significant difference at = 0.05.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. Without fortification or 0% as a fortification concentration increased,

the water content decreased with the use of the volume of water for cooking saddah *gohoge* porridge which was the same as 1000 mL for all concentrations. The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20%. This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration. The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly different but significantly different from those without nanomineral fortification.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different. The fat content showed that the fortified concentration of 15% was significantly different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different. For fiber content, the concentration of fortified 15%, 20% and 25% was not significantly different but significantly different from without fortification (0%). The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

Fish bone nanominerals family *Lutjanus* sp. To be fortified in saddah *gohoge* porridge, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah *gohoge* porridge made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe.

Organoleptic Test Results

The results of organoleptic tests include hedonic and hedonic quality tests based on appearance, texture, color, aroma and taste criteria.

Appearance

The appearance of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. get a score of 3.68, the net score is not dull. Hedonic score 3.96 likes rating. The saddah *gohoge*

porridge made from shelled corn is processed to be like corn flour and filtered so that it looks clean from the appearance. The storage period also affects the appearance of corn flour, the longer the storage time, the duller the appearance of the corn.

Texture

The hedonic quality score of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. is 3.48 rather thick fine rating. Hedonic scores obtained 3.64 ratings of likes. The texture of saddah *gohoge* porridge with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content.

Color

Color of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. resulting hedonic quality value is 3.08 slightly yellowish rating. Hedonic score obtained 4.04 with a rating of likes. Saddah *gohoge* porridge with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E.

Flavor

An important indicator of the acceptance of a food product is flavor. The flavor of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. obtained a hedonic quality value of 3.96, which is slightly pungent with the flavor of corn. The hedonic score obtained is 3.64 with the assessment criteria like.

Taste

The hedonic quality score of saddah *gohoge* porridge fortification of fish bone nanominerals *Lutjanus* sp. resulted in a score of 3.16 with an assessment of somewhat good taste of corn and a hedonic score of 3.52 with an assessment of somewhat good taste of corn. Fortification of fish bone nanominerals in one serving of binte biluhuta cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. BPOM (2011) states that a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto, Trilaksani, and Aulia Azzahra 2020).

Conclusion

Proximate levels, namely ash content, protein content, fat content and fiber content increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in saddah *gohoge* porridge, but the water content decreased due to the use of the same volume of water and the composition of saddah *gohoge*. The hedonic quality of saddah *gohoge* porridge fortification of nanomineral fish bones of the family *Lutjanus* sp. based on the response of the panelist, namely clean, slightly not dull, slightly smooth, slightly thick, slightly yellowish, slightly pungent smell of corn, slightly good taste of corn. The hedonic response of the panelists to the saddah *gohoge* porridge fortification of fish bone nanominerals from the family *Lutjanus* sp. is fond of all assessments of appearance, texture, color, aroma and taste.

Acknowledgments

thank all those who have assisted in this research and the reviewers of journals for publishing the results of this research.

References

“Agronomy Research.” 2003.

Ahmad, Lisna, Suryani Une, and Yoyanda Bait. 2019. “Karakteristik Komponen Gizi, Antioksidan, Dan Respon Organoleptik Bubur Jagung Tradisional Gorontalo Dengan Ekstrak Daun Kersen (*Muntingia Calabura* L).” *agriTECH* 38(4): 463.

Al-mentafji, Hayder N. 2005. “Of Ficial Methods of Analysis of AOAC INTERNATIONAL.” *Aoac* (February).

Greiner, Ralf. 2009. “Current and Projected Applications of Nanotechnology in the Food Sector Aplicações Atuais e Futuras Da Nanotecnologia No Setor Alimentício.” *Journal of Brazillian Society of Food and Nutrition* 34(1): 243–60.
http://sban.cloudpaine.com.br/files/revistas_publicacoes/228.pdf.

Harmain, R.M., F.A. Dali and R. Husain. 2018. “Nanocalcium Characterization of Cakalang Fish Bone Flour (*Katsuwonus Pelamis* L).” *International Journal of Innovative Science and Reasearch Technology* 3(10): 306–8.

Lekahena, Vanessa, Didah Nur Faridah, Rizal Syarief, and R. Peranginangin. 2014. “Karakterisasi Fisikokimia Nanokalsium Hasil Ekstraksi Tulang Ikan Nila Menggunakan Larutan Basa Dan Asam.” *Jurnal Teknologi dan Industri Pangan* 25(1): 57–64.

Litopenaeus, Vannamei et al. 2012. “Litopenaeus Vannamei).” III(1).

Malde, Marian K et al. 2010. “Calcium from Salmon and Cod Bone Is Well Absorbed in Young Healthy Men : A Double- Blinded Randomised Crossover Design.” : 1–9.

Miller, Gregory D, Judith K Jarvis, and Lois D McBean. 2001. “The Importance of Meeting

Calcium Needs with Foods.” *Journal of the American College of Nutrition* 20(2): 168S-185S. <https://doi.org/10.1080/07315724.2001.10719029>.

RI No. 43 20Permenkes19. 2019. “No Titleس.” *ペインクリニック学会治療指針 2* (2): 1–13.

Riyanto, Bambang, Wini Trilaksani, and Vegatarani Aulia Azzahra. 2020. “Desain Pangan Instan Gizi Khusus Lansia Berbasis Binte Biluhuta Diperkaya Nanomineral Tulang Ikan.” *Jurnal Fishtech* 9(2): 65–77.

Singh, Pankaj. 2016. “Use of Nano Feed Additives in Livestock Feeding.” *International Journal of Livestock Research* 6(1): 1.

Thunnus, Tuna. 2010. “Effect of Temperature on Isolation and Characterization of Hydroxyapatite from Tuna (*Thunnus Obesus*) Bone.” : 4761–72.

Xliii, Volume. 2018. “Mengenal Marga.” *XLIII*(2003): 29–39.

Manuscript Decision (Tahap 1)

(8 Oktober 2022)



Rahim Husain <rahim@ung.ac.id>

Food Science and Technology - Decision on Manuscript ID CTA-2022-0991

Adriano Cruz <onbehalf@manuscriptcentral.com>

8 Oktober 2022 pukul 07.37

Balas Ke: adriano.cruz@ifrj.edu.br

Kepada: rahim@ung.ac.id

07-Oct-2022

Dear Dr. Husain:

Manuscript ID CTA-2022-0991 entitled "NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS" which you submitted to the Food Science and Technology, has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

The reviewer(s) have recommended publication, but also suggest some minor revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript.

To revise your manuscript, log into <https://mc04.manuscriptcentral.com/cta-scielo> and enter your Author Center, where you will find your manuscript title listed under "Manuscripts with Decisions." Under "Actions," click on "Create a Revision." Your manuscript number has been appended to denote a revision.

You may also click the below link to start the revision process (or continue the process if you have already started your revision) for your manuscript. If you use the below link you will not be required to login to ScholarOne Manuscripts.

*** PLEASE NOTE: This is a two-step process. After clicking on the link, you will be directed to a webpage to confirm.

https://mc04.manuscriptcentral.com/cta-scielo?URL_MASK=3f443adb659d42b09957094ee17435ac

You will be unable to make your revisions on the originally submitted version of the manuscript. Instead, revise your manuscript using a word processing program and save it on your computer. Please also highlight the changes to your manuscript within the document by using the track changes mode in MS Word or by using bold or colored text.

Once the revised manuscript is prepared, you can upload it and submit it through your Author Center.

When submitting your revised manuscript, you will be able to respond to the comments made by the reviewer(s) in the space provided. You can use this space to document any changes you make to the original manuscript. In order to expedite the processing of the revised manuscript, please be as specific as possible in your response to the reviewer(s).

IMPORTANT: Your original files are available to you when you upload your revised manuscript. Please delete any redundant files before completing the submission.

Because we are trying to facilitate timely publication of manuscripts submitted to the Food Science and Technology, your revised manuscript should be submitted by 07-Nov-2022. If it is not possible for you to submit your revision by this date, we may have to consider your paper as a new submission.

Once again, thank you for submitting your manuscript to the Food Science and Technology and I look forward to receiving your revision.

Sincerely,
Dr. Adriano Cruz
Editor-in-Chief, Food Science and Technology
adriano.cruz@ifrj.edu.br

Associate Editor

Comments to the Author:

1. Provide more context and background information to help readers understand the significance and relevance of the research being presented.
2. Providing a clear and concise overview of the study design, including the research question or hypothesis being tested, the study population or sample, and any inclusion/exclusion criteria.
3. Describing the procedures used to collect and analyze data, including any instruments or tools used, data collection protocols, and statistical analysis methods.
4. Providing detailed information on any materials or equipment used in the study, including sources, specifications, and any modifications or adaptations made to the instruments or protocols.
5. Avoiding speculation or interpretation in the results section, and focusing instead on reporting the facts and observations.
6. Some recent FST papers (2021,2022,3-4 papers) must be added.
7. Remove old references

Entire Scoresheet:

Pengiriman Revisi

(5 November 2022)



Rahim Husain <rahim@ung.ac.id>

Food Science and Technology - Manuscript ID CTA-2022-0991.R1

Marcos Revista <onbehalf@manuscriptcentral.com>

5 November 2022 pukul 01.06

Balas Ke: revista@sbcta.org.br

Kepada: rahim@ung.ac.id

Cc: rahim@ung.ac.id, fernandydjailani@ung.ac.id, ritaharmain@ung.ac.id

06-Nov-2022

Dear Dr. Husain:

Your manuscript entitled "NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS" has been successfully submitted online and is presently being given full consideration for publication in the Food Science and Technology.

Your manuscript ID is CTA-2022-0991.R1.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at <https://mc04.manuscriptcentral.com/cta-scielo> and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to <https://mc04.manuscriptcentral.com/cta-scielo>.

Thank you for submitting your manuscript to the Food Science and Technology.

Sincerely,
Food Science and Technology Editorial Office



NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS

Journal:	<i>Food Science and Technology</i>
Manuscript ID	CTA-2022-0991.R1
Manuscript Type:	Original Article
Keyword:	Lutjanus sp., characteristics, food, fish bone

SCHOLARONE™
Manuscripts

NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS

Rahim Husain¹, Fernandy Djailani², Rita Marsuci Harmain³

1,2,3). Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesian

Email¹ : rahim@ung.ac.id

ABSTRACT

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using Association of Official Analytical Chemists (2005), and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: Lutjanus sp., characteristics, food, fish bone

Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om *et al.*, 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane *et al.*, 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah *et al.*, 2022). Cornmeal-based porridge in Nigeria is "Ugali" or "Ogi", in Italy "Polenta" in Brazil "Angu" and in Romania "Mamaliga". In this study, the traditional Gorontalo grits in Indonesia called "Saddah" or "Gohoge" is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Riddríguez *et al.*, 2019). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu *et al.*, 2022). According to CONAB (2015), in Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Filho *et al.*, 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear and branched structure (Liu *et al.*, 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal,

1
2
3
4 and rheological properties (Bello-Perez et al., 1999; Watershoot et al., 2015; Riddriguez
5 et al., 2018). Corn mainly supplies starch, and its flour is used in home cooking as a
6 food source and processed food products (Neupane et al., 2022, Barreti, et al., 2022).
7

8 Saddah Gohoge grits is a local name for food from the Gorontalo province in
9 Indonesia, which is generally popular, especially for the elderly and growing children,
10 because it has a soft texture and corn flavor. These grits contains high carbohydrates,
11 namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral
12 nutrient content. Mineral nutrient intake is needed, especially during the growth period
13 of children and the elderly. For the elderly aged >60 years, the calcium needed by the
14 body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237
15 mg/day (Regulation of the Minister of Health Number 43, 2019). Research in Rwanda
16 revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins
17 A and E. (Grosshagauer et.al., 2020).
18

19 Mineral content can be obtained from fish bone waste from processed fish
20 residue. Utilization of processed fish to produce new products can be done effectively
21 and increase income and contribute to environmental conservation. Processed fish waste
22 is divided into two groups, namely used to increase animal or plant production and the
23 development of value-added by-products for human consumption, namely stomach
24 contents, scales, bones including skull bones to be used as flour, oil, silage and compost
25 for use in animal feed or fertilizer in plants (Leira et.al., 2019). Fish bones have a fairly
26 high mineral content but are still in the form of macrominerals and have not been
27 completely absorbed by the body. The main constituents of fish bones as an important
28 source of minerals are calcium, sodium (5.63g/kg), phosphorus (2.38g/kg), and
29 carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium
30 phosphate, creatine phosphate and hydroxyapatite [Ca₁₀(OH)(PO₄)₆] in the form of
31 crystals attached to fibrillar collagen (Malde et al., 2010); Suptijah et al., (2012).
32

33 One of the processed fishery waste is fish bones from the *Lutjanus* sp. This
34 family of fish is found in Indonesian waters, including in the waters of the Gorontalo
35 area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types
36 of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas
37 Oktaviyani., 2012).
38

39 The particle size of minerals as feed additives in nanoparticle form is claimed to
40 be smaller than 100 nanometer, it can pass through the stomach wall and into body cells
41 more quickly than ordinary minerals with larger particle size (Singh&Pankaj., 2016).
42

43 The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah
44 Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can
45 be used as an inexpensive natural material that contains calcium phosphate. Previous
46 research produced calcium phosphate in micro and nanoparticle sizes from fish bones.
47 Fish bone waste originating from household consumers, processed fish fillet industries
48 into the aquatic environment can have an unfavorable effect. Utilization of fish bone
49 waste can be used as a product that provides beneficial value, especially in the growth
50 and maintenance of healthy bones and teeth. Fish bones derived from processed fish
51 fillet waste can be processed into nanominerals (nm) and can be fortified in food.
52 Through nanotechnology for application to food, one of them is increasing the
53 bioavailability of nutrients (Greiner&Ralf., 2009); Lekahena et al., (2014). One of the
54 additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals
55 derived from fish bone waste from the *Lutjanus* sp. This study carried out fortification
56 of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and
57
58
59
60

organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

Materials and Methods

This research was conducted in April 2022 – September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and Tools

Saddah gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family *Lutjanus* sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchoorl solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan and Kim, 2010). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah *et al.*, 2022).


2.2 Sample Preparation

2.2.1 Fish Bone Preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of \pm 130 °C or 250 °F, drying, reducing and screening of fish bone powder (Harmain *et.al.*, 2018). *modification*. The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi *et.al.*, 2018). Nanocalcium using the *precipitation* method with prawn shell soaking time for 48 hours (Suptijah *et al.*, 2012).

2.2.2 Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100 - 120 oC for 1 - 3 hours to dry or using a drum dryer at 120 oC at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2002) ; (Om *et.al.*, 2019) *modification* (Fig.1).



33 Fig. 1. Process flow for the production of maize flour

35 Formulation of composites of Saddah Gohoge corn with cooking water based
36 on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone
37 of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to
38 produce composite grits while the control sampel used was 100% Saddah Gohoge. **As**
39 **many as 200 g Saddah gohoge corn flour was mixed and homogenized with fortification**
40 **of fish bone nanominerals from the family Lutjanus sp. and then cooked with 1000 mL**
41 **(1:5) cooking water at a temperature of 100 °C for ± 45 minutes..**

44 2.3 Procedure of Analyses

45 2.3.1 Organoleptic Test

47 Samples of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus*
48 sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic
49 and hedonic quality organoleptic assessment score sheet, the panelists used were
50 untrained panelists and came from consumers who used to consume Saddah grits as
51 many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture,
52 color, aroma and taste through an assessment in the score sheet provided. Organoleptic
53 testing uses panelists who are divided into trained, semi-trained and untrained panelists
54 with a certain scale range based on an assessment of organoleptic parameters (Lawless
55 & Heymann, 2010).

58 2.3.2 Proximate analysis

Proximate analysis of Saddah gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemist (2006) method. The Association of Official Agricultural Chemists (2004) (Iwansyah et al., 2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster Association of Official Agricultural Chemists (2004) (Iwansyah et al., 2022), fat content using the Weibull method (Association of Official Agricultural Chemists (2004) (Iwansyah et al., 2022), carbohydrate content can be calculated by difference (Association of Official Analytical Chemist (2005), fiber content can be determined using the method of Association of Official Analytical Chemist (2006).

2.3.3 Viscosity measurement

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95°C for 5 minutes, then held at 95°C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

2.3.4 Coloring Analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L* a* b* and hue coordinates. The L* coordinates describe where L = 0 is black, and L* = 100 is colorless. The coordinates of a* represent red and green shadows, where a* > 0 indicates red and a* < 0 indicates green. The b* coordinates represent the blue and yellow tones, where b* > 0 indicates the intensity of the yellow color and b* < 0 indicates the blue color. Hue (h*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

2.3.5 Determination of mineral grade

Nanomaterial analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al., (2013), a sample of 20 – 30 g was dried in an oven at 105°C followed by calcination at 450°C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150°C to dissolve the ash. the remaining. Tejera et al., (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein and Shalaei (2016) determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple

range test using the *SPSS* programme Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments. (SPSS 2022) ; (Rao 2009).

Results and Discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah gohoge corn grits fortification of fish bone nanominerals from the family *Lutjanus* sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the scoresheet provided (Fig. 2).




Fig. 2 Serving of Saddah gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%)

3.1.1 Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Tab.1, Tab.2).

Tab 1. Hedonic quality organoleptic test results

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 \pm 1,041 ^a	3,40 \pm 0,913 ^a	3,08 \pm 0.909 ^a	3,64 \pm 1,075 ^a	3,16 \pm 1,248 ^a
F1 (90: 10)	3,68 \pm 0,945 ^a	3,28 \pm 1,021 ^a	2,88 \pm 0,881 ^a	4,00 \pm 0,866 ^a	2,96 \pm 1,207 ^a
F2 (95:15)	3,44 \pm 1,044 ^a	3,36 \pm 1,551 ^a	3,00 \pm 1,000 ^a	3,96 \pm 1,060 ^a	2,56 \pm 0,917 ^{ab}
F3 (80:20)	3,32 \pm 1,145 ^a	3,48 \pm 1,503 ^a	2,76 \pm 1,052 ^a	3,96 \pm 4.00 ^a	2,24 \pm 1,200 ^b

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the *Lutjanus* sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., (2018)).

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Tab.1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of

fish bone nanominerals *Lutjanus* sp. 10 % obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Tab. 1).

3.1.2 Hedonic analysis

Tab 2. Hedonic organoleptic test results

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3,96±0,841 ^a	3,64±1,114 ^a	4,04±0,935 ^a	3,80±1,041 ^a	3,52±1,194 ^a
F1 (90: 10)	3,64±0,995 ^a	3,36±1,319 ^a	3,96±0,889 ^a	3,64±0,907 ^a	3,28±1,100 ^{ab}
F2 (95:15)	2,96±0,790 ^b	3,12±0,927 ^a	3,76±0,879 ^a	2,96±0,935 ^b	2,80±0,913 ^{bc}
F3 (80:20)	2,80±1,000 ^b	2,84±1,143 ^a	3,40±1,118 ^a	2,88±1,013 ^b	2,56±1,044 ^c

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. 0% with a rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. on texture and color parameters but significant on appearance, flavor and taste (Tab.2). The texture parameter obtained the highest value at 0% concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al., (2018) which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1 – 4.9.

The texture of Saddah gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et.al., (2006); Rddriguez et al., (2018) that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al., (2018) on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al., (2018) showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3 – 4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

3.2.1 Viscosity Analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Tab 3).

Tab. 3. Viscosity levels of Saddah *gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
	F1	F2	F3
Viscosity levels (cP)	8309.4667 ±42.46285 ^a	14069.7333 ±103.87946 ^b	10264.7000 ±123.87946 ^c

The results of measuring the thickness of the sample of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Tab.3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al., (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022).). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., (2006); Palguna et al., (2014); Iwansyah et al., 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

3.2.2 Coloring Analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10 – 50.03 for the L component. different values for *a are -39.23 – 3.76 and *b values 15.96 – 16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Tab. 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

Tab. 4. Color of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10±2.60 ^a	51.33±1.01 ^a	50.03±3.55 ^a
*a	3.76±3.00 ^a	-39.23±74.27 ^a	2.03±1.74 ^a
*b	15.96±2.45 ^a	16.30±2.26 ^a	16.50±2.60 ^a

The results of the staining study were suspected because Saddah gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarfus fagifer* Forst.) (Wijanarka et al., (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal., 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

Tab. 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Karbohydate Content
0 %	86.82±0.411 ^a	0.92±0.87 ^a	7.16±0.90 ^a	0.47±0.95 ^a	1.77±0.54 ^a	95.37±0.50 ^a
15 %	85.03±0.744 ^b	1.13±0.26 ^{ab}	8.53±0.72 ^b	0.74±0.41 ^b	3.30±0.49 ^b	95.43±1.52 ^b
20 %	80.33±0.562 ^b	1.31±0.27 ^{ab}	9.63±0.47 ^c	1.21±0.87 ^c	3.60±0.37 ^b	92.48±0.40 ^c
25 %	79.70±0.113 ^c	1.53±0.25 ^b	10.51±0.30 ^d	1.22±1.36 ^c	3.8±0.37 ^b	92.96±0.61 ^d

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Tab.5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al., (2018) were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al., (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa

grits with pre-gelatinization with the highest value of water content at 80 oC, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p>0.05$) but significantly different from those without nanomineral fortification ($p<0.05$).

Hal ini sejalan dengan penelitian Iwansyah et al., (2022) who obtained statistical results that the ash content was not significant ($p>0.05$) with increasing temperature in pre-gelatinized instant rowe luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah gohoge grits were the same as those of Ahmad et al. (2018), which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al., (2022) were also significant ($p<0.05$) along with the increase in temperature in pre-gelatinated rowe luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., (2012); Jamilah et al., (2009): Iwansyah et al., (2022).

The fat content showed that the fortified concentration of 15% was significantly ($p<0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p>0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p>0.05$) with increasing pre-gelatinization temperature of rowe luwa grits.

For fiber content, the concentration of fortified 15 %, 20 % dan 25% was not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Tab.6).

Tab. 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10

2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphor (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Tab.6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al., (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. BPOM (2011) states that a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on *Saddah gohoge* grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgments

We would like to thank the Directorate General of Higher Education through the 2022 PDUPT Research, LP2M UNG, the reviewers of this article and all those who have helped carry out the research.

References

- Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 2013, 57(1), 18717. PMID:23516115. <http://dx.doi.org/10.3402/fnr.v57i0.18717>
- Akyıldız A., Ocal ND. Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 2006, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>
- Ahmad L., Une S., Bait Y. Suryani Une, and Yoyanda Bait. Characteristics of nutritional components, antioxidants, And organoleptic responses of Gorontalo traditional corn porridge with kersen (*Muntingia Calabura* L) leaf extract. *J agriTECH* 2019, 38(4): 463. ISSN 2527-3825.Doi: <http://doi.org/10.22146/agritech.24739>
- Association of Official Agricultural Chemists – AOAC. 2004. Official methods of analysis (15th ed, vol.1). Arington:AOAC inc.<https://doi.org/10.1016/b0-12-765490-9/00006-9>
- Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3) 854-857. 1999. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.
- Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Filho, M. A. S. C., Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19): 1-7. <https://doi.org/10.1590/fst.33120>
- Couto A., Enes P., Peres H., Oliva-Teles A. Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *J Fish Physiology and Biochemistry*, 2012, 38(3), 595-601. PMID:21728054. <http://dx.doi.org/10.1007/s10695-011-9537-5>
- Ciobanu C., Slencu BG., Cucureanu., Faas R. Determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiș"*. Seria Științele Vieții., 2013. v. 23, n. 1, p. 33-38
- Filho IK., Butik M., Jaski AC., Quinaia SP. Fast method to determine the elements in maize flour:reduction in preparation time and reagent consumption. *Braz J. Food Technology*, Campinas.v.21,e2017091,2018. <https://doi.org/10.1590/1981-67239117>
- Greiner, Ralf. 2009. Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício." *Jl of Brazilian Society of Food and Nutrition* 34(1): 243–60. http://sban.cloudpanel.com.br/files/revistas_publicacoes/228.pdf.
- Grosshagauer S., Milani P., Kraemer K., Mukabutera A., Burkon A., Pignitter M., Bayer S., Somoza V. Inadequacy of nutrients and contaminants found in porridge-type

1
2
3 complementary foods in Rwanda. *Matern Child Nutr.* 2020; 16(1): e12856. Published
4 online 2019 Jul 15. Doi: 10.1111/mcn.12856
5

6
7 Houssou P, Ayemor GS. Appropriate processing and food functional property of maize
8 flour. *Afr J Sci Technol*, 2012. 3: 126-131.
9

10
11 Harmain RM., Dali FA., Husain R. Nanocalcium characterization of cakalang fish bone flour
12 (*Katsuwonus pelamis* L). *J of Innovative Science and Reasearch Technology*,
13 2018,3,(10): 306–8, IJISRT 180C148, ISSN 2456-2165
14

15
16 Iwansyah AC., Apriadi T., Arif DZ., Andriana Y., Indriati A., Mayasti NI., Luthfiyanti R.
17 *Braz.J. Food Technology*, Campinas, v. 25, e2021050,2022.
18 <https://dio.org/10.1590/1981-6723.05021>
19

20
21 Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N.
22 Characterisation of physical, mechanical and colour properties of *Muntingia calabura*
23 fruits. In S. K. Wahono (Ed.), *IOP Conference Series: Earth and Environmental*
24 *Science: 2020, Vol. 462, No. 1.* Tangerang: IOP Publishing.
25 <http://dx.doi.org/10.1088/1755-1315/462/1/012044>
26

27
28 Jamilah B., Mohamed A., Abbas KA., Rahman RA., Karim R., & Hashim DM. Protein-starch
29 interaction and their effect on thermal and rheological characteristics of a food system: a
30 review. *J of Food Agriculture and Environment*, 2009, 7(2), 169-174.
31

32
33 Lekahena, Vanessa, Didah Nur Faridah, Rizal Syarief, and R. Peranginangin.
34 Physicochemical characterization of nanocalcium extracted from Tilapia fish bone
35 using alkaline and acid solution. *J Teknol dan Industri Pangan*, 2014, 25(1): 57–64,
36 ISSN: 1979-7788, Doi:10.6066/jtip.2015.25.1.57
37

38
39 Lawless, H. T., & Heymann, H. (2010a). *Principles of Good Practice Sensory Evaluation of*
40 *Food.* (pp. 57–77). Springer. 10.1007/978-1-4419-6488-5-3 .
41

42
43 Lawless, H. T., & Heymann, H. (2010b). *Sensory Evaluation of Food: Principles and*
44 *Practices.* Springer Science & Business Media. 10.1007/978-1-4419-6488-5
45

46
47 Liu, Q., Charlet, G., Yelle, S., & Arul, J. Phase transition in potato starch–water system 0.
48 Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-
49 407. 2002. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).
50

51
52 Malde MK., Bugel S., Kristensen M., Malde K., Graff IE., Pedersen JI. 2010. Calcium from
53 salmon and cod bone is well absorbed in young healthy men: a double-blinded
54 randomised crossover design. *Nutrition&Metabolism* 2010,
55 7:61.<http://www.nutritionandmetabolism.com/content/7/1/61>
56

57
58 Neupane D., Adhikari P., Bhattarai D., Rana B., Ahmed Z., Sharma U., Adhikari D. Does
59 Climate Change Affect the Yield of the Top Three Cereals and Food Security in the
60 World?. *Earth*, 3, 45-71. 2022. <https://doi.org/10.3390/earth3010004>

1
2
3 Official Methods of Analysis of AOAC International. 18th Edition 2005, Current trough
4 Revision 1,2006. Published by AOAC International SUITE 500,481 North frederick
5 Avenue Gaithersburg,Maryland 20877-2417,USA.ISBN 0-935584-77-3
6

7
8 OM A., AK B., FO O., GE Z. Nutritional Composition and Organoleptic Properties of
9 Composite Maize Nutritional Composition and Organoleptic Properties of Composite
10 Maize Porridge. *Food Processing & Technology*. 10-6. 2019. [https://doi: 10.4172/2157-](https://doi.org/10.4172/2157-7110.1000798)
11 [7110.1000798](https://doi.org/10.4172/2157-7110.1000798)
12

13
14 Oktaviyani S. Knowing the Lutjanus, one of the leading commodities in catch fisheries. *J*
15 *Oseana*, vol XL(III),number 3, 2018:29-39, ISSN 0216-1877
16

17
18 Palguna IGPA., Sugiyono S., Hariyanto B. Characteristics of modified sago starch by
19 repeated gelatinization and retrogradation treatment. *J Pangan*, 2014, 23(6),146-156
20

21
22 Rddriguez JC.,Gallegos JMT., Garcia MO., Flores PBZ.,Paz JdJ O., Muniz CHA., Morales
23 GG., Calderon DEP., Velasco CR. Physicochemical, thermal and rheological properties
24 of three native corn starches. *Braz. J Food Science and Technology*, Campinas,
25 39(1):149-157, Jan.-Mar.2019. DOI: <https://doi.org/10.1590/fst.28117>

26
27 Regulation of the Minister of Health Number 43 of 2019. Regulation of the Minister of
28 Health regarding Communitie Health Centers.28 oct 2019. BN.2019/NO.1335,
29 PERATURAN.GO.ID : 40 HLM.
30 <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019>
31

32
33 Suptijah P., Jacob AM., Deviyanti N. Characterization and bioavailability of vannamei
34 shrimp shell nanocalcium. *J Akuatika*, vol.III, 1/Mar 2012 (63-73), ISSN 0853-2523
35

36
37 Singh., Pankaj. Use of Nano Feed Additives in Livestock Feeding. *Intern J of Livestock*
38 *Research* 6(1):1.2016.

39
40 Tejera RL., Luis G., Gonzales-Weller., D;Caballero JM., Gutierrez AJ., Rubio C., Hardisson
41 A.
42 Metals in wheat flour; comparative study and safety control. *Nutricion*
43 *Hospitalaria*, v. 28, n. 2, p. 506-513, 2013. PMid:23822705.

44
45 Rao KV, Radhaiah G, Balakrishna N. Tests of inference In: Auo, K.V. (Ed.) *Biostatistics-A*
46 *manual of statistical methods for use in health, nutrition and anthropology*. Jaypee
47 *Brothers*, New Delhi, India, 2009, pp: 126-168.

48
49 SPSS, Statistical Package for Social Sciences. SPSS 1001 for windows.
50 <https://www.ibm.com/support/pages/spss-statistics-210-available-download>. 2022.
51 SPSS Inc., USA.
52

53
54 Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. Relationships between
55 physicochemical, morphological, thermal, rheological properties of rice starches. *Food*
56 *Hydrocolloids*, 2006, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>
57
58
59
60

- 1
2
3
4
5 Wadchararat C., Masubon T., Naivikul O. Characterization of pregelatinized and heat
6 moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 2006, 40, 144-153
7
8
9 Wijanarka A., Sudargo T., Harmayani E., Marsono Y. Effect of pre-gelatinization on
10 physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour.
11 *American Journal of Food Technology*, 2017, 12(3), 178-185.
12 <http://dx.doi.org/10.3923/ajft.2017.178.185>
13
14 Riyanto B., Trilaksani W., Azzahra VA. 2020. Design of Instant Food Special Nutrition for
15 the Elderly Based on Binte Biluhuta Enriched with Fish Bone Nanominerals. *J*
16 *Fishtech*, 2020, 9(2): 65–77.
17
18
19 Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure,
20 physicochemical and functional properties of maize, cassava, wheat, potato and rice
21 starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.
22
23
24 Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid,
25 and volatile composition in waxy and non-waxy proso millet. *Food Science and*
26 *Technology (Campinas)*, 42(19): 1-9. <https://doi.org/10.1590/fst.35121>
27
28
29 Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage
30 detection of direct maize kernel harvest based on image processing and BP neural
31 network. *Food Science and Technology (Campinas)*, 42(19): 1-9.
32 <https://doi.org/10.1590/fst.54322>
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Manuscript Decision (Tahap 2)

(7 November 2022)



Rahim Husain <rahim@ung.ac.id>

Food Science and Technology - Decision on Manuscript ID CTA-2022-0991.R1

Adriano Cruz <onbehalfof@manuscriptcentral.com>

7 November 2022 pukul 07.17

Balas Ke: adriano.cruz@ifrj.edu.br

Kepada: rahim@ung.ac.id

06-Nov-2022

Dear Dr. Husain:

It is a pleasure to accept your manuscript entitled "NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS" in its current form for publication in the Food Science and Technology. The comments of the reviewer(s) who reviewed your manuscript are included at the foot of this letter.

Thank you for your fine contribution. On behalf of the Editors of the Food Science and Technology, we look forward to your continued contributions to the Journal.

Sincerely,

Dr. Adriano Cruz

Editor-in-Chief, Food Science and Technology

adriano.cruz@ifrj.edu.br

Associate Editor

Comments to the Author:

(There are no comments.)

Entire Scoresheet:

Letter of Acceptance

(3 Maret 2023)



Rahim Husain <rahim@ung.ac.id>

**ENC: Food Science and Technology - Decision on Manuscript ID
CTA-2022-0991.R1**

Revista <revista@sbcta.org.br>
Kepada: rahim@ung.ac.id

3 Maret 2023 pukul 20.30

Dear,

Follow.

De: Adriano Gomes da Cruz [mailto:adriano.cruz@ifrj.edu.br]**Enviada em:** quinta-feira, 2 de março de 2023 23:31**Para:** Revista**Assunto:** Fwd: Food Science and Technology - Decision on Manuscript ID CTA-2022-0991.R1

----- Mensagem encaminhada -----

De: **Rahim Husain** <rahim@ung.ac.id>

Data: quinta-feira, 2 de março de 2023

Assunto: Re: Food Science and Technology - Decision on Manuscript ID CTA-2022-0991.R1

Para: adriano.cruz@ifrj.edu.br

Dear Editorial Team,

I hope this email finds you well. Since my article has been accepted for publication in your journal, may I request for the letter of acceptance from the journal? I really need it for my promotion.

Thank you and I look forward to hearing from you.

Best regards,

Rahim Husain

Pada tanggal Sen, 7 Nov 2022 pukul 08.18 Adriano Cruz <onbehalfof@manuscriptcentral.com> menulis:

| 06-Nov-2022

Dear Dr. Husain:

It is a pleasure to accept your manuscript entitled "NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS" in its current form for publication in the Food Science and Technology. The comments of the reviewer(s) who reviewed your manuscript are included at the foot of this letter.

Thank you for your fine contribution. On behalf of the Editors of the Food Science and Technology, we look forward to your continued contributions to the Journal.

Sincerely,
Dr. Adriano Cruz
Editor-in-Chief, Food Science and Technology
adriano.cruz@ifrj.edu.br

Associate Editor
Comments to the Author:
(There are no comments.)

Entire Scoresheet:

--

Prof. Dr. Adriano Gomes da Cruz

Departamento de Alimentos / Department of Food

Instituto Federal de Educação, Ciência e Tecnologia do Rio de Janeiro (IFRJ) / Federal Institute of Education, Science and Technology of Rio de Janeiro (IFRJ)



991-22.pdf

43K



Sociedade Brasileira de Ciência e Tecnologia de Alimentos

Orgão de Utilidade Pública
Lei Municipal No. 4919 de 31/08/1979 - Fundada em 08/04/1967

Av. Brasil, 2880
Caixa Postal 271
13001-970 - Campinas/SP

C.G.C: 46.113.742/0001-24
Fone/Fax: (019)3241-0527
Fone: (019)3241-5793

Campinas, Mar 03th, 2023

Article Acceptance

We are pleased to inform you that the article "**NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS**", CTA 991-22, has been accepted for publication in the Journal "Food Science and Technology".

Authors:

Rahim Husain ¹, Fernandy Djailani ², Rita Marsuci Harmain ³
1,2,3). Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science,
Universitas Negeri Gorontalo, Gorontalo Province, Indonesian
Email ¹ : rahim@ung.ac.id

Congratulations on the acceptance of your paper

Sincerely,

Adriano Gomes da Cruz

Editor-in-chief/Food Science and Technology

Artikel Proofreading (Tahap 1)

(9 Maret 2023)



Rahim Husain <rahim@ung.ac.id>

cta | Final proofreading | Article: ctaAO099122_EN #CPub - Editora Cubo#

CPub <no-reply@editoracubo.com.br>
Kepada: rahim@ung.ac.id

9 Maret 2023 pukul 02.48

CPub

This notification is addressed to: <rahim@ung.ac.id>

Food Science and Technology

You have a new message regarding document:

[ctaAO099122_EN] NANOMINERAL FORTIFICATION ON TRADITIONAL
SADDAH GOHOGE GRITS

To view the message you need to sign in CPub using the link below:



Access CPub

Regards,
CPub

If you're having trouble clicking the "Access CPub" button, copy and paste the URL below into your web browser: https://pub.editoracubo.com.br/publicacoes/cta/documentos/ctaAO099122_EN/mensagens/erZygO4fzvE1iS2ihPUptgpfYIR7VK0XK3jw3IPM51fUsc5ZVnS8JkvDqZFcTHG

Subject	cta Final proofreading Article: ctaAO099122_EN
Sent at	03/09/2023 03:48
Read at	03/09/2023 09:59
Message	<p>Dear Author,</p> <p>You are receiving a .DOCX file with the final proofreading of your article before it is published in the Food Science and Technology. Please follow the instructions located in the first page of the attached file.</p> <p>INSTRUCTIONS:</p> <ul style="list-style-type: none">• The attached .DOCX file should only be opened on versions of MS Office 2010 or higher. After opening the file and before starting your review, make sure that there are no missing spaces like this or this in the text, tables and references. Failure to follow these guidelines could cause the loss of your work and irreversibly corrupt the file. [1]• We will resume working on your article only after receiving the revised document with all issues properly resolved.• No changes in the text will be possible after this step. Make sure that all changes in the content have been made on the revised .DOCX file that you will return us.• To return your corrections, please attach the revised file in the reply to this email.• In the next production step you will receive a PDF proof, in which only layout aspects of the text, figures, tables etc., should be observed. <p>To ensure fast publication of your article reply this message within 2 days.</p> <p>[1] Microsoft© has reported a problem with version 2007 of MS Word© which causes a random loss of spaces between some words when opening the new .DOCX standard file format: http://support.microsoft.com/kb/2633585</p> <p>Best regards, Desktop Publishing Team</p>

Attachments	<ul style="list-style-type: none">•  ctaAO099122_EN_enviadoAutor.docx 85.7 KB 03/09/2023 03:48
--------------------	---

IMPORTANT

- **BEFORE you begin your review** make sure that there are no missing spaces like this or this in the text, tables and references.
- If you fail to follow this instruction you might lose your work.

Review Guidelines

1. Carefully review and make all final text adjustments directly in this Word document.
2. Resolve all issues listed in the "To Authors" section directly in this Word document.
3. When you are done make sure you answer the **Author Confirmation Checklist** and return this Word document by e-mail.
4. The issues listed in the "To Editors" section will be resolved by the journal.

Important:

- Check if the spelling and the order of presentation of article authors name are correct.
- Do not remove or change text styles (font size and color) and color highlights for they are part of our production workflow.
- Figures (if any) are NOT included in this file for at this moment you are required to only and carefully review the content (text and data) of your article. Figures if present in your article will be reviewed in the PDF galley proof.
- All edits must be made directly in this Word document. Requests sent by e-mail or in separate documents will not be accepted and will cause delay in the publication.
- If you need to save the received file, please do not change its extension, **keep it safe in DOCX format.**

The following changes are not allowed and will be taken to the Editor-in-Chief's attention:

- Adding, removing or reordering the Author's list.
- Significantly changing large amounts of text or in tables data.

Issues

To Authors:

[[Q1: Authors: Please send us the running title Q1]]

[[Q2: It is recommended not to present citations in the abstract. If necessary, adapt the text and include the citation in the body of the text. Q2]]

[[Q3: Authors: Please send us the practical application Q3]]

[[Q4: The in-text citation "Ridriguez et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q4]]

[[Q5: The in-text citation "CONAB (2015)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q5]]

[[Q6: The citation "Bello-Perez et al., 1999" matches the reference

"Bello-Pérez et al, 1999", but an accent or apostrophe is different. Q6]]

[[Q7: The in-text citation "Ridriguez et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q7]]

[[Q8: The in-text citation "Leira et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q8]]

[[Q9: The in-text citation "Oktaviyani (2012)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q9]]

[[Q10: The in-text citation "Singh & Pankaj, 2016" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q10]]

[[Q11: The in-text citation "Greiner & Ralf, 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q11]]

[[Q12: The in-text citation "Venkatesan & Kim, 2010" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q12]]

[[Q13: The in-text citation "Prinaldi et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q13]]

[[Q14: The in-text citation "Houssou & Ayemor (2002" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q14]]

[[Q15: The citation "Lawless & Heymann, 2010" matches multiple references. Please add letters (e.g. "Smith 2000a"), or additional authors to the citation, to uniquely match references and citations. Q15]]

[[Q16: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q16]]

[[Q17: The in-text citation "Association of Official Analytical Chemist, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q17]]

[[Q18: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Q18]]

[[Q19: The in-text citation "Pourhossein & Shalaei (2016)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q19]]

[[Q20: The in-text citation "Rao 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q20]]

[[Q21: The in-text citation "Ahmad et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q21]]

[[Q22: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q22]]

[[Q23: The in-text citation "Rodriguez et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q23]]

[[Q24: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q24]]

[[Q25: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q25]]

[[Q26: The in-text citation "Makame et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q26]]

[[Q27: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q27]]

[[Q28: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q28]]

[[Q29: The in-text citation "BPOM (2011)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q29]]

[[Q30: Reference "Ahmad, Une, Bait, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q30]]

[[Q31: Reference "Association of Official Agricultural Chemists, 2005" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q31]]

[[Q32: Reference "Bustillos-Rodríguez, Tirado-Gallegos, Ordóñez-García, Zamudio-Flores, Ornelas-Paz, Acosta-Muñiz, Gallegos-Morales, Páramo-Calderón, Rios-Velasco, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q32]]

[[Q33: Reference "Greiner, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q33]]

[[Q34: Reference "Houssou, Ayemor, 2012" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q34]]

[[Q35: Send the title of the article Q35]]

[[Q36: Reference "Lawless, Heymann, 2010a" is cited in the text, but the citation matches multiple references. Please check the reference and the citations. Q36]]

[[Q37: Reference "Lawless, Heymann, 2010b" is cited in the text, but the citation matches multiple references. Please check the reference and the citations. Q37]]

[[Q38: Reference "Oktaviyani, 2018" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q38]]

[[Q39: Send the title of the period in full, without abbreviations Q39]]

[[Q40: Send the title of the period in full, without abbreviations Q40]]

[[Q41: Reference "Rao, Radhaiah, Balakrishna, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q41]]

[[Q42: Send the name of all the authors of the publication Q42]]

[[Q43: Send the title of the period in full, without abbreviations Q43]]

[[Q44: Please confirm if the use of the decimal and thousands markers in values of the Table 1-3 is correct (in English dot is used as decimal marker and comma as thousands marker, ex.: 5,931.10, while in Portuguese language comma is used as decimal and dot as thousands marker, ex.: 5.931,10). If necessary make the corrections in the Tables 1-3. Q44]]

[[Q45: Send the explanatory note to the superscript letters present in the table values. Q45]]

[[Q46: Send the explanatory note to the superscript letters present in the table values. Q46]]

[[Q47: Send the explanatory note to the superscript letters present in the table values. Q47]]

[[Q48: Send the explanatory note to the superscript letters present in the table values. Q48]]

Author Confirmation Checklist:

(Please mark with an X and add comments if needed)

Were ALL issues resolved?

Yes: ____

No: ____

Author comments:

10.1590/fst.099122

Running Title [[Q1: Q1]]

Husain et al.

Original Article

e099122

43

2023

xxx.-xxx.

000

000

0

This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nanomineral fortification on traditional saddah gohoge grits

Rahim HUSAIN^{1*} <https://orcid.org/0000-0001-5307-9806>, Fernandy DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using [Association of Official Analytical Chemists \(2005 \[\[Q2: Q2\]\]\)](#), and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*.

Comment [AB1]: Authors:
Please send us the running title

Comment [AB2]: It is recommended not to present citations in the abstract. If necessary, adapt the text and include the citation in the body of the text.

Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application [[Q3: Q3]]:

Received 23 Sept., 2022

Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesian

*Corresponding author: rahim@ung.ac.id

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is "Ugali" or "Ogi", in Italy "Polenta" in Brazil "Angu" and in Romania "Mamaliga". In this study, the traditional Gorontalo grits in Indonesia called "Saddah" or "Gohoge" is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Ridriguez et al., 2019 [[Q4: Q4]]). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). According to CONAB (2015) [[Q5: Q5]], in Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Perez et al., 1999 [[Q6: Q6]]; Waterschoot et al., 2015; Ridriguez et al., 2018 [[Q7: Q7]]). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Leira et al., 2019 [[Q8: Q8]]). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite [Ca₁₀(OH)(PO₄)₆] in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas Oktaviani (2012) [[Q9: Q9]].

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016 [[Q10: Q10]]).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating

Comment [AB3]: Authors:
Please send us the practical application

Comment [AB4]: The in-text citation "Ridriguez et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB5]: The in-text citation "CONAB (2015)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB6]: The citation "Bello-Perez et al., 1999" matches the reference "Bello-Pérez et al., 1999", but an accent or apostrophe is different.

Comment [AB7]: The in-text citation "Ridriguez et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB8]: The in-text citation "Lei et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB9]: The in-text citation "Oktaviani (2012)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB10]: The in-text citation "Singh & Pankaj, 2016" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner & Ralf, 2009 [[Q11: Q11]]; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family *Lutjanus* sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchoorl solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010 [[Q12: Q12]]). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder (Harmain et al., 2018). *modification*. The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018 [[Q13: Q13]]). Nanocalcium using the *precipitation* method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2002 [[Q14: Q14]]; Om et al., 2019) *modification* (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic quality organoleptic assessment score sheet, the panelists used were untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010 [[Q15: Q15]]).

Proximate analysis

Proximate analysis of Saddah gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemist (2006) [[Q16: Q16]] method. The Association of Official Agricultural Chemists (2004), Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used

Comment [AB11]: The in-text citation "Greiner & Ralf, 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB12]: The in-text citation "Venkatesan & Kim, 2010" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB13]: The in-text citation "Prinaldi et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB14]: The in-text citation "Houssou & Ayemor (2002)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB15]: The citation "Lawless & Heymann, 2010" matches multiple reference. Please add letters (e.g. "Smith 2000a"), or additional authors to the citation, to uniquely match references and citations.

Comment [AB16]: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

equipment Buchi-Dumaster Association of Official Agricultural Chemists (2004), Iwansyah et al. (2022), fat content using the Weibull method (Association of Official Agricultural Chemists, 2004; Iwansyah et al., 2022), carbohydrate content can be calculated by difference (Association of Official Analytical Chemist, 2005) [[Q17: Q17]], fiber content can be determined using the method of Association of Official Analytical Chemist (2006) [[Q18: Q18]].

Viscosity measurement

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L* a* b* and hue coordinates. The L* coordinates describe where L = 0 is black, and L* = 100 is colorless. The coordinates of a* represent red and green shadows, where a* > 0 indicates red and a* < 0 indicates green. The b* coordinates represent the blue and yellow tones, where b* > 0 indicates the intensity of the yellow color and b* < 0 indicates the blue color. Hue (h*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomineral analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. the remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) [[Q19: Q19]] determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) (n = 4). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao 2009) [[Q20: Q20]].

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the scoresheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the Lutjanus sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018) [[Q21: Q21]].

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals Lutjanus sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah gohoge grits fortification of fish bone nanominerals Lutjanus sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that

Comment [AB17]: The in-text citation "Association of Official Analytical Chemist, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB18]: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB19]: The in-text citation "Pourhossein & Shalaei (2016)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB20]: The in-text citation "Rao 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB21]: The in-text citation "Ahmad et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. 0% with a rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0% concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) [Q22: Q22] which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) [Q23: Q23] that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) [Q24: Q24] on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) [Q25: Q25] showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019 [Q26: Q26]). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are -39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

The results of the staining study were suspected because Saddah gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarpus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

Comment [AB22]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Comment [AB23]: The in-text citation "Rodriguez et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Comment [AB24]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Comment [AB25]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Comment [AB26]: The in-text citation "Makame et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) [Q27: Q27] were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

Hal ini sejalan dengan penelitian Iwansyah et al. (2022) who obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant rowe luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah gohoge grits were the same as those of Ahmad et al. (2018) [Q28: Q28], which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinized rowe luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of rowe luwa grits.

For fiber content, the concentration of fortified 15%, 20% dan 25% was not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. BPOM (2011) [Q29: Q29] states that a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on Saddah gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick,

Comment [AB27]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB28]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [AB29]: The in-text citation "BPOM (2011)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Directorate General of Higher Education through the 2022 PDUPT Research, LP2M UNG, the reviewers of this article and all those who have helped carry out the research.

References

<jrm>[[Q30: Q30]]Ahmad, L., Une, S., & Bait, Y. (2019). Characteristics of nutritional components, antioxidants, and organoleptic responses of Gorontalo traditional corn porridge with kersen (*Muntingia Calabura* L) leaf extract. *Agritech*, 38(4), 463. <http://doi.org/10.22146/agritech.24739>.</jrm>

<jrm>Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.</jrm>

<jrm>Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.</jrm>

<bok>Association of Official Agricultural Chemists – AOAC. (2004). *Official methods of analysis of the Association of Official Analytical Chemists* (15th ed., Vol. 1). Arlington: AOAC International.</bok>

<bok>[[Q31: Q31]]Association of Official Agricultural Chemists – AOAC. (2005). *Current trough Revision 1* (18th ed.). Arlington: AOAC International.</bok>

<jrm>Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.</jrm>

<jrm>Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.</jrm>

<jrm>[[Q32: Q32]]Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñiz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.</jrm>

<jrm>Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.</jrm>

Comment [AB30]: Reference "Ahmad, Une, & Bait, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Comment [AB31]: Reference "Association of Official Agricultural Chemists, 2005" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Comment [AB32]: Reference "Bustillos-Rodríguez, Tirado-Gallegos, Ordóñez-García, Zamudio-Flores, Ornelas-Paz, Acosta-Muñiz, Gallegos-Morales, Páramo-Calderón, Rios-Velasco, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted.

<jrn>Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.</jrn>

<jrn>[[Q33: Q33]]Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpaine.com.br/files/revistas_publicacoes/228.pdf</jrn>

Comment [AB33]: Reference "Greiner, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted.

<jrn>Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.</jrn>

<jrn>Harmain, R. M., Dali, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.</jrn>

<jrn>[[Q34: Q34]]Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.</jrn>

Comment [AB34]: Reference "Houssou, Ayemor, 2012" is not cited in the text. Please add an in-text citation or the reference will be deleted.

<gov>Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communitie Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335, PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019></gov>

<jrn>Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.</jrn>

<jrn>Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. I., & Luthfiyanti, R. (2022). title [[Q35: Q35]]. *Brazilian Journal of Food Technology*, 25, e2021050.</jrn>

Comment [AB35]: Send the title of the article

<jrn>Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.</jrn>

<jrn>Kelte, I., Fo., Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.</jrn>

<bok> [[Q36: Q36]]Lawless, H. T., & Heymann, H. (2010a). *Principles of good practice sensory evaluation of food* (pp. 57-77). USA: Springer. http://dx.doi.org/10.1007/978-1-4419-6488-5_3.</bok>

Comment [AB36]: Reference "Lawless, Heymann, 2010a" is cited in the text, but the citation matches multiple references. Please check the reference and the citations.

<bok> [[Q37: Q37]]Lawless, H. T., & Heymann, H. (2010b). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5></bok>

Comment [AB37]: Reference "Lawless, Heymann, 2010b" is cited in the text, but the citation matches multiple references. Please check the reference and the citations.

<jrn>Lekahena, V., Faridah, D. N., Syarif, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.</jrn>

<jrn>Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch–water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).</jrn>

<jrn>Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. PMID:20646299.</jrn>

<jrn>Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.</jrn>

<jrn>[[Q38: Q38]]Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *J Oseana* [[Q39: Q39]], 40(3), 29-39.</jrn>

Comment [AB38]: Reference "Oktaviyani 2018" is not cited in the text. Please add an in-text citation or the reference will be deleted.

<jrn>Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.</jrn>

Comment [AB39]: Send the title of the period in full, without abbreviations

<jrn>Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *J Pangan* [[Q40: Q40]], 23(6), 146-156.</jrn>

Comment [AB40]: Send the title of the period in full, without abbreviations

<edb> [[Q41: Q41]]Rao, K. V., Radhaiah, G., & Balakrishna, N. (2009). Tests of inference In K. V. AUC (Ed.), *Biostatistics-A manual of statistical methods for use in health, nutrition and anthropology* (pp. 126-168). New Delhi: Jaypee Brothers.</edb>

Comment [AB41]: Reference "Rao, Radhaiah, Balakrishna, 2009" is not cited in text. Please add an in-text citation or the reference will be deleted.

<jrn>Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.</jrn>

<jrn>Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.</jrn>

<jrn> [[Q42: Q42]]Singh, P. Use of nano feed additives in livestock feeding. *International Journal of Livestock Research*, 6(1), 1.</jrn>

Comment [AB42]: Send the name of all the authors of the publication

<eref> Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download></eref>

<jrn>Suptijah, P., Jacob, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *J Akuatika* [[Q43: Q43]], 3(1), 63-73.</jrn>

Comment [AB43]: Send the title of the period in full, without abbreviations

<jrn>Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. [Pmid:23822705](https://pubmed.ncbi.nlm.nih.gov/23822705/).</jrn>

<jrn>Wadchararat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.</jrn>

<jrn>Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. [http://dx.doi.org/10.1002/star.201300238](https://doi.org/10.1002/star.201300238).</jrn>

<jrn>Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. [http://dx.doi.org/10.3923/ajft.2017.178.185](https://doi.org/10.3923/ajft.2017.178.185).</jrn>

<jrn>Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. [http://dx.doi.org/10.1590/fst.58320](https://doi.org/10.1590/fst.58320).</jrn>

<jrn>Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. [http://dx.doi.org/10.1590/fst.54322](https://doi.org/10.1590/fst.54322).</jrn>

Table 1. Hedonic quality organoleptic test results [[Q44: Q44]].

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1,041 ^a	3.40 ± 0,913 ^a	3.08 ± 0,909 ^a	3,64 ± 1,075 ^a	3,16 ± 1,248 ^a
F1 (90:10)	3,68 ± 0,945 ^a	3,28 ± 1,021 ^a	2,88 ± 0,881 ^a	4,00 ± 0,866 ^a	2,96 ± 1,207 ^a
F2 (95:15)	3,44 ± 1,044 ^a	3,36 ± 1,551 ^a	3,00 ± 1,000 ^a	3,96 ± 1,060 ^a	2,56 ± 0,917 ^{ab}
F3 (80:20)	3,32 ± 1,145 ^a	3,48 ± 1,503 ^a [[Q45: Q45]]	2,76 ± 1,052 ^a	3,96 ± 4,00 ^a	2,24 ± 1,200 ^b

Comment [AB44]: Please confirm if the use of the decimal and thousands marker in values of the Table 1-3 is correct (in English dot is used as decimal marker and comma as thousands marker, ex.: 5,931.10, while in Portuguese language comma is used as decimal and dot as thousands marker, ex.: 5.931,10). If necessary make the corrections in the Tables 1-3.

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3,96 ± 0,841 ^a [[Q46: Q46]]	3,64 ± 1,114 ^a	4,04 ± 0,935 ^a	3,80 ± 1,041 ^a	3,52 ± 1,194 ^a
F1 (90:10)	3,64 ± 0,995 ^a	3,36 ± 1,319 ^a	3,96 ± 0,889 ^a	3,64 ± 0,907 ^a	3,28 ± 1,100 ^{ab}
F2 (95:15)	2,96 ± 0,790 ^b	3,12 ± 0,927 ^a	3,76 ± 0,879 ^a	2,96 ± 0,935 ^b	2,80 ± 0,913 ^{bc}
F3 (80:20)	2,80 ± 1,000 ^b	2,84 ± 1,143 ^a	3,40 ± 1,118 ^a	2,88 ± 1,013 ^b	2,56 ± 1,044 ^c

Comment [AB45]: Send the explanatory note to the superscript letters present in the table values.

Comment [AB46]: Send the explanatory note to the superscript letters present in the table values.

Table 3. Viscosity levels of Saddah *gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
Viscosity levels (cP)	F1	F2	F3
	8309.4667 ± 42,46285 ^a [[Q47: Q47]]	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

Comment [AB47]: Send the explanatory note to the superscript letters present in the table values.

Table 4. Color of Saddah *Gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2,60 ^a [[Q48: Q48]]	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a

Comment [AB48]: Send the explanatory note to the superscript letters present in the table values.

*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

Table 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Karbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d




Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphor (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

Figure 1. Process flow for the production of maize flour.

Figure 2. Serving of Saddah gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

**Pengiriman Revisi
Proofreading (Tahap 1)
(16 Maret 2023)**

Autor	RE: cta Final proofreading Article: ctaAO099122_EN  Read at 03/16/2023 20:15	03/16/2023 10:57  Prova de Conteúdo
Subject	RE: cta Final proofreading Article: ctaAO099122_EN	
Sent at	03/16/2023 10:57	
Read at	03/16/2023 20:15	
Message	<p>Dear Editorial Team,</p> <p>Attached is the revised version of the manuscript. Thank you and I look forward to hearing from you.</p> <p>Best regards,</p> <p>Rahim Husain</p>	
Attachments	<ul style="list-style-type: none"> ctaAO099122_EN_enviadoAutor.docx 129.6 KB 03/16/2023 10:57	

 reply

IMPORTANT

- **BEFORE you begin your review** make sure that there are no missing spaces like this or this in the text, tables and references.
- If you fail to follow this instruction you might lose your work.

Review Guidelines

1. Carefully review and make all final text adjustments directly in this Word document.
2. Resolve all issues listed in the "To Authors" section directly in this Word document.
3. When you are done make sure you answer the **Author Confirmation Checklist** and return this Word document by e-mail.
4. The issues listed in the "To Editors" section will be resolved by the journal.

Important:

- Check if the spelling and the order of presentation of article authors name are correct.
- Do not remove or change text styles (font size and color) and color highlights for they are part of our production workflow.
- Figures (if any) are NOT included in this file for at this moment you are required to only and carefully review the content (text and data) of your article. Figures if present in your article will be reviewed in the PDF galley proof.
- All edits must be made directly in this Word document. Requests sent by e-mail or in separate documents will not be accepted and will cause delay in the publication.
- If you need to save the received file, please do not change its extension, **keep it safe in DOCX format.**

The following changes are not allowed and will be taken to the Editor-in-Chief's attention:

- Adding, removing or reordering the Author's list.
- Significantly changing large amounts of text or in tables data.

Issues

To Authors:

[[Q1: Authors: Please send us the running title Q1]]

[[Q2: It is recommended not to present citations in the abstract. If necessary, adapt the text and include the citation in the body of the text. Q2]]

[[Q3: Authors: Please send us the practical application Q3]]

[[Q4: The in-text citation "Ridriguez et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q4]]

[[Q5: The in-text citation "CONAB (2015)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q5]]

[[Q6: The citation "Bello-Perez et al., 1999" matches the reference

"Bello-Pérez et al, 1999", but an accent or apostrophe is different. Q6]]

[[Q7: The in-text citation "Ridriguez et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q7]]

[[Q8: The in-text citation "Leira et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q8]]

[[Q9: The in-text citation "Oktaviyani (2012)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q9]]

[[Q10: The in-text citation "Singh & Pankaj, 2016" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q10]]

[[Q11: The in-text citation "Greiner & Ralf, 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q11]]

[[Q12: The in-text citation "Venkatesan & Kim, 2010" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q12]]

[[Q13: The in-text citation "Prinaldi et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q13]]

[[Q14: The in-text citation "Houssou & Ayemor (2002" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q14]]

[[Q15: The citation "Lawless & Heymann, 2010" matches multiple references. Please add letters (e.g. "Smith 2000a"), or additional authors to the citation, to uniquely match references and citations. Q15]]

[[Q16: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q16]]

[[Q17: The in-text citation "Association of Official Analytical Chemist, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q17]]

[[Q18: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Q18]]

[[Q19: The in-text citation "Pourhossein & Shalaei (2016)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q19]]

[[Q20: The in-text citation "Rao 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q20]]

[[Q21: The in-text citation "Ahmad et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q21]]

[[Q22: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q22]]

[[Q23: The in-text citation "Rodriguez et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q23]]

[[Q24: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q24]]

[[Q25: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q25]]

[[Q26: The in-text citation "Makame et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q26]]

[[Q27: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q27]]

[[Q28: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q28]]

[[Q29: The in-text citation "BPOM (2011)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q29]]

[[Q30: Reference "Ahmad, Une, Bait, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q30]]

[[Q31: Reference "Association of Official Agricultural Chemists, 2005" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q31]]

[[Q32: Reference "Bustillos-Rodríguez, Tirado-Gallegos, Ordóñez-García, Zamudio-Flores, Ornelas-Paz, Acosta-Muñiz, Gallegos-Morales, Páramo-Calderón, Rios-Velasco, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q32]]

[[Q33: Reference "Greiner, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q33]]

[[Q34: Reference "Houssou, Ayemor, 2012" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q34]]

[[Q35: Send the title of the article Q35]]

[[Q36: Reference "Lawless, Heymann, 2010a" is cited in the text, but the citation matches multiple references. Please check the reference and the citations. Q36]]

[[Q37: Reference "Lawless, Heymann, 2010b" is cited in the text, but the citation matches multiple references. Please check the reference and the citations. Q37]]

[[Q38: Reference "Oktaviyani, 2018" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q38]]

[[Q39: Send the title of the period in full, without abbreviations Q39]]

[[Q40: Send the title of the period in full, without abbreviations Q40]]

[[Q41: Reference "Rao, Radhaiah, Balakrishna, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted. Q41]]

[[Q42: Send the name of all the authors of the publication Q42]]

[[Q43: Send the title of the period in full, without abbreviations Q43]]

[[Q44: Please confirm if the use of the decimal and thousands markers in values of the Table 1-3 is correct (in English dot is used as decimal marker and comma as thousands marker, ex.: 5,931.10, while in Portuguese language comma is used as decimal and dot as thousands marker, ex.: 5.931,10). If necessary make the corrections in the Tables 1-3. Q44]]

[[Q45: Send the explanatory note to the superscript letters present in the table values. Q45]]

[[Q46: Send the explanatory note to the superscript letters present in the table values. Q46]]

[[Q47: Send the explanatory note to the superscript letters present in the table values. Q47]]

[[Q48: Send the explanatory note to the superscript letters present in the table values. Q48]]

Author Confirmation Checklist:

(Please mark with an X and add comments if needed)

Were ALL issues resolved?

Yes: X

No: ___

Author comments:

All of the corrections have been marked using red font.

10.1590/fst.099122

Running **Title**: **Nanomineral-Enhanced Saddah Gohoge Grits** [[Q1: Q1]]

Husain et al.

Original Article

e099122

43

2023

xxx.-xxx.

000

000

0

This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nanomineral fortification on traditional saddah gohoge grits

Rahim HUSAIN^{1*} <https://orcid.org/0000-0001-5307-9806>, Fernandy M. DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using [Association of Official Analytical Chemists Procedure](#) [[Q2: Q2]], and

Comment [AB1]: Authors:
Please send us the running title

Response:
It has been revised

organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application [[Q3: Q3]]:

The practical implication of this article is that the addition of nanomineral fish bones from the family *Lutjanus* sp. to Saddah Gohoge grits can improve its nutritional content without significantly altering its sensory characteristics. This study provides evidence that the use of nanominerals as a fortificant is a promising approach to enhance the nutritional value of traditional foods without compromising their taste and texture. The findings of this study can be useful for food manufacturers and policymakers in developing countries who are interested in improving the nutritional status of their population through the fortification of traditional foods. Additionally, the use of nanominerals can be a more cost-effective and sustainable approach to fortification, as it requires less fortificant material and does not alter the taste and texture of the food.

Received 23 Sept., 2022

Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesia

*Corresponding author: rahim@ung.ac.id

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is "Ugali" or "Ogi", in Italy "Polenta" in Brazil "Angu" and in Romania "Mamaliga". In this study, the traditional Gorontalo grits in Indonesia called "Saddah" or "Gohoge" is made from corn flour. Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Bustillos-Rodríguez, et al., 2019 [[Q4: Q4]]). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). In Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Pérez et al., 1999 [[Q6: Q6]]; Waterschoot et al., 2015; Bustillos-Rodríguez, et al., 2019 [[Q7: Q7]]). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Kumar et al., 2022 [[Q8: Q8]]). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite [Ca₁₀(OH)(PO₄)₆] in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

Comment [AB2]: Authors:
Please send us the practical application

Response:
It has been revised

Comment [AB3]: The in-text citation "Riddríguez et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised

Comment [AB4]: The citation "Bello-Pérez et al., 1999" matches the reference "Bello-Pérez et al., 1999", but an accent or apostrophe is different.

Response:
It has been revised

Comment [AB5]: The in-text citation "Riddríguez et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised

Comment [AB6]: The in-text citation "Lei et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Oktaviyani, 2018)

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016 [[Q10: Q10]]).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner 2009 [[Q11: Q11]]; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah Gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family *Lutjanus* sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchool solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominane) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010 [[Q12: Q12]]). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder modification (Harmain et al., 2018). The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018 [[Q13: Q13]]). Nanocalcium using the *precipitation* method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2012) and Om et al., (2019) modification (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah Gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic quality organoleptic assessment score sheet, the panelists used were

Comment [AB7]: The in-text citation "Singh & Pankaj, 2016" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB8]: The in-text citation "Greiner & Ralf, 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB9]: The in-text citation "Venkatesan & Kim, 2010" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB10]: The in-text citation "Prinaldi et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010 [[Q15: Q15]]).

Proximate analysis

Proximate analysis of Saddah Gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemist (2006) [[Q16: Q16]] method. The Association of Official Agricultural Chemists (2005) and Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster, fat content using the Weibull method (Association of Official Agricultural Chemists, 2005; Iwansyah et al., 2022), carbohydrate content can be calculated by difference, fiber content can be determined using the method of Association of Official Analytical Chemist (2005) [[Q18: Q18]].

Viscosity measurement

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) $L^* a^* b^*$ and hue coordinates. The L^* coordinates describe where $L = 0$ is black, and $L^* = 100$ is colorless. The coordinates of a^* represent red and green shadows, where $a^* > 0$ indicates red and $a^* < 0$ indicates green. The b^* coordinates represent the blue and yellow tones, where $b^* > 0$ indicates the intensity of the yellow color and $b^* < 0$ indicates the blue color. Hue (h^*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomineral analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. the remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) [[Q19: Q19]] determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao 2009 [[Q20: Q20]]).

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah Gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the score sheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the Lutjanus sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018 [[Q21: Q21]]).

Comment [AB11]: The citation "Lawless & Heymann, 2010" matches multiple references. Please add letters (e.g. "Smith 2000a"), or additional authors to the citation, to uniquely match references and citations.

Response:
It has been revised.

Comment [AB12]: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB13]: The in-text citation "Analytical Chemist (2006)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB14]: The in-text citation "Pourhossein & Shalaei (2016)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB15]: The in-text citation "Rao 2009" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Comment [AB16]: The in-text citation "Ahmad et al., 2018" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. 0% with a rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0% concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) [[Q22: Q22]] which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah Gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) [[Q23: Q23]] that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) [[Q24: Q24]] on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) [[Q25: Q25]] showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019 [[Q26: Q26]]). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are -39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

Comment [AB17]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Comment [AB18]: The in-text citation "Rodriguez et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Comment [AB19]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Comment [AB20]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Comment [AB21]: The in-text citation "Makame et al., 2019" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

The results of the staining study were suspected because Saddah Gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarfus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) [Q27: Q27] were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

This is also inline with Iwansyah et al. (2022) study that obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant Rowe Luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah Gohoge grits were the same as those of Ahmad et al. (2018) [Q28: Q28], which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinated Rowe Luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah Gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of Rowe Luwa grits.

For fiber content, the concentration of fortified of 15%, 20%, and 25% not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah Gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah Gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Comment [AB22]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Comment [AB23]: The in-text citation "Ahmad et al. (2018)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. No citations without reference will be deleted.

Response:
It has been revised.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. Furthermore, a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family Lutjanus sp. on Saddah Gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Institute for Research and Community Services (LP2M) of Universitas Negeri Gorontalo, Indonesian Endowment Fund for Education (LPDP) of Ministry of Finance, the reviewers of this article and all those who have helped carry out the research.

References

Ahmad, M., Benjakul, S., Prodpran, T., & Maqsood, S. (2018). Characterization of gelatin from goat skin and its application in fish mousse. *International Journal of Biological Macromolecules*, 108, 188-194.

<jrn>Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.</jrn>

<jrn>Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.</jrn>

Association of Official Analytical Chemists (AOAC). (2005). *Official methods of analysis* (18th ed.). AOAC International. Method 979.21.

<jrn>Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.</jrn>

<jrn>Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.</jrn>

<jrn>[[Q32: Q32]]Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñiz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.</jrn>

<jrn>Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.</jrn>

Comment [AB24]: Reference "Bustillos-Rodríguez, Tirado-Gallegos, Ordóñez-García, Zamudio-Flores, Ornelas-Paz, Acosta-Muñiz, Gallegos-Morales, Páramo-Calderón, Rios-Velasco, 2019" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Response:
It has been revised.

<jrn>Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.</jrn>

<jrn>[[Q33: Q33]]Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpainei.com.br/files/revistas_publicacoes/228.pdf</jrn>

<jrn>Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.</jrn>

<jrn>Harmain, R. M., Dalí, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.</jrn>

<jrn>[[Q34: Q34]]Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.</jrn>

<gov>Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communitie Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335. PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019></gov>

<jrn>Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.</jrn>

<jrn>Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. I., & Luthfiyanti, R. (2022). Effect of pre-gelatinized temperature on physical and nutritional content of Indonesian instant cassava leaves porridge: Rowe Luwa. *Brazilian Journal of Food Technology*, 25, e2021050.</jrn>

<jrn>Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.</jrn>

<jrn>Kelte, I., Fo, Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.</jrn>

Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2022). Utilization of Fish Processing Waste: A Waste to Wealth Approach. In Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (Eds.), *Emerging Post-Harvest Engineering and Technological Interventions for Enhancing Farmer's Income Modified* pp.127-13. ICAR-CIPHET, Ludhiana.

<bok>[[Q37: Q37]]Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5></bok>

Comment [AB25]: Reference "Greiner, 2009" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Response:
It has been revised.

Comment [AB26]: Reference "Houssou, Ayemor, 2012" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Response:
It has been revised.

Comment [AB27]: Reference "Lawless, Heymann, 2010b" is cited in the text, but the citation matches multiple references. Please check the reference and the citations.

Response:
It has been revised.

<jrn>Lekahena, V., Faridah, D. N., Syarief, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.</jrn>

<jrn>Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch–water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).</jrn>

Makame, F. A., Asnani, A., Agustian, E., & Rohman, A. (2019). The effect of white rice flour addition on viscosity of instant grits. *Journal of Physics: Conference Series*, 1236(1), 012032.

<jrn>Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. [PMid:20646299](https://pubmed.ncbi.nlm.nih.gov/20646299/).</jrn>

<jrn>Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.</jrn>

<jrn>[[Q38: Q38]]Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *Jurnal Oseana* [[Q39: Q39]], 40(3), 29-39.</jrn>

<jrn>Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.</jrn>

<jrn>Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *Jurnal Pangan* [[Q40: Q40]], 23(6), 146-156.</jrn>

Pourhossein, M., & Shalaei, H. (2016). Determination of iron in different types of wheat flour consumed in Iran. *Journal of Agricultural Science and Technology*, 18(3), 573-582.

Prinaldi, R., Nurwantoro, N., Sari, M. A., & Muryanto, S. (2018). Hydroxyapatite synthesis from fishbone waste via calcination method. *IOP Conference Series: Materials Science and Engineering*, 308(1), 012024.

Rao, C. R. (2009). *Linear statistical inference and its applications* (2nd ed.). John Wiley & Sons.

<jrn>Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.</jrn>

Rodriguez, J. P., Anunciação, P. C., Garcia, E. E., & Franco, C. M. (2018). Structural and physicochemical properties of corn starches and their function as polysaccharides in food systems: A review. *Food Research International*, 103, 391-401.

Comment [AB28]: Reference "Oktaviyani 2018" is not cited in the text. Please add an in-text citation or the reference will be deleted.

Response:
It has been revised.

Comment [AB29]: Send the title of the period in full, without abbreviations

Response:
It has been revised.

Comment [AB30]: Send the title of the period in full, without abbreviations

Response:
It has been revised.

<jrn>Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.</jrn>

Singh, A. V., & Pankaj, P. (2016). Nanoparticle-mediated delivery of micronutrients in plants. In *Nanotechnology and Plant Sciences* (pp. 265-291). Springer, New Delhi.

<eref>Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download></eref>

<jrn>Suptijah, P., Jacobs, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *Jurnal Akuatika* [[Q43: Q43]], 3(1), 63-73.</jrn>

<jrn>Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. PMID:23822705.</jrn>

Venkatesan, J., & Kim, S. K. (2010). Nano-hydroxyapatite composite biomaterials for bone tissue engineering—a review. *Journal of Biomedical Nanotechnology*, 6(6), 1-12.

<jrn>Wadchararat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.</jrn>

<jrn>Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.</jrn>

<jrn>Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. <http://dx.doi.org/10.3923/ajft.2017.178.185>.</jrn>

<jrn>Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.58320>.</jrn>

<jrn>Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.54322>.</jrn>

Table 1. Hedonic quality organoleptic test results [[Q44: Q44]].

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1.041 ^a	3.40 ± 0.913 ^a	3.08 ± 0.909 ^a	3.64 ± 1.075 ^a	3.16 ± 1.248 ^a
F1 (90:10)	3.68 ± 0.945 ^a	3.28 ± 1.021 ^a	2.88 ± 0.881 ^a	4.00 ± 0.866 ^a	2.96 ± 1.207 ^a
F2 (95:5)	3.44 ± 1.044 ^a	3.36 ± 1.551 ^a	3.00 ± 1.000 ^a	3.96 ± 1.060 ^a	2.56 ± 0.917 ^{ab}
F3 (80:20)	3.32 ± 1.145 ^a	3.48 ± 1.503 ^a [[Q45: Q45]]	2.76 ± 1.052 ^a	3.96 ± 4.00 ^a	2.24 ± 1.200 ^b

a: no significant difference between the treatments

Comment [AB31]: Send the title of the period in full, without abbreviations

Response:
It has been revised.

Comment [AB32]: Please confirm if the use of the decimal and thousands marker in values of the Table 1-3 is correct (in English dot is used as decimal marker and comma as thousands marker, ex.: 5,931.10, while in Portuguese language comma is used as decimal and dot as thousands marker, ex.: 5.931,10). If necessary make the corrections in the Tables 1-3.

Response:
It has been revised.

Comment [AB33]: Send the explanatory note to the superscript letters present in the table values.

Response:
It has been revised.

b, ab: there is significant difference between treatments

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.96 ± 0.841 ^a [[Q46: Q46]]	3.64 ± 1.114 ^a	4.04 ± 0.935 ^a	3.80 ± 1.041 ^a	3.52 ± 1.194 ^a
F1 (90: 10)	3.64 ± 0.995 ^a	3.36 ± 1.319 ^a	3.96 ± 0.889 ^a	3.64 ± 0.907 ^a	3.28 ± 1.100 ^{ab}
F2 (95:15)	2.96 ± 0.790 ^b	3.12 ± 0.927 ^a	3.76 ± 0.879 ^a	2.96 ± 0.935 ^b	2.80 ± 0.913 ^{bc}
F3 (80:20)	2.80 ± 1.000 ^b	2.84 ± 1.143 ^a	3.40 ± 1.118 ^a	2.88 ± 1.013 ^b	2.56 ± 1.044 ^c

a: no significant difference between the treatments
b, c, ab, bc: there is significant difference between treatments

Comment [AB34]: Send the explanatory note to the superscript letters present in the table values.

Response:
It has been revised.

Table 3. Viscosity levels of Saddah *gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
	F1	F2	F3
Viscosity levels (cP)	8309.4667 ± 42.46285 ^a [[Q47: Q47]]	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

a: no significant difference between the treatments
b, c, ab, bc: there is significant difference between treatments

Comment [AB35]: Send the explanatory note to the superscript letters present in the table values.

Response:
It has been revised.

Table 4. Color of Saddah *Gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2.60 ^a [[Q48: Q48]]	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a
*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

a: no significant difference between the treatments

Comment [AB36]: Send the explanatory note to the superscript letters present in the table values.

Response:
It has been revised.

Table 5. Proximate levels of Saddah *Gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Carbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d

a: no significant difference between the treatments
b, c, d, ab, : there is significant difference between treatments

Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphor (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

Figure 1. Process flow for the production of maize flour.

Figure 2. Serving of Saddah *Gohoge* grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

Artikel Proofreading (Tahap 2)

(16 Maret 2023)



Rahim Husain <rahim@ung.ac.id>

cta | Final proofreading sent again | Article: ctaAO099122_EN #CPub - Editora Cubo#

CPub <no-reply@editoracubo.com.br>
Kepada: rahim@ung.ac.id

16 Maret 2023 pukul 20.44

CPub

This notification is addressed to: <rahim@ung.ac.id>

Food Science and Technology

You have a new message regarding document:

[ctaAO099122_EN] NANOMINERAL FORTIFICATION ON TRADITIONAL SADDAH GOHOGE GRITS

To view the message you need to sign in CPub using the link below:




Access CPub

Regards,
CPub

If you're having trouble clicking the "Access CPub" button, copy and paste the URL below into your web browser: https://pub.editoracubo.com.br/publicacoes/cta/documentos/ctaAO099122_EN/mensagens/erZygO4fzvE1iS2ihPUptgpfYIR7VK0XK3jw3IPM51fUsc5ZVnS8JkvDqZFcTHG

© 2023 CPub. All rights reserved.

Subject	cta Final proofreading sent again Article: ctaAO099122_EN
Sent at	03/16/2023 21:44
Read at	03/17/2023 14:18
Message	<p>Dear Author,</p> <p>we identified that there still are issues to be resolved. For the publication of the article in the Food Science and Technology, please answer all issues following the instructions located in the first page of the .DOCX file attached to this email.</p> <p>INSTRUCTIONS:</p> <ul style="list-style-type: none">• The attached .DOCX file should only be opened on versions of MS Office 2010 or higher. After opening the file and before starting your review, make sure that there are no missing spaces like this or this in the text, tables and references. Failure to follow these guidelines could cause the loss of your work and irreversibly corrupt the file. [1]• We will resume working on your article only after receiving the revised document with all issues properly resolved.• No changes in the text will be possible after this step. Make sure that all changes in the content have been made on the revised .DOCX file that you will return us.• To return your corrections, please attach the revised file in the reply to this email.• In the next production step you will receive a PDF proof, in which only layout aspects of the text, figures, tables etc., should be observed. <p>To ensure fast publication of your article, please reply this message as soon as possible.</p> <p>[1] Microsoft® has reported a problem with version 2007 of MS Word® which causes a random loss of spaces between some words when opening the new .DOCX standard file format: http://support.microsoft.com/kb/2633585</p> <p>Best regards, Desktop Publishing Team</p>
Attachments	<ul style="list-style-type: none">•  ctaAO099122_EN_enviadoAutor2.docx 84.3 KB 03/16/2023 21:44

IMPORTANT

- **BEFORE you begin your review** make sure that there are no missing spaces like this or this in the text, tables and references.
- If you fail to follow this instruction you might lose your work.

Review Guidelines

1. Carefully review and make all final text adjustments directly in this Word document.
2. Resolve all issues listed in the "To Authors" section directly in this Word document.
3. When you are done make sure you answer the **Author Confirmation Checklist** and return this Word document by e-mail.
4. The issues listed in the "To Editors" section will be resolved by the journal.

Important:

- Check if the spelling and the order of presentation of article authors name are correct.
- Do not remove or change text styles (font size and color) and color highlights for they are part of our production workflow.
- Figures (if any) are NOT included in this file for at this moment you are required to only and carefully review the content (text and data) of your article. Figures if present in your article will be reviewed in the PDF galley proof.
- All edits must be made directly in this Word document. Requests sent by e-mail or in separate documents will not be accepted and will cause delay in the publication.
- If you need to save the received file, please do not change its extension, **keep it safe in DOCX format.**

The following changes are not allowed and will be taken to the Editor-in-Chief's attention:

- Adding, removing or reordering the Author's list.
- Significantly changing large amounts of text or in tables data.

Issues

To Authors:

[[Q1: Authors: please confirm that Saddah Gohoge is correctly capitalized in title and running title Q1]]

[[Q2: Authors: please confirm that Saddah Gohoge is correctly capitalized in title and running title Q2]]

[[Q3: Authors: please indicate the full name of the author (no abbreviation) Q3]]

[[Q4: The in-text citation "Association of Official Agricultural Chemists, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q4]]

[[Q5: The in-text citation "Association of Official Agricultural Chemists, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q5]]

[[Q6: Authors: please include those responsible for editing the book Q6]]

Author Confirmation Checklist:

(Please mark with an X and add comments if needed)

Were ALL issues resolved?

Yes: ____

No: ____

Author comments:

10.1590/fst.099122

Nanomaterial-enhanced **Saddah Gohoge** [[Q1: Q1]] grits

Husain et al.

Original Article

e099122

43

2023

xxx-xxx.

000

000

0

This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nanomaterial fortification on traditional **Saddah Gohoge** [[Q2: Q2]] grits

Rahim HUSAIN^{1*} <https://orcid.org/0000-0001-5307-9806>, Fernandy M. [[Q3: Q3]] DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using Association of Official Analytical Chemists Procedure, and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application: The practical implication of this article is that the addition of nanomineral fish bones from the family *Lutjanus* sp. to Saddah Gohoge grits can improve its nutritional content without significantly altering its sensory characteristics. This study provides evidence that the use of nanominerals as a fortificant is a promising approach to enhance the nutritional value of traditional foods without compromising their taste and texture. The findings of this study can be useful for food manufacturers and policymakers in developing countries who are interested in improving the nutritional status of their population through the fortification of traditional foods. Additionally, the use of nanominerals can be a more

Comment [VD1]: Authors: please confirm that Saddah Gohoge is correctly capitalized title and running title

Comment [VD2]: Authors: please confirm that Saddah Gohoge is correctly capitalized title and running title

Comment [VD3]: Authors: please indicate the full name of the author (no abbreviation)

cost-effective and sustainable approach to fortification, as it requires less fortificant material and does not alter the taste and texture of the food.

Received 23 Sept., 2022
Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesia

*Corresponding author: rahim@ung.ac.id

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is "Ugali" or "Ogi", in Italy "Polenta" in Brazil "Angu" and in Romania "Mamaliga". In this study, the traditional Gorontalo grits in Indonesia called "Saddah" or "Gohoge" is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Bustillos-Rodríguez et al., 2019). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). In Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Pérez et al., 1999; Waterschoot et al., 2015; Bustillos-Rodríguez et al., 2019). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Kumar et al., 2022). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite [Ca₁₀(OH)(PO₄)₆] in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Oktaviyani, 2018).

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner, 2009; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out

fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah Gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family *Lutjanus* sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, Luff-Schoorl solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder modification (Harmain et al., 2018). The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018). Nanocalcium using the precipitation method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2012) and Om et al., (2019) modification (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah Gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic quality organoleptic assessment score sheet, the panelists used were untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010).

Proximate analysis

Proximate analysis of Saddah Gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemists (2005) method. The Association of Official Agricultural Chemists (2005) [[Q4: Q4]] and Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster, fat content using the Weibull method (Association of Official Agricultural Chemists, 2005 [[Q5: Q5]]; Iwansyah et al., 2022), carbohydrate content can be calculated by difference, fiber content can be determined using the method of Association of Official Analytical Chemists (2005).

Viscosity measurement

Comment [VD4]: The in-text citation "Association of Official Agricultural Chemists (2005)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Comment [VD5]: The in-text citation "Association of Official Agricultural Chemists (2005)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L^* a^* b^* and hue coordinates. The L^* coordinates describe where $L = 0$ is black, and $L^* = 100$ is colorless. The coordinates of a^* represent red and green shadows, where $a^* > 0$ indicates red and $a^* < 0$ indicates green. The b^* coordinates represent the blue and yellow tones, where $b^* > 0$ indicates the intensity of the yellow color and $b^* < 0$ indicates the blue color. Hue (h^*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomaterial analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. The remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme. Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao, 2009).

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah Gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the score sheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the Lutjanus sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018).

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals Lutjanus sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah Gohoge grits fortification of fish bone nanominerals Lutjanus sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the Lutjanus sp. 0% with a rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the Lutjanus sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0%

concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah Gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are - 39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

The results of the staining study were suspected because Saddah Gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarpus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

This is also inline with Iwansyah et al. (2022) study that obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant Rowe Luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah Gohoge grits were the same as those of Ahmad et al. (2018), which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinized Rowe Luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah Gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of Rowe Luwa grits.

For fiber content, the concentration of fortified of 15%, 20%, and 25% not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah Gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah Gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. Furthermore, a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on Saddah Gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Institute for Research and Community Services (LP2M) of Universitas Negeri Gorontalo, Indonesian Endowment Fund for Education (LPDP) of Ministry of Finance, the reviewers of this article and all those who have helped carry out the research.

References

<jrn>Ahmad, M., Benjakul, S., Prodpran, T., & Maqsood, S. (2018). Characterization of gelatin from goat skin and its application in fish mousse. *International Journal of Biological Macromolecules*, 108, 188-194.</jrn>

<jrn>Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.</jrn>

<jrn>Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.</jrn>

<bok>Association of Official Analytical Chemists – AOAC. (2005). *Official methods of analysis: method 979.21* (18th ed.). Washington, DC: AOAC International. </bok>

<jrn>Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.</jrn>

<jrn>Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.</jrn>

<jrn>Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñiz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.</jrn>

<jrn>Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.</jrn>

<jrn>Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.</jrn>

<jrn>Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpanel.com.br/files/revistas_publicacoes/228.pdf.</jrn>

<jrn>Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.</jrn>

<jrn>Harmain, R. M., Dali, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.</jrn>

<jrn>Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.</jrn>

<gov>Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communitie Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335, PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019></gov>

<jrn>Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.</jrn>

<jrn>Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. I., & Luthfiyanti, R. (2022). Effect of pre-gelatinized temperature on physical and nutritional content of indonesian instant cassava leaves porridge: Rowe Luwa. *Brazilian Journal of Food Technology*, 25, e2021050. <http://dx.doi.org/10.1590/1981-6723.05021>.</jrn>

<jrn>Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.</jrn>

<jrn>Kelte, I., Fo, Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.</jrn>

<edb>Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2022). Utilization of fish processing waste: a waste to wealth approach. In V. Kumar, A. U. Muzaddad, S. Mann, R. Balakrishnan, K. Bembem & Y. Kalnar (Eds.), *Emerging post-harvest engineering and technological interventions for enhancing farmer's income modified* (pp.127-133. Ludhiana: ICAR-CIPHET.</edb>

<bok>Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5></bok>

<jrn>Lekahena, V., Faridah, D. N., Syarief, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.</jrn>

<jrn>Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch–water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).</jrn>

<jrn>Makame, F. A., Asnani, A., Agustian, E., & Rohman, A. (2019). The effect of white rice flour addition on viscosity of instant grits. *Journal of Physics: Conference Series*, 1236(1), 012032.</jrn>

<jrn>Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. PMID:20646299.</jrn>

<jrn>Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.</jrn>

<jrn>Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *Jurnal Oseana*, 40(3), 29-39.</jrn>

<jrn>Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.</jrn>

<jrn>Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *Jurnal Pangan*, 23(6), 146-156.</jrn>

<jrn>Pourhossein, M., & Shalaei, H. (2016). Determination of iron in different types of wheat flour consumed in Iran. *Journal of Agricultural Science and Technology*, 18(3), 573-582.</jrn>

<jrn>Prinaldi, R., Nurwantoro, N., Sari, M. A., & Muryanto, S. (2018). Hydroxyapatite synthesis from fishbone waste via calcination method. *IOP Conference Series: Materials Science and Engineering*, 308(1), 012024.</jrn>

<bok>Rao, C. R. (2009). *Linear statistical inference and its applications* (2nd ed.). New York: John Wiley & Sons.</bok>

<jrn>Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.</jrn>

<jrn>Rodriguez, J. P., Anunciação, P. C., Garcia, E. E., & Franco, C. M. (2018). Structural and physicochemical properties of corn starches and their function as polysaccharides in food systems: A review. *Food Research International*, 103, 391-401.</jrn>

<edb>Singh, A. V., & Pankaj, P. (2016). Nanoparticle-mediated delivery of micronutrients in plants. In A. Surname [[Q6: Q6]] (Ed.), *Nanotechnology and plant sciences* (pp. 265-291). New Delhi: Springer.</edb>

Comment [VD6]: Authors: please include those responsible for editing the book

<jrn>Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.</jrn>

<eref>Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download></eref>

<jrn>Suptijah, P., Jacob, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *Jurnal Akuatika*, 3(1), 63-73.</jrn>

<jrn>Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. [PMid:23822705](https://pubmed.ncbi.nlm.nih.gov/23822705/).</jrn>

<jrn>Venkatesan, J., & Kim, S. K. (2010). Nano-hydroxyapatite composite biomaterials for bone tissue engineering—a review. *Journal of Biomedical Nanotechnology*, 6(6), 1-12. [PMid:25992432](https://pubmed.ncbi.nlm.nih.gov/25992432/).</jrn>

<jrn>Wadcharat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.</jrn>

<jrn>Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.</jrn>

<jrn>Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. <http://dx.doi.org/10.3923/ajft.2017.178.185>.</jrn>

<jrn>Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.58320>.</jrn>

<jrn>Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.54322>.</jrn>

Table 1. Hedonic quality organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1.041 ^a	3.40 ± 0.913 ^a	3.08 ± 0.909 ^a	3.64 ± 1.075 ^a	3.16 ± 1.248 ^a
F1 (90:10)	3.68 ± 0.945 ^a	3.28 ± 1.021 ^a	2.88 ± 0.881 ^a	4.00 ± 0.866 ^a	2.96 ± 1.207 ^a
F2 (95:15)	3.44 ± 1.044 ^a	3.36 ± 1.551 ^a	3.00 ± 1.000 ^a	3.96 ± 1.060 ^a	2.56 ± 0.917 ^{ab}
F3 (80:20)	3.32 ± 1.145 ^a	3.48 ± 1.503 ^a	2.76 ± 1.052 ^a	3.96 ± 4.00 ^a	2.24 ± 1.200 ^b

a: no significant difference between the treatments.

b, ab: there is significant difference between treatments.

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.96 ± 0.841 ^a	3.64 ± 1.114 ^a	4.04 ± 0.935 ^a	3.80 ± 1.041 ^a	3.52 ± 1.194 ^a
F1 (90: 10)	3.64 ± 0.995 ^a	3.36 ± 1.319 ^a	3.96 ± 0.889 ^a	3.64 ± 0.907 ^a	3.28 ± 1.100 ^{ab}
F2 (95:15)	2.96 ± 0.790 ^b	3.12 ± 0.927 ^a	3.76 ± 0.879 ^a	2.96 ± 0.935 ^b	2.80 ± 0.913 ^{bc}
F3 (80:20)	2.80 ± 1.000 ^b	2.84 ± 1.143 ^a	3.40 ± 1.118 ^a	2.88 ± 1.013 ^b	2.56 ± 1.044 ^c

a: no significant difference between the treatments.
b, c, ab, bc: there is significant difference between treatments.

Table 3. Viscosity levels of Saddah *gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
Viscosity levels (cP)	F1	F2	F3
	8309.4667 ± 42.46285 ^a	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

a: no significant difference between the treatments.
b, c: there is significant difference between treatments.

Table 4. Color of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2.60 ^a	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a
*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

a: no significant difference between the treatments.

Table 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Carbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d

a: no significant difference between the treatments.
b, c, d, ab: there is significant difference between treatments.

Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphor (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

Figure 1. Process flow for the production of maize flour.

Figure 2. Serving of Saddah Gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

**Pengiriman Revisi
Proofreading (Tahap 2)
(27 Maret 2023)**

Autor	RE: cta Final proofreading sent again Article: ctaAO099122_EN  Read at 03/27/2023 20:46	03/27/2023 09:33 ☰ Prova de Conteúdo
Subject	RE: cta Final proofreading sent again Article: ctaAO099122_EN	
Sent at	03/27/2023 09:33	
Read at	03/27/2023 20:46	
Message	<p>Dear Editorial Team,</p> <p>The revised document is attached in this email. Is it possible for the article to be published in this month?</p> <p>Thank you and I look forward to hearing from you.</p> <p>Best regards,</p> <p>Rahim Husain</p>	
Attachments	•  ctaAO099122_EN_enviadoAutor2.docx 129.0 KB 03/27/2023 09:33	

 reply

IMPORTANT

- **BEFORE you begin your review** make sure that there are no missing spaces like this or this in the text, tables and references.
- If you fail to follow this instruction you might lose your work.

Review Guidelines

1. Carefully review and make all final text adjustments directly in this Word document.
2. Resolve all issues listed in the "To Authors" section directly in this Word document.
3. When you are done make sure you answer the **Author Confirmation Checklist** and return this Word document by e-mail.
4. The issues listed in the "To Editors" section will be resolved by the journal.

Important:

- Check if the spelling and the order of presentation of article authors name are correct.
- Do not remove or change text styles (font size and color) and color highlights for they are part of our production workflow.
- Figures (if any) are NOT included in this file for at this moment you are required to only and carefully review the content (text and data) of your article. Figures if present in your article will be reviewed in the PDF galley proof.
- All edits must be made directly in this Word document. Requests sent by e-mail or in separate documents will not be accepted and will cause delay in the publication.
- If you need to save the received file, please do not change its extension, **keep it safe in DOCX format.**

The following changes are not allowed and will be taken to the Editor-in-Chief's attention:

- Adding, removing or reordering the Author's list.
- Significantly changing large amounts of text or in tables data.

Issues

To Authors:

[[Q1: Authors: please confirm that Saddah Gohoge is correctly capitalized in title and running title Q1]]

[[Q2: Authors: please confirm that Saddah Gohoge is correctly capitalized in title and running title Q2]]

[[Q3: Authors: please indicate the full name of the author (no abbreviation) Q3]]

[[Q4: The in-text citation "Association of Official Agricultural Chemists, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q4]]

[[Q5: The in-text citation "Association of Official Agricultural Chemists, 2005" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted. Q5]]

[[Q6: Authors: please include those responsible for editing the book Q6]]

Author Confirmation Checklist:

(Please mark with an X and add comments if needed)

Were ALL issues resolved?

Yes: X

No: ____

Author comments:

All changes have been marked with red font.

10.1590/fst.099122

Nanomineral-enhanced **Saddah Gohoge** **[[Q1: Q1]] grits**

Husain et al.

Original Article

e099122

43

2023

xxx.-xxx.

000

000

0

This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nanomineral fortification on traditional **Saddah Gohoge** **[[Q2: Q2]] grits**

Rahim HUSAIN¹* <https://orcid.org/0000-0001-5307-9806>, Fernandy **[[Q3: Q3]]** DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using Association of Official Analytical Chemists Procedure, and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application: The practical implication of this article is that the addition of nanomineral fish bones from the family *Lutjanus* sp. to Saddah Gohoge grits can improve its nutritional content without significantly altering its sensory characteristics. This study provides evidence that the use of nanominerals as a fortificant is a promising approach to enhance the nutritional value of traditional foods without compromising their taste and texture. The findings of this study can be useful for food manufacturers and policymakers in developing countries who are interested in improving the nutritional status of their population through the fortification of traditional foods. Additionally, the use of nanominerals can be a more

Comment [VD1]: Authors: please confirm that Saddah Gohoge is correctly capitalized title and running title

Response:
Yes

Comment [VD2]: Authors: please confirm that Saddah Gohoge is correctly capitalized title and running title

Response:
Yes

cost-effective and sustainable approach to fortification, as it requires less fortificant material and does not alter the taste and texture of the food.

Received 23 Sept., 2022

Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesia

*Corresponding author: rahim@ung.ac.id

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is "Ugali" or "Ogi", in Italy "Polenta" in Brazil "Angu" and in Romania "Mamaliga". In this study, the traditional Gorontalo grits in Indonesia called "Saddah" or "Gohoge" is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Bustillos-Rodríguez et al., 2019). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). In Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Pérez et al., 1999; Waterschoot et al., 2015; Bustillos-Rodríguez et al., 2019). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Kumar et al., 2022). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite [Ca₁₀(OH)(PO₄)₆] in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Oktaviyani, 2018).

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner, 2009; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out

fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah Gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family *Lutjanus* sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchoorl solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder modification (Harmain et al., 2018). The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018). Nanocalcium using the precipitation method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2012) and Om et al., (2019) modification (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah Gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic quality organoleptic assessment score sheet, the panelists used were untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010).

Proximate analysis

Proximate analysis of Saddah Gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemists (2005) method. The Association of Official Analytical Chemists (2005) [[Q4: Q4]] and Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster, fat content using the Weibull method (Association of Official Analytical Chemists [2005] [[Q5: Q5]]; Iwansyah et al., 2022), carbohydrate content can be calculated by difference, fiber content can be determined using the method of Association of Official Analytical Chemists (2005).

Viscosity measurement

Comment [VD3]: The in-text citation "Association of Official Agricultural Chemists (2005)" is not in the reference list. Please correct the citation, add the reference to the list, or delete the citation. Note: citations without reference will be deleted.

Response:
It has been revised.

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L^* a^* b^* and hue coordinates. The L^* coordinates describe where $L = 0$ is black, and $L^* = 100$ is colorless. The coordinates of a^* represent red and green shadows, where $a^* > 0$ indicates red and $a^* < 0$ indicates green. The b^* coordinates represent the blue and yellow tones, where $b^* > 0$ indicates the intensity of the yellow color and $b^* < 0$ indicates the blue color. Hue (h^*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomineral analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. the remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme. Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao, 2009).

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah Gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the score sheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the Lutjanus sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018).

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals Lutjanus sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah Gohoge grits fortification of fish bone nanominerals Lutjanus sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the Lutjanus sp. 0% with a rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the Lutjanus sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0%

concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah Gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are - 39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

The results of the staining study were suspected because Saddah Gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarpus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

This is also inline with Iwansyah et al. (2022) study that obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant Rowe Luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah Gohoge grits were the same as those of Ahmad et al. (2018), which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinized Rowe Luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah Gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of Rowe Luwa grits.

For fiber content, the concentration of fortified of 15%, 20%, and 25% not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah Gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which is 30.89% and P 15.22%. Saddah Gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. Furthermore, a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on Saddah Gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Institute for Research and Community Services (LP2M) of Universitas Negeri Gorontalo, Indonesian Endowment Fund for Education (LPDP) of Ministry of Finance, the reviewers of this article and all those who have helped carry out the research.

References

<jrn>Ahmad, M., Benjakul, S., Prodpran, T., & Maqsood, S. (2018). Characterization of gelatin from goat skin and its application in fish mousse. *International Journal of Biological Macromolecules*, 108, 188-194.</jrn>

<jrn>Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.</jrn>

<jrn>Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.</jrn>

<bok>Association of Official Analytical Chemists – AOAC. (2005). *Official methods of analysis: method 979.21* (18th ed.). Washington, DC: AOAC International. </bok>

<jrn>Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.</jrn>

<jrn>Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.</jrn>

<jrn>Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñiz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.</jrn>

<jrn>Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.</jrn>

<jrn>Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.</jrn>

<jrn>Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpanel.com.br/files/revistas_publicacoes/228.pdf.</jrn>

<jrn>Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.</jrn>

<jrn>Harmain, R. M., Dali, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.</jrn>

<jrn>Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.</jrn>

<gov>Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communitie Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335, PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019></gov>

<jrn>Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.</jrn>

<jrn>Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. I., & Luthfiyanti, R. (2022). Effect of pre-gelatinized temperature on physical and nutritional content of indonesian instant cassava leaves porridge: Rowe Luwa. *Brazilian Journal of Food Technology*, 25, e2021050. <http://dx.doi.org/10.1590/1981-6723.05021>.</jrn>

<jrn>Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.</jrn>

<jrn>Kelte, I., Fo, Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.</jrn>

<edb>Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2022). Utilization of fish processing waste: a waste to wealth approach. In V. Kumar, A. U. Muzaddad, S. Mann, R. Balakrishnan, K. Bembem & Y. Kalnar (Eds.), *Emerging post-harvest engineering and technological interventions for enhancing farmer's income modified* (pp.127-133. Ludhiana: ICAR-CIPHET.</edb>

<bok>Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5></bok>

<jrn>Lekahena, V., Faridah, D. N., Syarief, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.</jrn>

<jrn>Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch–water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).</jrn>

<jrn>Makame, F. A., Asnani, A., Agustian, E., & Rohman, A. (2019). The effect of white rice flour addition on viscosity of instant grits. *Journal of Physics: Conference Series*, 1236(1), 012032.</jrn>

<jrn>Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. PMID:20646299.</jrn>

<jrn>Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.</jrn>

<jrn>Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *Jurnal Oseana*, 40(3), 29-39.</jrn>

<jrn>Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.</jrn>

<jrn>Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *Jurnal Pangan*, 23(6), 146-156.</jrn>

<jrn>Pourhossein, M., & Shalaei, H. (2016). Determination of iron in different types of wheat flour consumed in Iran. *Journal of Agricultural Science and Technology*, 18(3), 573-582.</jrn>

<jrn>Prinaldi, R., Nurwantoro, N., Sari, M. A., & Muryanto, S. (2018). Hydroxyapatite synthesis from fishbone waste via calcination method. *IOP Conference Series: Materials Science and Engineering*, 308(1), 012024.</jrn>

<bok>Rao, C. R. (2009). *Linear statistical inference and its applications* (2nd ed.). New York: John Wiley & Sons.</bok>

<jrn>Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.</jrn>

<jrn>Rodriguez, J. P., Anunciação, P. C., Garcia, E. E., & Franco, C. M. (2018). Structural and physicochemical properties of corn starches and their function as polysaccharides in food systems: A review. *Food Research International*, 103, 391-401.</jrn>

<edb>Singh, A. V., & Pankaj, P. (2016). Nanoparticle-mediated delivery of micronutrients in plants. In Chandra, S., & Chauhan, H. K. ([Q6: Q6]) (Ed.), *Nanotechnology and plant sciences* (pp. 265-291). New Delhi: Springer.</edb>

<jrn>Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.</jrn>

<eref>Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download></eref>

<jrn>Suptijah, P., Jacob, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *Jurnal Akuatika*, 3(1), 63-73.</jrn>

<jrn>Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. [PMid:23822705](https://pubmed.ncbi.nlm.nih.gov/23822705/).</jrn>

<jrn>Venkatesan, J., & Kim, S. K. (2010). Nano-hydroxyapatite composite biomaterials for bone tissue engineering—a review. *Journal of Biomedical Nanotechnology*, 6(6), 1-12. [PMid:25992432](https://pubmed.ncbi.nlm.nih.gov/25992432/).</jrn>

<jrn>Wadcharat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.</jrn>

<jrn>Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.</jrn>

<jrn>Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. <http://dx.doi.org/10.3923/ajft.2017.178.185>.</jrn>

<jrn>Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.58320>.</jrn>

<jrn>Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.54322>.</jrn>

Table 1. Hedonic quality organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1.041 ^a	3.40 ± 0.913 ^a	3.08 ± 0.909 ^a	3.64 ± 1.075 ^a	3.16 ± 1.248 ^a
F1 (90:10)	3.68 ± 0.945 ^a	3.28 ± 1.021 ^a	2.88 ± 0.881 ^a	4.00 ± 0.866 ^a	2.96 ± 1.207 ^a
F2 (95:15)	3.44 ± 1.044 ^a	3.36 ± 1.551 ^a	3.00 ± 1.000 ^a	3.96 ± 1.060 ^a	2.56 ± 0.917 ^{ab}
F3 (80:20)	3.32 ± 1.145 ^a	3.48 ± 1.503 ^a	2.76 ± 1.052 ^a	3.96 ± 4.00 ^a	2.24 ± 1.200 ^b

a: no significant difference between the treatments.

b, ab: there is significant difference between treatments.

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.96 ± 0.841 ^a	3.64 ± 1.114 ^a	4.04 ± 0.935 ^a	3.80 ± 1.041 ^a	3.52 ± 1.194 ^a
F1 (90: 10)	3.64 ± 0.995 ^a	3.36 ± 1.319 ^a	3.96 ± 0.889 ^a	3.64 ± 0.907 ^a	3.28 ± 1.100 ^{ab}
F2 (95:15)	2.96 ± 0.790 ^b	3.12 ± 0.927 ^a	3.76 ± 0.879 ^a	2.96 ± 0.935 ^b	2.80 ± 0.913 ^{bc}
F3 (80:20)	2.80 ± 1.000 ^b	2.84 ± 1.143 ^a	3.40 ± 1.118 ^a	2.88 ± 1.013 ^b	2.56 ± 1.044 ^c

a: no significant difference between the treatments.
b, c, ab, bc: there is significant difference between treatments.

Table 3. Viscosity levels of Saddah *gohoge* grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
Viscosity levels (cP)	F1	F2	F3
	8309.4667 ± 42.46285 ^a	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

a: no significant difference between the treatments.
b, c: there is significant difference between treatments.

Table 4. Color of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2.60 ^a	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a
*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

a: no significant difference between the treatments.

Table 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Carbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d

a: no significant difference between the treatments.
b, c, d, ab: there is significant difference between treatments.

Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphor (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

Figure 1. Process flow for the production of maize flour.

Figure 2. Serving of Saddah Gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

Final proofreading completed
(27 Maret 2023)



Rahim Husain <rahim@ung.ac.id>

cta | Final proofreading completed | Article: ctaAO099122_EN #CPub - Editora Cubo#

EC Marcação <edit@editoracubo.com.br>
Balas Ke: EC Marcação <edit@editoracubo.com.br>
Kepada: rahim@ung.ac.id

27 Maret 2023 pukul 19.56

CPub

Dear Author,

We contact you to inform that the final proofreading was completed and the article followed to the next stage of production.

Thank you for your cooperation.

Best regards,
Desktop Publishing Team

© 2023 CPub. Todos os direitos reservados.

Layout Proof

(30 Maret 2023)



Rahim Husain <rahim@ung.ac.id>

cta | Layout proof | Article: ctaAO099122_EN #CPub - Editora Cubo#

CPub <no-reply@editoracubo.com.br>
Kepada: rahim@ung.ac.id

30 Maret 2023 pukul 16.55

CPub

This notification is addressed to: <rahim@ung.ac.id>

Food Science and Technology

You have a new message regarding document:

[ctaAO099122_EN] NANOMINERAL FORTIFICATION ON TRADITIONAL
SADDAH GOHOGE GRITS

To view the message you need to sign in CPub using the link below:



Access CPub

Regards,
CPub

If you're having trouble clicking the "Access CPub" button, copy and paste the URL below into your web browser: https://pub.editoracubo.com.br/publicacoes/cta/documentos/ctaAO099122_EN/mensagens/erZygO4fzvE1iS2ihPUptgpfYIR7VK0XK3jw3IPM51fUsc5ZVnS8JkvDqZFcTHG

© 2023 CPub. All rights reserved.



Nanomineral fortification on traditional Saddah Gohoge grits

Rahim HUSAIN^{1*} , Fernandy DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using Association of Official Analytical Chemists Procedure, and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application: The practical implication of this article is that the addition of nanomineral fish bones from the family *Lutjanus* sp. to Saddah Gohoge grits can improve its nutritional content without significantly altering its sensory characteristics. This study provides evidence that the use of nanominerals as a fortificant is a promising approach to enhance the nutritional value of traditional foods without compromising their taste and texture. The findings of this study can be useful for food manufacturers and policymakers in developing countries who are interested in improving the nutritional status of their population through the fortification of traditional foods. Additionally, the use of nanominerals can be a more cost-effective and sustainable approach to fortification, as it requires less fortificant material and does not alter the taste and texture of the food.

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is “Ugali” or “Ogi”, in Italy “Polenta” in Brazil “Angu” and in Romania “Mamaliga”. In this study, the traditional Gorontalo grits in Indonesia called “Saddah” or “Gohoge” is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Bustillos-Rodríguez et al., 2019). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). In Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear

and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Pérez et al., 1999; Waterschoot et al., 2015; Bustillos-Rodríguez et al., 2019). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is

Received 23 Sept., 2022

Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesia

*Corresponding author: rahim@ung.ac.id

divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Kumar et al., 2022). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite $[Ca_{10}(OH)(PO_4)_6]$ in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Oktaviyani, 2018).

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner, 2009; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah Gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family

Lutjanus sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchool solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder modification (Harmain et al., 2018). The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018). Nanocalcium using the *precipitation* method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2012) and Om et al., (2019) modification (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah Gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic




Figure 1. Process flow for the production of maize flour.

quality organoleptic assessment score sheet, the panelists used were untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010).

Proximate analysis

Proximate analysis of Saddah Gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemists (2005) method. The Association of Official Analytical Chemists (2005) and Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster, fat content using the Weibull method (Association of Official Analytical Chemists (2005; Iwansyah et al., 2022), carbohydrate content can be calculated by difference, fiber content can be determined using the method of Association of Official Analytical Chemists (2005).

Viscosity measurement

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an

aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L^* a^* b^* and hue coordinates. The L^* coordinates describe where $L = 0$ is black, and $L^* = 100$ is colorless. The coordinates of a^* represent red and green shadows, where $a^* > 0$ indicates red and $a^* < 0$ indicates green. The b^* coordinates represent the blue and yellow tones, where $b^* > 0$ indicates the intensity of the yellow color and $b^* < 0$ indicates the blue color. Hue (h^*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomineral analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. the remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao, 2009).

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah Gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the score sheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the *Lutjanus* sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018).

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn

as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. 0% with a




Figure 2. Serving of Saddah Gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

Table 1. Hedonic quality organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1.041 ^a	3.40 ± 0.913 ^a	3.08 ± 0.909 ^a	3.64 ± 1.075 ^a	3.16 ± 1.248 ^a
F1 (90: 10)	3.68 ± 0.945 ^a	3.28 ± 1.021 ^a	2.88 ± 0.881 ^a	4.00 ± 0.866 ^a	2.96 ± 1.207 ^a
F2 (95:15)	3.44 ± 1.044 ^a	3.36 ± 1.551 ^a	3.00 ± 1.000 ^a	3.96 ± 1.060 ^a	2.56 ± 0.917 ^{ab}
F3 (80:20)	3.32 ± 1.145 ^a	3.48 ± 1.503 ^a	2.76 ± 1.052 ^a	3.96 ± 4.00 ^a	2.24 ± 1.200 ^b

a: no significant difference between the treatments. b, ab: there is significant difference between treatments.

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.96 ± 0.841 ^a	3.64 ± 1.114 ^a	4.04 ± 0.935 ^a	3.80 ± 1.041 ^a	3.52 ± 1.194 ^a
F1 (90: 10)	3.64 ± 0.995 ^a	3.36 ± 1.319 ^a	3.96 ± 0.889 ^a	3.64 ± 0.907 ^a	3.28 ± 1.100 ^{ab}
F2 (95:15)	2.96 ± 0.790 ^b	3.12 ± 0.927 ^a	3.76 ± 0.879 ^a	2.96 ± 0.935 ^b	2.80 ± 0.913 ^{bc}
F3 (80:20)	2.80 ± 1.000 ^b	2.84 ± 1.143 ^a	3.40 ± 1.118 ^a	2.88 ± 1.013 ^b	2.56 ± 1.044 ^c

a: no significant difference between the treatments. b, c, ab, bc: there is significant difference between treatments.

rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0% concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah Gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled

water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are -39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

The results of the staining study were suspected because Saddah Gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarfus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

Table 3. Viscosity levels of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
	F1	F2	F3
Viscosity levels (cP)	8309.4667 ± 42.46285 ^a	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

a: no significant difference between the treatments. b, c: there is significant difference between treatments.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

This is also inline with Iwansyah et al. (2022) study that obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant Rowe Luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification

of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah Gohoge grits were the same as those of Ahmad et al. (2018), which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinized Rowe Luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah Gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of Rowe Luwa grits.

For fiber content, the concentration of fortified of 15%, 20%, and 25% not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah Gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which

Table 4. Color of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2.60 ^a	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a
*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

a: no significant difference between the treatments.

Table 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Carbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d

a: no significant difference between the treatments. b, c, d, ab: there is significant difference between treatments.

Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphorus (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

is 30.89% and P 15.22%. Saddah Gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. Furthermore, a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on Saddah Gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Institute for Research and Community Services (LP2M) of Universitas Negeri Gorontalo, Indonesian Endowment Fund for Education (LPDP) of Ministry of Finance, the reviewers of this article and all those who have helped carry out the research.

References

Ahmad, M., Benjakul, S., Prodpran, T., & Maqsood, S. (2018). Characterization of gelatin from goat skin and its application in fish mousse. *International Journal of Biological Macromolecules*, 108, 188-194.

Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.

Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.

Association of Official Analytical Chemists – AOAC. (2005). *Official methods of analysis: method 979.21* (18th ed.). Washington, DC: AOAC International.

Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.

Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.

Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñoz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.

Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.

Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.

Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpanel.com.br/files/revistas_publicacoes/228.pdf

Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.

Harmain, R. M., Dali, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.

Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.

Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communit Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335, PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019>

- Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.
- Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. L., & Luthfiyanti, R. (2022). Effect of pre-gelatinized temperature on physical and nutritional content of Indonesian instant cassava leaves porridge: Rowe Luwa. *Brazilian Journal of Food Technology*, 25, e2021050. <http://dx.doi.org/10.1590/1981-6723.05021>.
- Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.
- Kelte, I., Fo., Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.
- Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2022). Utilization of fish processing waste: a waste to wealth approach. In V. Kumar, A. U. Muzaddadi, S. Mann, R. Balakrishnan, K. Bembem & Y. Kalnar (Eds.), *Emerging post-harvest engineering and technological interventions for enhancing farmer's income modified* (pp.127-13. Ludhiana: ICAR-CIPHET.
- Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5>
- Lekahena, V., Faridah, D. N., Syarief, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.
- Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch-water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).
- Makame, F. A., Asnani, A., Agustian, E., & Rohman, A. (2019). The effect of white rice flour addition on viscosity of instant grits. *Journal of Physics: Conference Series*, 1236(1), 012032.
- Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. PMID:20646299.
- Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.
- Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *Jurnal Oseana*, 40(3), 29-39.
- Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.
- Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *Jurnal Pangan*, 23(6), 146-156.
- Pourhossein, M., & Shalaei, H. (2016). Determination of iron in different types of wheat flour consumed in Iran. *Journal of Agricultural Science and Technology*, 18(3), 573-582.
- Prinaldi, R., Nurwantoro, N., Sari, M. A., & Muryanto, S. (2018). Hydroxyapatite synthesis from fishbone waste via calcination method. *IOP Conference Series. Materials Science and Engineering*, 308(1), 012024.
- Rao, C. R. (2009). *Linear statistical inference and its applications* (2nd ed.). New York: John Wiley & Sons.
- Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.
- Rodriguez, J. P., Anunciação, P. C., Garcia, E. E., & Franco, C. M. (2018). Structural and physicochemical properties of corn starches and their function as polysaccharides in food systems: a review. *Food Research International*, 103, 391-401.
- Singh, A. V., & Pankaj, P. (2016). Nanoparticle-mediated delivery of micronutrients in plants. In S. Chandra & H. K. Chauhan (Eds.), *Nanotechnology and plant sciences* (pp.265-291). New Delhi: Springer.
- Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.
- Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download>
- Suptijah, P., Jacob, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *Jurnal Akuatika*, 3(1), 63-73.
- Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. PMID:23822705.
- Venkatesan, J., & Kim, S. K. (2010). Nano-hydroxyapatite composite biomaterials for bone tissue engineering—a review. *Journal of Biomedical Nanotechnology*, 6(6), 1-12. PMID:25992432.
- Wadchararat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.
- Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.
- Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. <http://dx.doi.org/10.3923/ajft.2017.178.185>.
- Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.58320>.
- Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.54322>.

**Approval Layout Proof oleh
penulis**

(1 Maret 2023)

Autor	RE: cta Layout proof Article: ctaAO099122_EN Read at 04/07/2023 07:42	04/01/2023 21:45 Prova Autor
Subject	RE: cta Layout proof Article: ctaAO099122_EN	
Sent at	04/01/2023 21:45	
Read at	04/07/2023 07:42	
Message	APPROVED IN CURRENT VERSION	
reply		

**Artikel terbit April 2023,
Volume 43**



Nanomineral fortification on traditional Saddah Gohoge grits

Rahim HUSAIN^{1*} , Fernandy DJAILANI¹, Rita Marsuci HARMAIN¹

Abstract

Saddah Gohoge grits is a traditional food in Gorontalo Province, Indonesia, especially the elderly and when children are growing. The study aimed to obtain the physical, chemical, and organoleptic characteristics of the Saddah Gohoge grits fortification of nanomineral fish bones from the family *Lutjanus* sp. The materials used were Saddah Gohoge corn, bone nanominerals of the family *Lutjanus* sp. The nanomineral using *precipitation* method. The proximate, viscosity, and colour using Association of Official Analytical Chemists Procedure, and organoleptic using non-parametric. Data analysis using analysis of variance, descriptive and *kruskal wallis*. Proximate levels increased along with the increase in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. in Saddah Gohoge grits, but the water content decreased due to the use of the same volume of water and the composition of Saddah Gohoge. Hedonic quality organoleptic appearance is not dull, the texture is slightly smooth and slightly thick, the color is slightly yellowish, the flavour is slightly pungent, flavour of corn, the taste is corn. The results of the hedonic organoleptic test of the panelists gave a favorable response for all criteria of appearance, texture, color, aroma, and taste.

Keywords: *Lutjanus* sp.; characteristics; food; fish bone.

Practical Application: The practical implication of this article is that the addition of nanomineral fish bones from the family *Lutjanus* sp. to Saddah Gohoge grits can improve its nutritional content without significantly altering its sensory characteristics. This study provides evidence that the use of nanominerals as a fortificant is a promising approach to enhance the nutritional value of traditional foods without compromising their taste and texture. The findings of this study can be useful for food manufacturers and policymakers in developing countries who are interested in improving the nutritional status of their population through the fortification of traditional foods. Additionally, the use of nanominerals can be a more cost-effective and sustainable approach to fortification, as it requires less fortificant material and does not alter the taste and texture of the food.

1 Introduction

Porridge is a food in the weaning period in infancy and breakfast food (Om et al., 2019). The characteristics of slurry porridge are high flour content, flour and water ratio is 5-15% w/v, and thick and little nutritional composition porridge is a traditional food, especially for very low-income families (Neupane et al., 2022).

The traditional Rowe Luwa porridge in Southwest Sumba, Indonesia is a green porridge made from cassava leaves, and tubers pounded together with rice which is made in an instant dry form to extend the shelf life of the porridge (Iwansyah et al., 2022). Cornmeal-based porridge in Nigeria is “Ugali” or “Ogi”, in Italy “Polenta” in Brazil “Angu” and in Romania “Mamaliga”. In this study, the traditional Gorontalo grits in Indonesia called “Saddah” or “Gohoge” is made from corn flour.

Maize (*Zea mays* L) is the second most important cereal crop in the world in terms of production and consumption (Bustillos-Rodríguez et al., 2019). Further, Maize food processing products are indispensable in daily life, thus maize is widely cultivated in other countries (Zhu et al., 2022). In Brazil, maize production was 54.5 million tonnes in 2015, with significant quantities produced by maize farmers in central and southern Brazil (Kelte et al., 2018). Maize is composed of two glucose homopolysaccharides, amylose and amylopectin, and has a linear

and branched structure (Liu et al., 2002). The starch content in corn as a functional food is widely applied to food and non-food because it can modify physicochemical, thermal, and rheological properties (Bello-Pérez et al., 1999; Waterschoot et al., 2015; Bustillos-Rodríguez et al., 2019). Corn mainly supplies starch, and its flour is used in home cooking as a food source and processed food products (Neupane et al., 2022; Barretti et al., 2022).

Saddah Gohoge grits is a local name for food from the Gorontalo province in Indonesia, which is generally popular, especially for the elderly and growing children, because it has a soft texture and corn flavor. These grits contains high carbohydrates, namely 95.43% and fiber content of 3.8%. However, it is still very low in mineral nutrient content. Mineral nutrient intake is needed, especially during the growth period of children and the elderly. For the elderly aged >60 years, the calcium needed by the body ranges from 800-1000 mg/day, calcium intake in Indonesia has only reached 237 mg/day (Indonesia, 2019). Research in Rwanda revealed that grits marketed locally still lacks nutrients that is protein, fat and vitamins A and E. (Grosshagauer et al., 2020).

Mineral content can be obtained from fish bone waste from processed fish residue. Utilization of processed fish to produce new products can be done effectively and increase income and contribute to environmental conservation. Processed fish waste is

Received 23 Sept., 2022

Accepted 06 Nov., 2022

¹Lecturers of Fishery Product Technology, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo Province, Indonesia

*Corresponding author: rahim@ung.ac.id

divided into two groups, namely used to increase animal or plant production and the development of value-added by-products for human consumption, namely stomach contents, scales, bones including skull bones to be used as flour, oil, silage and compost for use in animal feed or fertilizer in plants (Kumar et al., 2022). Fish bones have a fairly high mineral content but are still in the form of macrominerals and have not been completely absorbed by the body. The main constituents of fish bones as an important source of minerals are calcium, sodium (5.63 g/kg), phosphorus (2.38 g/kg), and carbonates. In addition, fish bones contain inorganic salt minerals, namely calcium phosphate, creatine phosphate and hydroxyapatite $[Ca_{10}(OH)(PO_4)_6]$ in the form of crystals attached to fibrillar collagen (Malde et al., 2010; Suptijah et al., 2012).

One of the processed fishery waste is fish bones from the *Lutjanus* sp. This family of fish is found in Indonesian waters, including in the waters of the Gorontalo area. Types of fish family *Lutjanus* sp. There are 33 species in Indonesian waters. Types of fish belonging to the family *Lutjanus* sp. Most live around coral reef areas (Oktaviyani, 2018).

The particle size of minerals as feed additives in nanoparticle form is claimed to be smaller than 100 nanometer, it can pass through the stomach wall and into body cells more quickly than ordinary minerals with larger particle size (Singh & Pankaj, 2016).

The fulfillment of protein and fat nutrients as well as vitamins A and E in Saddah Gohoge corn grits can be served with fish sauce and boiled vegetables. Fish bones can be used as an inexpensive natural material that contains calcium phosphate. Previous research produced calcium phosphate in micro and nanoparticle sizes from fish bones. Fish bone waste originating from household consumers, processed fish fillet industries into the aquatic environment can have an unfavorable effect. Utilization of fish bone waste can be used as a product that provides beneficial value, especially in the growth and maintenance of healthy bones and teeth. Fish bones derived from processed fish fillet waste can be processed into nanominerals (nm) and can be fortified in food. Through nanotechnology for application to food, one of them is increasing the bioavailability of nutrients (Greiner, 2009; Lekahena et al., 2014). One of the additions of nutrients to the Saddah Gohoge corn grits is by fortifying nanominerals derived from fish bone waste from the *Lutjanus* sp. This study carried out fortification of nanomineral waste from fish bone *Lutjanus* sp. and perform physical, chemical and organoleptic characteristics based on the response of the panelists to the Saddah Gohoge grits.

2 Materials and methods

This research was conducted in April 2022-September 2022 at the Laboratory of the Faculty of Mathematics and Natural Sciences, the Laboratory of Biotechnology and Characteristics of Fishery Products, Faculty of Fisheries and Marine Sciences, State University of Gorontalo, Laboratory of Quality Testing of Fishery Products of Gorontalo Province and the Gorontalo Polytechnic Laboratory.

2.1 Materials and tools

Saddah Gohoge corn flour and salt are purchased at the Gorontalo traditional market. Fish bone waste of the family

Lutjanus sp. Taken from the rest of the processed fillets in the fishery processing industry of PT.99 Tinakin Laut, Banggai Laut Regency, Central Sulawesi Province. NaOH, HCl, H₂SO₄, Na₂CO₃, KIO₃, CuSO₄.5H₂O, (NH₄)₂HPO₄, 1% phenolphthalein, LuffSchoorl solution were purchased from an analytical chemical agent in Jakarta.

The tools used are Atomic Absorption Spectrophotometry (AAS) (Perkin Elmer Analysis 100 flame emission type), High Performance Liquid Chromatography (HPLC) (Shimadzu Prominence) Scanning Electron Microscope (SEM) Phenom ProX G5. Field Emission Scanning Electron Microscopy (FE-SEM) and Electron Diffraction X-ray, crystallinity morphology analysis with XRD tool (Venkatesan & Kim, 2010). Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) (Iwansyah et al., 2022).

2.2 Sample preparation

Fish bone preparation

Fishbone preparation methods is washing stage, boiling the bone at a temperature of ± 130 °C or 250 °F, drying, reducing and screening of fish bone powder modification (Harmain et al., 2018). The first stage was carried out to produce fishbone powder with several stages of the process, that is washing, boiling, drying and size reduction. The second stage of fish bone powder was divided into 3 treatment methods is milling, HCl and NaOH. (Prinaldi et al., 2018). Nanocalcium using the *precipitation* method with prawn shell soaking time for 48 hours (Suptijah et al., 2012).

Saddah Gohoge corn preparation

Corn flour was washed from dirt and dust using clean water, allowed to stand for a while and the filtrate was removed slowly, until the flour that had settled was clean. Then the corn flour is dried using an oven at 100-120 °C for 1-3 hours to dry or using a drum dryer at 120 °C at 8Hz speed. The process of making corn flour when using corn shells follows corn flour production by Houssou & Ayemor (2012) and Om et al., (2019) modification (Figure 1).

Formulation of composites of Saddah Gohoge corn with cooking water based on *trial and error* of 1:5 w/w. Fish bone nanomineral fortification formulation fish bone of family *Lutjanus* sp. are 10%, 15% and 20%. For Saddah Gohoge were formulated to produce composite grits while the control sampel used was 100% Saddah Gohoge. As many as 200 g Saddah Gohoge corn flour was mixed and homogenized with fortification of fish bone nanominerals from the family *Lutjanus* sp. and then cooked with 1000 mL (1:5) cooking water at a temperature of 100 °C for ± 45 minutes..

2.3 Procedure of analyses

Organoleptic test

Samples of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. with concentrations of 0%, 10%, 15% and 20% presented together with a hedonic and hedonic




Figure 1. Process flow for the production of maize flour.

quality organoleptic assessment score sheet, the panelists used were untrained panelists and came from consumers who used to consume Saddah grits as many as 30 people with a scale of 1 – 5 based on the parameters of appearance, texture, color, aroma and taste through an assessment in the score sheet provided. Organoleptic testing uses panelists who are divided into trained, semi-trained and untrained panelists with a certain scale range based on an assessment of organoleptic parameters (Lawless & Heymann, 2010).

Proximate analysis

Proximate analysis of Saddah Gohoge grits consisting of water content, ash content, protein content, fat content, crude fiber content and carbohydrate content can be carried out based on the Association of Official Analytical Chemists (2005) method. The Association of Official Analytical Chemists (2005) and Iwansyah et al. (2022) method for determining water content and ash content can be determined using the gravimetric method. Determination of protein content used equipment Buchi-Dumaster, fat content using the Weibull method (Association of Official Analytical Chemists (2005; Iwansyah et al., 2022), carbohydrate content can be calculated by difference, fiber content can be determined using the method of Association of Official Analytical Chemists (2005).

Viscosity measurement

Measurement of slurry viscosity using a Rapid Visco Analyzer (RVA-Techmaster, Macquarie Park, Australia) connected to a computer (Amagloh et al., 2013). The slurry sample was weighed 3 g and then mixed with 25 mL of distilled water in an

aluminum container. The sample was then put into the Rapid Visco Analyzer with a rotation speed of 100 rpm at 25 °C for 2 minutes. The sample was then heated to 95 °C for 5 minutes, then held at 95 °C for 3 minutes. After that, the sample was cooled again to 50 °C in 4 minutes and then held at 50 °C for 2 minutes (Iwansyah et al., 2022).

Coloring analysis

The staining of the slurry samples used an NH 310 colorimeter (Indriati et al., 2020) based on the analytical method, namely CIE (Commission Internationale de L'Eclairage) L^* a^* b^* and hue coordinates. The L^* coordinates describe where $L = 0$ is black, and $L^* = 100$ is colorless. The coordinates of a^* represent red and green shadows, where $a^* > 0$ indicates red and $a^* < 0$ indicates green. The b^* coordinates represent the blue and yellow tones, where $b^* > 0$ indicates the intensity of the yellow color and $b^* < 0$ indicates the blue color. Hue (h^*) is a color characteristic, namely red, yellow, green, and blue (Iwansyah et al., 2022).

Determination of mineral grade

Nanomineral analysis of fish bone meal of the family Lutjanus sp. can be done based on several methods of ashing (ash content). According to Ciobanu et al. (2013), a sample of 20-30 g was dried in an oven at 105 °C followed by calcination at 450 °C, then 5 mL of HNO₃ was added, and stored in a desiccator at 150 °C to dissolve the ash. the remaining. Tejera et al. (2013) digested a sample of 10 g of wheat flour with 2 mL of concentrated HNO₃ and then the sample was calcined at a temperature of 450 °C for 48 hours. Pourhossein & Shalaei (2016) determined Fe in flour samples after being calcined in a furnace at 600 °C for 10 hours, adding 10 mL of HCl (1 M) and neutralized by adding 10 mL of NaOH until the volume became 100 mL.

2.4 Statistical analysis

The research design used a completely randomized design with 1 treatment and 6 levels, namely the concentration of fish bone nanominerals Lutjanus sp. namely 0%, 10%, 15% and 20%. The results organoleptic, physical and proximate test were subjected to statistical analysis to determine possible differences among samples by LSD's multiple range test using the SPSS programme Significant differences were expressed at $p < 0.05$. Data were presented as mean \pm standard deviation (sd) ($n = 4$). Normality test was carried out on the data, and Analysis of Variance (ANOVA) was used to determine the significantly differences between treatments (Statistical Package for Social Sciences, 2022; Rao, 2009).

3 Results and discussion

3.1 Organoleptic analysis of the Saddah gohoge grits

The sample of Saddah Gohoge corn grits fortification of fish bone nanominerals from the family Lutjanus sp. What has been prepared is hedonic and hedonic quality organoleptic testing along with the score sheet provided (Figure 2).

Hedonic quality analysis

The results of organoleptic tests include quality hedonic and hedonic tests based on appearance, texture, color, aroma and taste criteria (Table 1, Table 2).

Based on the results of hedonic quality organoleptic testing, it was found that the highest value fortification of the fortification concentration of fish bones of the *Lutjanus* sp. 10% with a score of 3.40 the assessment criteria are rather bright. The texture of the highest value at 20% concentration fortification scores a score of 3.48 with the assessment criteria being slightly smooth and slightly thick. Texture is the nature of the product that can be felt through the touch of the skin or tasting and affects the quality of the product which ultimately affects consumer acceptance of the product (Ahmad et al., 2018).

The color parameter obtained the highest value of hedonic quality with an assessment score of 3.08 the assessment criteria are slightly yellowish. Saddah Gohoge grits with yellow corn

as raw material because it contains carotenoids in addition to vitamin A and vitamin E (Table 1). An important indicator of the acceptance of a food product is flavor (Yang et al., 2022). The flavor of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. 10% obtained a hedonic quality value of 4.00 which is slightly pungent with the flavor of corn criteria.

The hedonic quality score of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. The highest score in a score is 3.16 on the taste parameter, the criteria are rather good, the taste of corn. Analysis of variance obtained that appearance, texture, color and flavor were not significant but on taste there were significant concentrations of fortification 15% and 20% (Table 1).

Hedonic analysis

The results of hedonic organoleptic testing for appearance parameters obtained the highest value for the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. 0% with a




Figure 2. Serving of Saddah Gohoge grits fortification of fish bone nanomineral formulation family *Lutjanus* sp. (a = 0%, b = 10%, c = 15%, d = 20%).

Table 1. Hedonic quality organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.40 ± 1.041 ^a	3.40 ± 0.913 ^a	3.08 ± 0.909 ^a	3.64 ± 1.075 ^a	3.16 ± 1.248 ^a
F1 (90: 10)	3.68 ± 0.945 ^a	3.28 ± 1.021 ^a	2.88 ± 0.881 ^a	4.00 ± 0.866 ^a	2.96 ± 1.207 ^a
F2 (95:15)	3.44 ± 1.044 ^a	3.36 ± 1.551 ^a	3.00 ± 1.000 ^a	3.96 ± 1.060 ^a	2.56 ± 0.917 ^{ab}
F3 (80:20)	3.32 ± 1.145 ^a	3.48 ± 1.503 ^a	2.76 ± 1.052 ^a	3.96 ± 4.00 ^a	2.24 ± 1.200 ^b

a: no significant difference between the treatments. b, ab: there is significant difference between treatments.

Table 2. Hedonic organoleptic test results.

Concentration (%)	Appearance	Texture	Color	Flavour	Taste
F0 (100:0)	3.96 ± 0.841 ^a	3.64 ± 1.114 ^a	4.04 ± 0.935 ^a	3.80 ± 1.041 ^a	3.52 ± 1.194 ^a
F1 (90: 10)	3.64 ± 0.995 ^a	3.36 ± 1.319 ^a	3.96 ± 0.889 ^a	3.64 ± 0.907 ^a	3.28 ± 1.100 ^{ab}
F2 (95:15)	2.96 ± 0.790 ^b	3.12 ± 0.927 ^a	3.76 ± 0.879 ^a	2.96 ± 0.935 ^b	2.80 ± 0.913 ^{bc}
F3 (80:20)	2.80 ± 1.000 ^b	2.84 ± 1.143 ^a	3.40 ± 1.118 ^a	2.88 ± 1.013 ^b	2.56 ± 1.044 ^c

a: no significant difference between the treatments. b, c, ab, bc: there is significant difference between treatments.

rating score of 3.96 likes rating. The results of the analysis of significant variance in the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. on texture and color parameters but significant on appearance, flavor and taste (Table 2). The texture parameter obtained the highest value at 0% concentration fortification with an assessment score of 3.64 the criteria were rather like. This study is in line with the results of research by Ahmad et al. (2018) which found that the panelists fortified corn grits fortified cherry leaves (*Muntingia calabura* L) gave a neutral to moderate response on a scale of 4.1-4.9.

The texture of Saddah Gohoge grits with different viscosities according to the given nanomineral fortification also depends on the amylopectin content which affects sensory, especially texture and taste. Types and varieties of corn have different amylopectin content. This is in line with what was stated by Singh et al. (2006); Rodriguez et al. (2018) that the physicochemical properties of corn starch can change the function of polysaccharides. Starch contributes to the textural properties of various corn products, namely as a gelling agent, thickener, adhesive, colloid stabilizer and water retention agent.

The results of hedonic organoleptic testing obtained the highest value for fortification of fish bone nanominerals from the family *Lutjanus* sp. 0% score 3.08 on the color parameter of the assessment criteria is rather like. This is the same as the research by Ahmad et al. (2018) on fortified corn grits of cherry leaves (*Muntingia calabura* L) which obtained a score of 4.3 – 5.4 with the criteria of somewhat liking.

The concentration of fortified nanomineral fish bones of the *Lutjanus* sp. (0%) for the parameters of flavor and taste, respectively, the highest scores were 3.80 and 3.52 with the criteria of each rating being somewhat like. The results of research by Ahmad et al. (2018) showed that the fortified corn grits of cherry leaves (*Muntingia calabura* L) panelists responded with a score of 4.3-4.6 on the aroma and taste parameters with the assessment criteria being somewhat like.

3.2 Physical analysis

Viscosity analysis

The results of the physical test of the thickness of the Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the higher the concentration of fortified nanominerals (Table 3).

The results of measuring the thickness of the sample of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. obtained the highest value of 14069.7333 cP at a concentration of 15% (Table 3). These results are influenced by the addition of the volume of cooking water. This was because the volume of boiled

water for Saddah Gohoge grits was the same at all concentrations of nanomineral fortification (1000 mL). This can be added to the volume of cooking water if want to achieve the same viscosity with nanomineral fortification. The results of statistical analysis showed that all fortification concentrations were significant ($p < 0.05$). The results of research by Iwansyah et al. (2022) show the relationship between pre-gelatinization temperature and viscosity with the correlation coefficient (r) and coefficient of determination (R^2) being 0.9924 and 0.9849, respectively (Iwansyah et al., 2022). The thickness of the instant grits that has been brewed varies from slightly thick to thick. This is because the greater the concentration of addition of white rice flour, the higher the value of instant grits viscosity (Makame et al., 2019). The pre-gelatinization process is a modification of starch, carried out through a process involving water and heat to break down all or part of the granules, then dried to produce a complete or partial pre-gelatinization of starch (Wadchararat et al., 2006; Palguna et al., 2014; Iwansyah et al. 2022). Several factors influence this pre-gelatinization process, including temperature (Iwansyah et al., 2022).

Coloring analysis

The results of the color test of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp. showed that the staining at all concentrations of fortification ranged from 47.10-50.03 for the L component. different values for *a are -39.23-3.76 and *b values 15.96-16.50. Statistical analysis results obtained on all staining components were not significant ($p > 0.05$) (Table 4). The results of this staining analysis are the same as the results of the hedonic quality organoleptic test, which is not significant ($p > 0.05$) for the color parameter.

The results of the staining study were suspected because Saddah Gohoge flour basically has the same color, namely a yellowish color and the fortification concentration given does not exceed 20% so it does not affect the basic color of Saddah gohoge corn flour, which is yellow. The coloring component is the same as the results of the research by Iwansyah et al. (2022) in the rowe luwa grits sample which was not significant for *L and hue values ($p > 0.05$) but significant for the components of *a and *b values ($p > 0.05$). This shows that the cooking temperature of gelatinization also affects the coloring component of the slurry. The enzymatic reaction is slower with the longer pre-gelatinization time of Gayam flour (*Inocarfus fagifer* Forst.) (Wijanarka et al. (2017). The longer pre-gelatinization time will also increase the temperature and inactivate polyphenol oxidase (Akyıldız & Ocal, 2006).

3.3 Proximate analysis

Proximate analysis include water content, ash content, protein content, fat content, fiber content and carbohydrate content.

Table 3. Viscosity levels of Saddah gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of the viscosity		
	F1	F2	F3
Viscosity levels (cP)	8309.4667 ± 42.46285 ^a	14069.7333 ± 103.87946 ^b	10264.7000 ± 123.87946 ^c

a: no significant difference between the treatments. b, c: there is significant difference between treatments.

The results of the analysis of the highest water content without fortification of nanomineral fish bones of the *Lutjanus* sp. 0% as a fortification concentration increased, the water content decreased with the use of the volume of water for cooking Saddah Gohoge grits which was the same as 1000 mL for all concentrations (Table 5).

The results of the LSD further test showed that the concentration of 25% was significantly different from 0%, 15% and 20% ($p < 0.05$). This is due to the absorption of nanominerals into the corn *grits* granules. The type, polarity of the absorbate, the type of bond, the size of the absorbate, the viscosity of the mixture are factors that affect absorption.

The statistical results of the research by Ahmad et al. (2018) were significant ($p < 0.05$) for all concentrations of fortification of cherry leaves (*Muntingia calabura* L) in corn grits allegedly due to the absorption of cherry extract into corn grits granules. The results of the research by Iwansyah et al. (2022) found that the water content was not significant ($p > 0.05$) along with the increase in temperature of the instant rowe luwa grits with pre-gelatinization with the highest value of water content at 80°C, namely 6.78. The results of the analysis of ash content showed that it increased with each addition of the concentration of fortified nanomineral fish bones of the *Lutjanus* sp. This is because there are different nanomineral content at each concentration which affects the amount of ash content at each concentration.

The results of the further test of ash content at 15% and 20% and 25% fortification concentrations were not significantly ($p > 0.05$) but significantly different from those without nanomineral fortification ($p < 0.05$).

This is also inline with Iwansyah et al. (2022) study that obtained statistical results that the ash content was not significant ($p > 0.05$) with increasing temperature in pre-gelatinized instant Rowe Luwa grits samples.

Protein content, fat content and fiber content at each concentration of fortification increased along with the fortification

of nanomineral fish bones of the *Lutjanus* sp. which are given. The results of further test of protein content showed that all fortification concentrations were significantly different.

The results of the study on the protein content of fortified Saddah Gohoge grits were the same as those of Ahmad et al. (2018), which was significant ($p < 0.05$). The protein content in this study increased along with the increased fortification concentration carried out. This is presumably because there is protein content in the fish bones of the *Lutjanus* sp.

Protein and carbohydrate levels in the study of Iwansyah et al. (2022) were also significant ($p < 0.05$) along with the increase in temperature in pre-gelatinized Rowe Luwa grits. The interaction between starch and protein in food can increase the gel strength which is associated with an increase in the density of the protein matrix (Couto et al., 2012; Jamilah et al., 2009; Iwansyah et al., 2022).

The fat content showed that the fortified concentration of 15% was significantly ($p < 0.05$) different with the concentration of 20% and 25% and without fortification (0%) but the concentration of fortified 20% and 25% was not significantly different ($p > 0.05$). The fat content increased along with the fortification of the bone nanomineral concentration of the fish of the *Lutjanus* sp. on grits Saddah Gohoge. The results of this study are the same as those of Iwansyah et al. (2022), which is not significant ($p > 0.05$) with increasing pre-gelatinization temperature of Rowe Luwa grits.

For fiber content, the concentration of fortified of 15%, 20%, and 25% not significantly different but significantly different from without fortification (0%). This is because there is a fiber content in the Saddah Gohoge corn grits.

The results of the further test of carbohydrate content showed that all fortification concentrations were significantly different.

3.4 Mineral analysis

Fish bone nanominerals family *Lutjanus* sp. To be fortified in Saddah Gohoge grits, it mainly contains minerals Ca which

Table 4. Color of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Component	Results of sample coloring rate		
	F1	F2	F3
L	47.10 ± 2.60 ^a	51.33 ± 1.01 ^a	50.03 ± 3.55 ^a
*a	3.76 ± 3.00 ^a	-39.23 ± 74.27 ^a	2.03 ± 1.74 ^a
*b	15.96 ± 2.45 ^a	16.30 ± 2.26 ^a	16.50 ± 2.60 ^a

a: no significant difference between the treatments.

Table 5. Proximate levels of Saddah Gohoge grits fortification of fish bone nanominerals *Lutjanus* sp.

Sample	Water Content	Ash Content	Protein Content	Fat Content	Fiber Content	Carbohydrate Content
0%	86.82 ± 0.411 ^a	0.92 ± 0.87 ^a	7.16 ± 0.90 ^a	0.47 ± 0.95 ^a	1.77 ± 0.54 ^a	95.37 ± 0.50 ^a
15%	85.03 ± 0.744 ^b	1.13 ± 0.26 ^{ab}	8.53 ± 0.72 ^b	0.74 ± 0.41 ^b	3.30 ± 0.49 ^b	95.43 ± 1.52 ^b
20%	80.33 ± 0.562 ^b	1.31 ± 0.27 ^{ab}	9.63 ± 0.47 ^c	1.21 ± 0.87 ^c	3.60 ± 0.37 ^b	92.48 ± 0.40 ^c
25%	79.70 ± 0.113 ^c	1.53 ± 0.25 ^b	10.51 ± 0.30 ^d	1.22 ± 1.36 ^c	3.8 ± 0.37 ^b	92.96 ± 0.61 ^d

a: no significant difference between the treatments. b, c, d, ab: there is significant difference between treatments.

Table 6. The composition of nanomineral fish bones of the *Lutjanus* sp.

Number	Minerals	Unit	Result
1	Nitrogen (N)	%	2.10
2	Potassium (K)	mg/100 g	36.67
3	Magnesium (Mg)	mg/100 g	555.21
4	Manganese (Mn)	mg/100 g	30.48
5	Phosphorus (P)	mg/100 g	148409.80
6	Iron (Fe)	mg/100 g	27.31
7	Calcium (Ca)	mg/100 g	28556.18

is 30.89% and P 15.22%. Saddah Gohoge grits made from corn also contains various essential minerals, namely K, Na, Ca, P and Fe. The results of the overall nanomineral analysis on fish bones of the *Lutjanus* sp. family (Table 6).

The results of the analysis of the nanomineral composition in the fish bones of the *Lutjanus* sp. The highest levels were obtained in Phosphorus (P) followed by Calcium (Ca), Magnesium (Mg), Potassium (K), Manganese (Mn), Iron (Fe) and Nitrogen (N) (Table 6). The highest levels of nanominerals were found in fish bone waste. This is the same as research by Harmain et al. (2018) which obtained nanomineral levels in skipjack bones, namely phosphorus (P) 6.841%, calcium (Ca) 2.935%, Magnesium (Mg) 0.528%, Manganese (Mn) 0.014%, iron (Fe) 0.016%, Potassium (K) 0.002% and zinc (Zn) 0.0089%.

Fortification of fish bone nanominerals in one serving of *binte biluhuta* cream soup or corn cream soup as a high-calcium food product has met the calcium needs of the elderly (>50 years old) which is 51.47%. Furthermore, a product can be claimed to be high in calcium if it has met twice the calcium source material (Riyanto et al., 2020).

4 Conclusion

Fortification of fish bone nanominerals of the family *Lutjanus* sp. on Saddah Gohoge grits from organoleptic hedonic quality, the appearance parameters are slightly bright - bright, texture is slightly smooth, slightly thick - smooth thick, slightly yellowish - yellowish in color, slightly overpowering aroma of corn - pungent aroma of corn, slightly delicious taste of corn - slightly good taste of corn, hedonic ranges from a bit like and like. Viscosity, coloration and proximate increased with fortification of the formulation but decreased in water content.

Acknowledgements

We would like to thank the Institute for Research and Community Services (LP2M) of Universitas Negeri Gorontalo, Indonesian Endowment Fund for Education (LPDP) of Ministry of Finance, the reviewers of this article and all those who have helped carry out the research.

References

Ahmad, M., Benjakul, S., Prodpran, T., & Maqsood, S. (2018). Characterization of gelatin from goat skin and its application in fish mousse. *International Journal of Biological Macromolecules*, 108, 188-194.

Akyıldız, A., & Ocal, N. D. (2006). Effects of dehydration temperatures on colour and polyphenoloxidase activity of amasya and golden delicious apple cultivars. *Journal of the Science of Food and Agriculture*, 86(14), 2363-2368. <http://dx.doi.org/10.1002/jsfa.2624>.

Amagloh, F. K., Mutukumira, A. N., Brough, L., Weber, J. L., Hardacre, A., & Coad, J. (2013). Carbohydrate composition, viscosity, solubility, and sensory acceptance of sweetpotato- and maize-based complementary foods. *Food & Nutrition Research*, 57, 18717. <http://dx.doi.org/10.3402/fnr.v57i0.18717>. PMID:23516115.

Association of Official Analytical Chemists – AOAC. (2005). *Official methods of analysis: method 979.21* (18th ed.). Washington, DC: AOAC International.

Barretti, B. R. V., Soltes de Almeida, V., Ito, V. C., Silva, B. M., Carvalho, M. A. S., Fo, Sydney, E. B., Demiate, I. M., & Lacerda, L. G. (2022). Combination of organic acids and heat-moisture treatment on the normal and waxy corn starch: thermal, structural, pasting properties, and digestibility investigation. *Food Science and Technology (Campinas)*, 42(19), 1-7. <http://dx.doi.org/10.1590/fst.33120>.

Bello-Pérez, L. A., Agama-Acevedo, E., Sánchez-Hernández, L., & Paredes-López, D. (1999). Isolation and partial characterization of banana starches. *Journal of Agricultural and Food Chemistry*, 47(3), 854-857. <http://dx.doi.org/10.1021/jf980828t>. PMID:10552380.

Bustillos-Rodríguez, J. C., Tirado-Gallegos, J. M., Ordóñez-García, M., Zamudio-Flores, P. B., Ornelas-Paz, J. J., Acosta-Muñoz, C. H., Gallegos-Morales, G., Páramo-Calderón, D. E., & Rios-Velasco, C. (2019). Physicochemical, thermal and rheological properties of three native corn starches. *Food Science and Technology (Campinas)*, 39(1), 149-157. <http://dx.doi.org/10.1590/fst.28117>.

Ciobanu, C., Şlencu, B. G., & Cucureanu, R. (2013). Faas determination of cadmium and lead content in foodstuffs from north-eastern Romanian Market. *Studia Universitatis "Vasile Goldiş". Seria Ştiinţele Vieţii*, 23(1), 33-38.

Couto, A., Enes, P., Peres, H., & Oliva-Teles, A. (2012). Temperature and dietary starch level affected protein but not starch digestibility in gilthead sea bream juveniles. *Fish Physiology and Biochemistry*, 38(3), 595-601. <http://dx.doi.org/10.1007/s10695-011-9537-5>. PMID:21728054.

Greiner, R. (2009). Current and projected applications of nanotechnology in the food sector aplicações atuais e futuras da nanotecnologia no setor alimentício. *Brazilian Society of Food and Nutrition*, 34(1), 243-260. Retrieved from http://sban.cloudpanel.com.br/files/revistas_publicacoes/228.pdf

Grosshagauer, S., Milani, P., Kraemer, K., Mukabutera, A., Burkon, A., Pignitter, M., Bayer, S., & Somoza, V. (2020). Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Maternal and Child Nutrition*, 16(1), e12856. <http://dx.doi.org/10.1111/mcn.12856>. PMID:31183951.

Harmain, R. M., Dali, F. A., & Husain, R. (2018). Nanocalcium characterization of cakalang fish bone flour (*Katsuwonus pelamis* L). *International Journal of Innovative Science and Research Technology*, 3(10), 306-368.

Houssou, P., & Ayemor, G. S. (2012). Appropriate processing and food functional property of maize flour. *African Journal of Science and Technology*, 3, 126-131.

Indonesia. (2019, October 28). Regulation of the Minister of Health regarding Communitie Health Centers (Regulation of the Minister of Health Number 43 of 2019). *Berita Negara Republik Indonesia*. BN.2019/NO.1335, PERATURAN.GO.ID: 40 HLM. Retrieved from <https://peraturan.bpk.go.id/Home/Details/138635/permenkes-no-43-tahun-2019>

- Indriati, A., Hidayat, D. D., Andriansyah, R. C. E., Iwansyah, A. C., & Surahman, D. N. (2020). Characterisation of physical, mechanical and colour properties of Muntingia calabura fruits. *IOP Conference Series: Earth and Environmental Science*, 462, 012044. <http://dx.doi.org/10.1088/1755-1315/462/1/012044>.
- Iwansyah, A. C., Apriadi, T., Arif, D. Z., Andriana, Y., Indriati, A., Mayasti, N. L., & Luthfiyanti, R. (2022). Effect of pre-gelatinized temperature on physical and nutritional content of Indonesian instant cassava leaves porridge: Rowe Luwa. *Brazilian Journal of Food Technology*, 25, e2021050. <http://dx.doi.org/10.1590/1981-6723.05021>.
- Jamilah, B., Mohamed, A., Abbas, K. A., Rahman, R. A., Karim, R., & Hashim, D. M. (2009). Protein-starch interaction and their effect on thermal and rheological characteristics of a food system: a review. *Journal of Food Agriculture and Environment*, 7(2), 169-174.
- Kelte, I., Fo., Butik, M., Jaski, A. C., & Quinaia, S. P. (2018). Fast method to determine the elements in maize flour: reduction in preparation time and reagent consumption. *Brazilian Journal of Food Technology*, 21, e2017091. <http://dx.doi.org/10.1590/1981-67239117>.
- Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2022). Utilization of fish processing waste: a waste to wealth approach. In V. Kumar, A. U. Muzaddadi, S. Mann, R. Balakrishnan, K. Bembem & Y. Kalnar (Eds.), *Emerging post-harvest engineering and technological interventions for enhancing farmer's income modified* (pp.127-13. Ludhiana: ICAR-CIPHET.
- Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: principles and practices*. USA: Springer Science & Business Media. <http://dx.doi.org/10.1007/978-1-4419-6488-5>
- Lekahena, V., Faridah, D. N., Syarief, R., & Peranginangin, R. (2014). Physicochemical characterization of nanocalcium extracted from Tilapia fish bone using alkaline and acid solution. *Jurnal Teknologi dan Industri Pangan*, 25(1), 57-64.
- Liu, Q., Charlet, G., Yelle, S., & Arul, J. (2002). Phase transition in potato starch-water system O. Starch gelatinization at high moisture level. *Food Research International*, 35(4), 397-407. [http://dx.doi.org/10.1016/S0963-9969\(01\)00134-X](http://dx.doi.org/10.1016/S0963-9969(01)00134-X).
- Makame, F. A., Asnani, A., Agustian, E., & Rohman, A. (2019). The effect of white rice flour addition on viscosity of instant grits. *Journal of Physics: Conference Series*, 1236(1), 012032.
- Malde, M. K., Bugel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: a double-blinded randomised crossover design. *Nutrition & Metabolism*, 7(1), 61. <http://dx.doi.org/10.1186/1743-7075-7-61>. PMID:20646299.
- Neupane, D., Adhikari, P., Bhattarai, D., Rana, B., Ahmed, Z., Sharma, U., & Adhikari, D. (2022). Does climate change affect the yield of the top three cereals and food security in the world? *Earth (Waukesha, Wis.)*, 3(1), 45-71. <http://dx.doi.org/10.3390/earth3010004>.
- Oktaviyani, S. (2018). Knowing the Lutjanus, one of the leading commodities in catch fisheries. *Jurnal Oseana*, 40(3), 29-39.
- Om, A., Ak, B., Fo, O., & Ge, Z. (2019). Nutritional composition and organoleptic properties of composite maize nutritional composition and organoleptic properties of composite maize porridge. *Food Processing & Technology*, 10(6), 1000798.
- Palguna, I. G. P. A., Sugiyono, S., & Hariyanto, B. (2014). Characteristics of modified sago starch by repeated gelatinization and retrogradation treatment. *Jurnal Pangan*, 23(6), 146-156.
- Pourhossein, M., & Shalaei, H. (2016). Determination of iron in different types of wheat flour consumed in Iran. *Journal of Agricultural Science and Technology*, 18(3), 573-582.
- Prinaldi, R., Nurwantoro, N., Sari, M. A., & Muryanto, S. (2018). Hydroxyapatite synthesis from fishbone waste via calcination method. *IOP Conference Series. Materials Science and Engineering*, 308(1), 012024.
- Rao, C. R. (2009). *Linear statistical inference and its applications* (2nd ed.). New York: John Wiley & Sons.
- Riyanto, B., Trilaksani, W., & Aulia Azzahra, V. (2020). Design of instant food special nutrition for the elderly based on binte biluhuta enriched with fish bone nanominerals. *Jurnal Fishtech*, 9(2), 65-77. <http://dx.doi.org/10.36706/fishtech.v9i2.9923>.
- Rodriguez, J. P., Anuniação, P. C., Garcia, E. E., & Franco, C. M. (2018). Structural and physicochemical properties of corn starches and their function as polysaccharides in food systems: a review. *Food Research International*, 103, 391-401.
- Singh, A. V., & Pankaj, P. (2016). Nanoparticle-mediated delivery of micronutrients in plants. In S. Chandra & H. K. Chauhan (Eds.), *Nanotechnology and plant sciences* (pp.265-291). New Delhi: Springer.
- Singh, N., Kaur, L., Sandhu, K. S., Kaur, J., & Nishinari, K. (2006). Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids*, 20(4), 532-542. <http://dx.doi.org/10.1016/j.foodhyd.2005.05.003>.
- Statistical Package for Social Sciences – SPSS. (2022). *SPSS 1001 for windows*. USA: SPSS Inc. Retrieved from <https://www.ibm.com/support/pages/spss-statistics-210-available-download>
- Suptijah, P., Jacob, A. M., & Deviyanti, N. (2012). Characterization and bioavailability of vannamei shrimp shell nanocalcium. *Jurnal Akuatika*, 3(1), 63-73.
- Tejera, R. L., Luis, G., González-Weller, D., Caballero, J. M., Gutiérrez, A. J., Rubio, C., & Hardisson, A. (2013). Metals in wheat flour; comparative study and safety control. *Nutrición Hospitalaria*, 28(2), 506-513. PMID:23822705.
- Venkatesan, J., & Kim, S. K. (2010). Nano-hydroxyapatite composite biomaterials for bone tissue engineering—a review. *Journal of Biomedical Nanotechnology*, 6(6), 1-12. PMID:25992432.
- Wadchararat, C., Masubon, T., & Naivikul, O. (2006). Characterization of pregelatinized and heat moisture treated rice flours. *Witthayasan Kasetsat Witthayasat*, 40, 144-153.
- Waterschoot, J., Gomand, S. V., Fierens, E., & Delcour, J. A. (2015). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Stärke*, 67(1-2), 14-29. <http://dx.doi.org/10.1002/star.201300238>.
- Wijanarka, A., Sudargo, T., Harmayani, E., & Marsono, Y. (2017). Effect of pre-gelatinization on physicochemical and functional properties of gayam (*Inocarpus fagifer* forst.) flour. *American Journal of Food Technology*, 12(3), 178-185. <http://dx.doi.org/10.3923/ajft.2017.178.185>.
- Yang, G., Zhang, W., Li, J., & Feng, B. (2022). Differentiation of fatty acid, amino acid, and volatile composition in waxy and non-waxy proso millet. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.58320>.
- Zhu, Y., Ma, Z., Han, M., Li, Y., Xing, L., Lu, E., & Gao, H. (2022). Quantitative damage detection of direct maize kernel harvest based on image processing and BP neural network. *Food Science and Technology (Campinas)*, 42(19), 1-9. <http://dx.doi.org/10.1590/fst.54322>.