



**BUKTI KORESPONDENSI**  
**ARTIKEL JURNAL TERAKREDITASI INTERNASIONAL**

Judul Artikel : Organic Fertilizer Application to Improve Bacterial Quality and Rice Production  
Jurnal : Jurnal International Journal of Life Science Research Archive  
ISSN : 0799-6640 (Online)  
Journal Homepage : <https://sciresjournals.com/>  
Volume : Volume 4 - Issue 1 (January - March 2023)  
Halaman : 117 - 122  
Article DOI : 10.53771/ijlsra.2023.4.1.0017  
DOI url : <https://doi.org/10.53771/ijlsra.2023.4.1.0017>  
Penulis : Zulzain Ilahude  
Hayatiningsih Gubali  
Mohamad Lihawa  
Mohamad Ikbah Bahua  
Corresponding author : Mohamad Ikbah Bahua  
Email Correspondent : [mohamad.bahua@ung.ac.id](mailto:mohamad.bahua@ung.ac.id)

## 1. Bukti Submit Artikel Tanggal 18 Januari 2023



## 2. Bukti Konfirmasi Submit Artikel dan Artikel yang di Submit Tanggal 19 Januari 2023



# Organic Fertilizer Application to Improve Bacterial Quality and Rice Production

Zulzain Ilahude<sup>1</sup>, Hayatiningsih Gubali<sup>2</sup>, Mohamad Lihawa<sup>3</sup>, Mohamad Ikbah Bahua<sup>4\*</sup>

<sup>1,2,3,4</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas Negeri Gorontalo, Indonesia

\*Corresponding author. Mohamad Ikbah Bahua, Email: [mohamad.bahua@ung.ac.id](mailto:mohamad.bahua@ung.ac.id)

## Abstract

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil. This study aims to determine the increase in the quality of bacteria and production of rice plants. The research was carried out in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The research using Completely Randomized Block Design with 6 treatments 3 replications. The plant used is rice Cakrabuana variety. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo. The plot size used is (10 x 5) m. The bacterial population was calculated using the SANICHECK kit bacterial test. The dose of Granul organic fertilizer that uses 10 tons and 20 tons/hectare. The research results show that Granule organic fertilizer 20 tons/hectare can improve the quality of bacteria from 10.13 to 10.17 cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is the best treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare.

**Keywords:** Organic Fertilizers, Bacterial Quality, Rice Production

## 1. Introduction

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil (Milosevic *et al.*, 2022). According to Weerahewa and Dayananda (2023), organic fertilizers can reduce environmental pollution and improve land quality in a sustainable manner. In principle, the organic farming system is a farming system that adheres to natural principles in building agro-ecosystem balance, so that it is beneficial to soil, water, air, plants and all living things, and is able to provide healthy food for human needs (Puech and Stark, 2023).

The addition of organic fertilizers to the soil, especially in soils with low organic matter, is an effort to ameliorate the soil so that the provision of plant nutrients can be more effective (Chen *et al.*, 2022). In general, the application of organic fertilizers to the soil will improve the physical, chemical and biological properties of the soil (Hui *et al.*, 2022). According to Gao *et al.* (2021), in soils that lack organic matter and degraded soils, organic fertilizer is the main requirement for soil amelioration, so that the provision of nutrient inputs is more efficient and effective.

Technology that can increase fertilization efficiency and the sustainability of rice production systems is microbial fertilizer technology contained in organic fertilizers (Hameed *et al.*, 2023). Supply of some of the N, P, K nutrients needed by rice plants can be carried out by rhizosphere bacteria, endophytic, green manure which have the ability to fix N<sub>2</sub> from the air, Phosphate solubilizing microbes can help mine Phosphate in the soil, so that it becomes available P for plants (Negi *et al.*, 2022). Microbial groups as biological agents that are widely used in organic fertilizers include Azotobacter Sp, Lactobacillus Sp, Alkaligenes sp, Rhizobium Sp, Pseudomonas Sp and Endophytic Bacteria such as *Ovhrbactrum pseudogignonense*, *Azospirillum*, and *Penicillium* Sp (Allouzi *et al.*, 2022).

According to Sahab *et al* (2021), in paddy fields waterlogging greatly affects the microbial metabolic processes of rice plants, such as; eukaryotic, aquatic weeds, heterotrophic microflora and fauna including roots and rhizosphere which is a place of microbial activity. Microbial activity in paddy soil can cause changes in soil biochemical functions such as; dissolution, binding, mineralization, immobilization, oxidation and reduction (Afzal and Singh, 2022).

Based on the explanation of the various theoretical concepts above, the purpose of this study was to determine the increase in the quality of bacteria and the production of rice plants.

## **2. Materials and Methods**

### **2.1. Place and Time of Implementation**

The research was conducted in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The types of rice and organic fertilizers used were Cakrabuana varieties and Granul fertilizers. This research was carried out on alluvial soil types which have a high organic matter content, brown to gray in color with slightly loose soil structure. According to Swetha *et al* (2022), alluvial soils fall into the Vertic Endoaquepts classification, with a sandy loam texture class (15% clay and 43% silt on the treated layer, 23% clay and 40% silt on the bottom layer). The soil in the tilled layer (0-20 cm) has a slightly acidic reaction (pH-H<sub>2</sub>O 5.6). Low total N content (0.3 mg/g), low C-organic (3.2 mg/g), low Bray 1 extracted P (5.06 ppm P), low CEC (6.96 cmol(+)/kg), cations K, Na, Ca, Low exchangeable Mg 0.12 each; 0.24; 3.05; and 0.61 cmol(+)/kg.

### **2.2. Research Implementation**

The study was arranged in a randomized block design with three replications and six treatments which included:

1. No fertilizer
2. NPK recommendation
3. Granular Fertilizer 10 tons/hectare
4. Granular Fertilizer 10 tons/hectare + recommended NPK
5. Granular Fertilizer 20 tons/hectare
6. Granular Fertilizer 20 tons/hectare + recommended NPK

The recommended dose of inorganic fertilizer for alluvial rice field soil in Gorontalo is 112.5 kg N; 45 kg P<sub>2</sub>O<sub>5</sub>; 60 kg K<sub>2</sub>O/hectare. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo.

The first stage of fertilizer application was carried out 10 days after planting for P fertilizer at once, 1/3 dose N, and ½ dose K. The second stage fertilizer application was carried out 21 days after planting for 1/3 dose N and ½ dose K, and the second stage was given the third is done 42 days after planting for 1/3 dose of N. Granular organic fertilizer is given simultaneously with soil processing.

### **2.3. Observed variable**

1. Bacterial quality, calculated using a kid bacterial test
2. Plant height, measured from 12 clump samples until just before harvest
3. The number of tillers, measured from 12 samples of clumps until just before harvest
4. Production of dry harvested grain and dry milled grain, measured by the size of the tiles (3m x 3m) m.

## **3. Results and Discussion**


### **3.1. Rice Plant Growth**

Plant growth depends on genetic characteristics and the ability to adapt to environmental conditions of cultivation (Jiang *et al*, 2022). Observations of plant height during the rice growth period were observed from

plants aged 14 days after planting until harvest. The growth of rice plant height from the age of 14 days after planting increased, the rice plant age increased, but after the plant was 94 days after planting, the rice plant height began to decrease, because the growth of rice grains had begun to bend and turn yellow. The best growth in rice plant height was in the treatment of granulated organic fertilizer 20 tons/hectare + NPK.


The increase in plant height growth is due to the addition of organic matter to the soil. The higher the dose of organic fertilizer used, the higher the plant growth, when compared to the control treatment. The ability of organic fertilizers to bind water and increase soil porosity will improve soil respiration, so that it can support root growth to help plant height growth. The high growth of rice plants is described in Figure 1.

**Image 1.** Growth of rice plant height at various plant ages and all treatments



The increase in rice plant height growth was in line with the growth in the number of tillers at various plant ages. The highest number of tillers was in the treatment of using granulated organic fertilizer 20 tons/hectare + NPK, followed by the treatment using granulated organic fertilizer 20 tons/hectare. Organic fertilizers contain lots of humic acid and folic acid as well as growth regulators which can accelerate the growth of rice tillers (Brambilla *et al*, 2022). According to Jiang *et al* (2023), the application of organic fertilizers is beneficial in providing and absorbing nutrients, as well as activating soil microorganisms, so that the soil structure becomes crumbly which causes an expansion of root reach and increases the number of rice tillers. The growth of the number of rice tillers is described in Figure 2.

**Figure 2.** The growth of the number of rice tillers at various ages of observation and all treatments



Provision of organic fertilizers can improve the quality of bacteria in the soil. Granul organic fertilizer dosage has a significant effect on improving the quality of bacteria in the soil. Soils that are not given organic matter of bacterial quality tend to be low, because the availability and activity of bacteria is needed in the process of absorbing and providing nutrients for plants. Organic fertilizers play an important role in the mineral cycle, especially the nitrogen, phosphorus, sulfur and carbon cycles (Cheng and Wan, 2023). Bacteria that play a role in the nitrogen cycle include Azotobacter and Azospirillum, these bacteria are non-symbiotic in nature

that are able to bind free N<sub>2</sub> which has an influence on the physical and chemical properties of the soil so as to increase soil fertility (Harindintwali *et al*, 2022).

The quality of the bacteria at the beginning of the treatment ranged from 10<sup>2</sup> to 10<sup>3</sup> and the quality of the bacteria continued to increase after being given treatment. The treatment of using granular organic fertilizer 20 tons hectare<sup>-1</sup> can improve the quality of bacteria from 10.1<sup>3</sup> cfu/ml to 10.1<sup>7</sup> cfu/ml. In this study it was seen that the more doses of organic fertilizer given to the soil, the higher the quality of the bacteria produced. According to Zainuddin *et al* (2022), the application of organic fertilizers in the long term increases the amount of nutrients such as available nitrogen, available phosphorus and total zinc in the soil and will increase the number and diversity of bacteria in the soil for plant growth and production processes. The increase in the quality of bacteria at the beginning and end of the study is described in Table 1.

**Table 1.** Bacterial quality at the beginning and end of the study

No	Treatment	Initial bacterial quality (cfu/ml)	Final bacterial quality (cfu/ml)
1	No fertilizer/control	10,2 <sup>2</sup>	10,2 <sup>2</sup>
2	NPK recommendations	10,2 <sup>2</sup>	10,2 <sup>2</sup>
3	Granular Fertilizer 10 tons/hectare	10,1 <sup>3</sup>	10,1 <sup>6</sup>
4	Granular Fertilizer 10 tons/hectare + NPK recommendation	10,1 <sup>3</sup>	10,1 <sup>6</sup>
5	Granular Fertilizer 20 tons/hectare	10,1 <sup>3</sup>	10,1 <sup>7</sup>
6	Granular Fertilizer 20 tons/hectare + NPK recommendation	10,1 <sup>3</sup>	10,1 <sup>7</sup>

### 3.2. Rice Production

The use of organic fertilizers can restore and increase soil fertility in addition to increasing crop production. Rice plant production through grain yield indicators is the role of organic fertilizer application which can facilitate the physiological processes of plant growth and production. The production results of rice plants are described in Figure 3.

**Figure 3.** Production of harvested dry unhulled rice and dry milled unhusked rice (tonnes/hectare)




Figure 3 explains that the highest rice production was found in the treatment of granular organic fertilizer 20 tons/hectare + NPK with a production of 7.4 tons/hectare harvested dry grain, and 4.24 tons/hectare dry milled grain. The weight production of harvested dry grain and dry milled grain in rice plants reflects the accumulation of organic compounds that plants have successfully synthesized from organic compounds, water and carbon dioxide which will contribute to plant dry weight (Alvarez *et al*, 2021). Active division of meristem cells in roots will spur good root development, so that active microbial activity in the root area and energy generated from the respiration process becomes more and supports plant canopy growth, thereby increasing plant dry weight (Olenska *et al*, 2020).

#### 4. Conclusion

Granular organic fertilizer application 20 tons/hectare can improve the quality of bacteria from  $10.1^3$  to  $10.1^7$  cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is a treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare., when compared to production in the organic fertilizer treatment Granul 20 ton/hectare, as well as the recommended NPK fertilizer treatment.

#### Compliance with ethical standards:

##### *Thank-you note*

The author would like to thank the Agriculture Office of Bone Bolango Regency, Gorontalo Province, Indonesia, for facilitating the implementation of this research.

##### *Disclosure of conflicts of interest*

The author declares no conflict of interest.

#### Reference

- [1] Milosevic, T., Milosevic, N. and Mladenovic, J. The influence of organic, organo-mineral and mineral fertilizers on tree growth, yielding, fruit quality and leaf nutrient composition of apple cv. 'Golden Delicious Renders'. Journal of Scientia Horticulturee 2022. 297:110 - 120.<https://doi.org/10.1016/j.scienta.2022.110978>.
- [2] Weerahewa, J. and Dayananda, D. Land use changes and economic effects of alternative fertilizer policies: A simulation analysis with a bio-economic model for a Tank Village of Sri Lanka. Journal of Agricultural Systems 2023. 205:563 - 514.<https://doi.org/10.1016/j.agry.2022.103563>.
- [3] Puech, T. and Stark, F. Diversification of an integrated crop-livestock system: Agroecological and food production assessment at farm scale. Journal of Agriculture, Ecosystems & Environment 2023. 344: 108 - 118.<https://doi.org/10.1016/j.agee.2022.108300>.
- [4] Chen, D., Wang, X., Carrion, VJ, Yin, S., Yue, Z., Liao, Y., Dong, Y. and Li, X. Acidic amelioration of soil amendments improves soil health by impacting rhizosphere microbial assemblies. Journal of Soil Biology and Biochemistry 2022. 167: 599 - 609.<https://doi.org/10.1016/j.soilbio.2022.108599>.
- [5] Hui, K., Tan, W. and Song, Q. Long-term application of nitrogen fertilizer alters the properties of dissolved soil organic matter and increases the accumulation of polycyclic aromatic hydrocarbons. Journal of Environmental Research 2022. 215: 114 - 125.<https://doi.org/10.1016/j.envres.2022.114267>.
- [6] Gao, Y., Shao, G., Yang, Z., Zhang, K., Lu, J., Wang, Z., Wu, S. and Xu, D. Influences of soil and biochar properties and amount of biochar and fertilizer on the performance of biochar in improving plant photosynthetic rate: A meta-analysis. European Journal of Agronomy 2021. 130: 126 - 138.<https://doi.org/10.1016/j.eja.2021.126345>.
- [7] Hameed, A., Chen, YP, Shen, FT, Lin, SY, Huang, HI, Lin, YW and Young, CC Evaluation of a subtropical maize-rice rotation system maintained under sustainable long-term fertilizer inputs for intensification of agriculture. Journal of Applied Soil Ecology 2023. 184: 104 - 114.<https://doi.org/10.1016/j.apsoil.2022.104772>.
- [8] Negi, R., Kaur, T., Devi, R., Kour, D. and Yadav, AN Assessment of nitrogen-fixing endophytic and mineral solubilizing rhizospheric bacteria as a multifunctional microbial consortium for growth promotion of wheat and wild wheat relative Aegilops kotschy. Journal Heliyon 2022. 8:125 - 135.<https://doi.org/10.1016/j.heliyon.2022.e12579>.



- [9] Allouzi, MMA, Allouzi, SMA, Keng, ZX, Supramaniam, CV, Singh, A. and Chong, S. Liquid biofertilizers as a sustainable solution for agriculture. *Journal Heliyon* 2022. 8:609 - 619.<https://doi.org/10.1016/j.heliyon.2022.e12609>.
- [10] Sahab, S., Suhani, I., Srivastava, V., Chauhan, PS, Singh, RP and Prasad, V. Potential risk assessment of soil salinity to agroecosystem sustainability: Current status and management strategies. *Journal of Science of The Total Environment* 2021. 764:144 - 154.<https://doi.org/10.1016/j.scitotenv.2020.144164>.
- [11] Afzal, S. and Singh, NK Effect of zinc and iron oxide nanoparticles on plant physiology, seed quality and microbial community structure in a rice-soil-microbial ecosystem. *Journal of Environmental Pollution* 2022. 314: 120 - 131.<https://doi.org/10.1016/j.envpol.2022.120224>.
- [12] Swetha, RK, Dasgupta, S., Chakraborty, S., Li, B., Weindorf, DC, Mancini, M., Silva, SHG, Ribeiro, BT, Curi, N. and Ray, DP Using Nix color sensor and Munsell soil color variables to classify contrasting soil types and predict soil organic carbon in Eastern India. *Journal of Computers and Electronics in Agriculture* 2022. 199: 107 - 117.<https://doi.org/10.1016/j.compag.2022.107192>.
- [13] Jiang, L., Wu, L., Wang, Y., Xu, Q., Xu, Z. and Chen, W. Research progress on the divergence and genetic basis of agronomic traits in Xian and Gang rice. *The Crop Journal* 2022. 10:924 - 931.<https://doi.org/10.1016/j.cj.2022.02.006>.
- [14] Brambilla, S., Stritzler, M., Soto, G. and Ayub, N. A synthesis of functional contributions of rhizobacteria to growth promotion in diverse crops. *Journal of the Rhizosphere* 2022. 24: 100 - 110.<https://doi.org/10.1016/j.rhisph.2022.100611>.
- [15] Jiang, Z., Li, Y., Cao, B., Li, J., Ren, Z., Qu, J. and Zhang, Y. Noval bio-organic fertilizer containing *Arthrobacter* sp. DNS10 alleviates atrazine-induced growth inhibition on soybean by improving atrazine removal and nitrogen accumulation. *Chemosphere Journal* 2023. 313: 137 - 147.<https://doi.org/10.1016/j.chemosphere.2022.137575>.
- [16] Cheng, Y. and Wan, W. Strong linkage between nutrient-cycling functional gene diversity and ecosystem multifunctionality during winter composting with pig manure and fallen leaves. *Journal Science of The Total Environment* 2023. 867:161 - 172.<https://doi.org/10.1016/j.scitotenv.2023.161529>.
- [17] Harindintwali, JD, Wang, F., Yang, W., Zhou, J., Muhoza, B., Mugabowindekwe, M. and Yu, X. Harnessing the power of cellulolytic nitrogen-fixing bacteria for biovalorization of lignocellulosic biomass. *Journal of Industrial Crops and Products* 2022. 186: 115 - 125.<https://doi.org/10.1016/j.indcrop.2022.115235>.
- [18] Zainuddin, N., Keni, MF, Ibrahim, SAS and Masri, MMM Effect of integrated biofertilizers with chemical fertilizers on oil palm growth and soil microbial diversity. *Journal of Biocatalysis and Agricultural Biotechnology* 2022. 39: 102 - 112.<https://doi.org/10.1016/j.bcab.2021.102237>.
- [19] Alvarez, AL, Weyers, SL, Goemann, HM, Peyton, BM and Gardner, RD Microalgae, soil and plants: A critical review of microalgae as a renewable resource for agriculture. *Journal of Algal Research* 2021. 54:102 - 112.<https://doi.org/10.1016/j.algal.2021.102200>.
- [20] Olenska, E., Malek, W., Wojcik, M., Swiecicka, I., Thijs, S. and Vangronsveld, J. Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review . *Journal of Science of The Total Environment* 2020. 743:140 - 150.<https://doi.org/10.1016/j.scitotenv.2020.140682>.

### 3. Bukti Konfirmasi Review dan Hasil Review Pertama Tanggal 28 Januari 2023

File Edit View History Bookmarks Tools Help

ULSRA-2023-0017 Article proof | X

https://mail.google.com/mail/u/0/#search/editor-ijlsra%40sciresjournals.com/FMfcgzGrcPHLtFttMMKzzzVFfsPFjtvk 90%

Getting Started Other Bookmarks

Gmail editor-ijlsra@sciresjournals.com

Mail Chat Spaces Meet

4 dari 15

IJLSRA-2023-0017 Article proof for final check and approval Eksternal Kotak Masuk x

International Journal of Life Science Research Archive <editor-ijlsra@sciresjournals.com> 28 Jan 2023, 14:01 ☆ ↶ ⋮  
kepada saya ▾

Inggris > Indonesia ▾ Terjemahkan pesan Nonaktifkan untuk: Inggris x

Dear Author,  
Greetings for the day...

We are grateful for your association with us. Please find herewith attached final edited version (as a word file) of your manuscript (ref no. IJLSRA-2023-0017) entitled ***"Organic Fertilizer Application to Improve Bacterial Quality and Rice Production"*** which is accepted for publication in International Journal of Life Science Research Archive (IJLSRA).

Download Author proof copy of your manuscript: <https://sciresjournals.com/ijlsra/sites/default/files/Attached%20Files/L...>  
Download Author query form: <https://sciresjournals.com/ijlsra/content/downloads>

**Special Note:**

1. Provide the name(s) of authors as First Name (Given name)-Middle Name- Last name (Family name/Surname) for each author and check that all names are accurately spelled. It is important from Indexing purpose.
2. Author should provide 4-6 keywords which is important for indexing.
3. Scientific names of plant/animal/microbes should be written in italics and follow nomenclature rules like *Genus species*. Check this throughout the manuscript.
4. While writing numerical results in table author should use decimal point instead of comma.
5. Mention the X-axis and Y-axis title (label) in the figure.
6. Provide appropriate acknowledgments in the manuscript.
7. Ensure that, all the references appearing in the text should be enlisted at the end in reference section and vice versa.

Without this we will not be able to process your manuscript further.

**We hereby request you that,**

1. Read the proof at least thrice, note the errors and make necessary corrections (in blue colored text) to make the manuscript error free.
2. Do not change the format of the word file, just check the spelling and other errors in the attached text part.
3. Use this proof for checking the typesetting, editing, completeness and correctness of the text, tables and figures.
4. Significant changes to the article will only be considered at this stage with permission from the Editor.
5. Make sure that you send final approved copy of article proof along with duly filled Author proof form within 3 days after receipt of this e-mail.
6. **Your manuscript will not be further processed for publication until and unless we receive your final approved article proof and Author query form.**
7. **It is responsibility to author to check each and every word and spelling including italic and normal formatting of specific names. No changes will be accepted after this. If any changes are expected after publication of manuscript, author will have to pay charges accordingly.**

We would appreciate to receive your corrected proof as a word file by e-mail within stipulated time period so that we can process your article quickly and efficiently.

Thank you for submitting your work to this journal.

If you need more information, please do not hesitate to contact us.

With Best Regards,  
Managing Editor  
International Journal of Life Science Research Archive (IJLSRA)  
[editor-ijlsra@sciresjournals.com](mailto:editor-ijlsra@sciresjournals.com)

Thank you for your mail. Approved. Thank you for your email.

Activate Windows  
Go to Settings to activate Windows.

Windows Taskbar: 8:49 AM 6/6/2023



(RESEARCH ARTICLE)



# Organic fertilizer application to improve bacterial quality and rice production

Zulzain Ilahude, Hayatiningsih Gubali, Mohamad Lihawa and Mohamad Ikbah Bahuha\*

*Department of Agrotechnology, Faculty of Agriculture, Universitas Negeri Gorontalo, Indonesia.*

International Journal of Life Science Research Archive, 2023, XX(XX), XXX-XXX

Publication history: Received on 10 December 2022; revised on 25 January 2023; accepted on 27 January 2023

[illegible]

## Abstract

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil. This study aims to determine the increase in the quality of bacteria and production of rice plants. The research was carried out in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The research using Completely Randomized Block Design with 6 treatments 3 replications. The plant used is rice Cakrabuana variety. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo. The plot size used is (10 x 5) m. The bacterial population was calculated using the SANICHECK kit bacterial test. The dose of Granul organic fertilizer that uses 10 tons and 20 tons/hectare. The research results show that Granule organic fertilizer 20 tons/hectare can improve the quality of bacteria from 10.13 to 10.17 cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is the best treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare.

**Keywords:** Organic Fertilizers; Bacterial Quality; Rice Production

## 1 Introduction

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil (Milosevic *et al.*, 2022). According to Weerahewa and Dayananda (2023), organic fertilizers can reduce environmental pollution and improve land quality in a sustainable manner. In principle, the organic farming system is a farming system that adheres to natural principles in building agroecosystem balance, so that it is beneficial to soil, water, air, plants and all living things, and is able to provide healthy food for human needs (Puech and Stark, 2023). ).

The addition of organic fertilizers to the soil, especially in soils with low organic matter, is an effort to ameliorate the soil so that the provision of plant nutrients can be more effective (Chen *et al*, 2022). In general, the application of organic fertilizers to the soil will improve the physical, chemical and biological properties of the soil (Hui *et al*, 2022). According to Gao *et al* (2021), in soils that lack organic matter and degraded soils, organic fertilizer is the main requirement for soil amelioration, so that the provision of nutrient inputs is more efficient and effective.

Technology that can increase fertilization efficiency and the sustainability of rice production systems is microbial fertilizer technology contained in organic fertilizers (Hameed *et al*, 2023). Supply of some of the N, P, K nutrients needed by rice plants can be carried out by rhizosphere bacteria, endophytic, green manure which have the ability to fix N<sub>2</sub> from the air, Phosphate solubilizing microbes can help mine Phosphate in the soil, so that it becomes available P for plants (Negi *et al*, 2022). Microbial groups as biological agents that are widely used in organic fertilizers include *Azotobacter* Sp, *Lactobacillus* Sp, *Alkaligenes* sp, *Rhizobium* Sp, *Pseudomonas* Sp and Endophytic Bacteria such as *Ovhrobactrum pseudogignonense*, *Azospirillum*, and *Penicillium* Sp (Allouzi *et al*, 2022).

\* Corresponding author: Mohamad Ikbah Bahua

According to Sahab *et al* (2021), in paddy fields waterlogging greatly affects the microbial metabolic processes of rice plants, such as; eukaryotic, aquatic weeds, heterotrophic microflora and fauna including roots and rhizosphere which is a place of microbial activity. Microbial activity in paddy soil can cause changes in soil biochemical functions such as; dissolution, binding, mineralization, immobilization, oxidation and reduction (Afzal and Singh, 2022).

Based on the explanation of the various theoretical concepts above, the purpose of this study was to determine the increase in the quality of bacteria and the production of rice plants.

## 2 Material and methods

### 2.1 Place and Time of Implementation

The research was conducted in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The types of rice and organic fertilizers used were Cakrabuana varieties and Granul fertilizers. This research was carried out on alluvial soil types which have a high organic matter content, brown to gray in color with slightly loose soil structure. According to Swetha *et al* (2022), alluvial soils fall into the Vertic Endoaquepts classification, with a sandy loam texture class (15% clay and 43% silt on the treated layer, 23% clay and 40% silt on the bottom layer). The soil in the tilled layer (0-20 cm) has a slightly acidic reaction (pH-H<sub>2</sub>O 5.6). Low total N content (0.3 mg/g), low C-organic (3.2 mg/g), low Bray 1 extracted P (5.06 ppm P), low CEC (6.96 cmol(+)/kg), cations K, Na, Ca, Low exchangeable Mg 0.12 each; 0.24; 3.05; and 0.61 cmol(+)/kg.

### 2.2 Research Implementation

The study was arranged in a randomized block design with three replications and six treatments which included:

- No fertilizer
- NPK recommendation
- Granular Fertilizer 10 tons/hectare
- Granular Fertilizer 10 tons/hectare + recommended NPK
- Granular Fertilizer 20 tons/hectare
- Granular Fertilizer 20 tons/hectare + recommended NPK

The recommended dose of inorganic fertilizer for alluvial rice field soil in Gorontalo is 112.5 kg N; 45 kg P<sub>2</sub>O<sub>5</sub>; 60 kg K<sub>2</sub>O/hectare. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo.

The first stage of fertilizer application was carried out 10 days after planting for P fertilizer at once, 1/3 dose N, and 1/2 dose K. The second stage fertilizer application was carried out 21 days after planting for 1/3 dose N and 1/2 dose K, and the second stage was given the third is done 42 days after planting for 1/3 dose of N. Granular organic fertilizer is given simultaneously with soil processing.

### 2.3 Observed variable


- Bacterial quality, calculated using a kid bacterial test
- Plant height, measured from 12 clump samples until just before harvest
- The number of tillers, measured from 12 samples of clumps until just before harvest
- Production of dry harvested grain and dry milled grain, measured by the size of the tiles (3m x 3m) m.

## 3 Results and Discussion

### 3.1 Rice Plant Growth


Plant growth depends on genetic characteristics and the ability to adapt to environmental conditions of cultivation (Jiang *et al*, 2022). Observations of plant height during the rice growth period were observed from plants aged 14 days after planting until harvest. The growth of rice plant height from the age of 14 days after planting increased, the rice plant age increased, but after the plant was 94 days after planting, the rice plant height began to decrease, because the growth of rice grains had begun to bend and turn yellow. The best growth in rice plant height was in the treatment of granulated organic fertilizer 20 tons/hectare + NPK.

The increase in plant height growth is due to the addition of organic matter to the soil. The higher the dose of organic fertilizer used, the higher the plant growth, when compared to the control treatment. The ability of organic fertilizers to bind water and increase soil porosity will improve soil respiration, so that it can support root growth to help plant height growth. The high growth of rice plants is described in Figure 1.



**Figure 1** Growth of rice plant height at various plant ages and all treatments

The increase in rice plant height growth was in line with the growth in the number of tillers at various plant ages. The highest number of tillers was in the treatment of using granulated organic fertilizer 20 tons/ hectare + NPK, followed by the treatment using granulated organic fertilizer 20 tons/hectare. Organic fertilizers contain lots of humic acid and folic acid as well as growth regulators which can accelerate the growth of rice tillers (Brambilla *et al*, 2022). According to Jiang *et al* (2023), the application of organic fertilizers is beneficial in providing and absorbing nutrients, as well as activating soil microorganisms, so that the soil structure becomes crumbly which causes an expansion of root reach and increases the number of rice tillers. The growth of the number of rice tillers is described in Figure 2.



**Figure 2** The growth of the number of rice tillers at various ages of observation and all treatments

Provision of organic fertilizers can improve the quality of bacteria in the soil. Granul organic fertilizer dosage has a significant effect on improving the quality of bacteria in the soil. Soils that are not given organic matter of bacterial quality tend to be low, because the availability and activity of bacteria is needed in the process of absorbing and providing nutrients for plants. Organic fertilizers play an important role in the mineral cycle, especially the nitrogen, phosphorus, sulfur and carbon cycles (Cheng and Wan, 2023). Bacteria that play a role in the nitrogen cycle include Azotobacter and Azospirillum, these bacteria are non-symbiotic in nature that are able to bind free N<sub>2</sub> which has an influence on the physical and chemical properties of the soil so as to increase soil fertility (Harindintwali *et al*, 2022).

The quality of the bacteria at the beginning of the treatment ranged from 10<sup>2</sup> to 10<sup>3</sup> and the quality of the bacteria continued to increase after being given treatment. The treatment of using granular organic fertilizer 20 tons hectare<sup>-1</sup> can improve the quality of bacteria from 10.1<sup>3</sup> cfu/ml to 10.1<sup>7</sup> cfu/ml. In this study it was seen that the more doses of organic fertilizer given to the soil, the higher the quality of the bacteria produced. According to Zainuddin *et al* (2022), the application of organic fertilizers in the long term increases the amount of nutrients such as available nitrogen, available phosphorus and total zinc in the soil and will increase the number and diversity of bacteria in the soil for plant growth and production processes. The increase in the quality of bacteria at the beginning and end of the study is described in Table 1.

**Table 1** Bacterial quality at the beginning and end of the study

No	Treatment	Initial bacterial quality (cfu/ml)	Final bacterial quality (cfu/ml)
1	No fertilizer/control	10,2 <sup>2</sup>	10,2 <sup>2</sup>
2	NPK recommendations	10,2 <sup>2</sup>	10,2 <sup>2</sup>
3	Granular Fertilizer 10 tons/hectare	10,1 <sup>3</sup>	10,1 <sup>6</sup>
4	Granular Fertilizer 10 tons/hectare + NPK recommendation	10,1 <sup>3</sup>	10,1 <sup>6</sup>
5	Granular Fertilizer 20 tons/hectare	10,1 <sup>3</sup>	10,1 <sup>7</sup>
6	Granular Fertilizer 20 tons/hectare + NPK recommendation	10,1 <sup>3</sup>	10,1 <sup>7</sup>

### 3.2 Rice Production

The use of organic fertilizers can restore and increase soil fertility in addition to increasing crop production. Rice plant production through grain yield indicators is the role of organic fertilizer application which can facilitate the physiological processes of plant growth and production. The production results of rice plants are described in Figure 3.


**Figure 3** Production of harvested dry unhulled rice and dry milled unhulled rice (tonnes/hectare)

Figure 3 explains that the highest rice production was found in the treatment of granular organic fertilizer 20 tons/hectare + NPK with a production of 7.4 tons/hectare harvested dry grain, and 4.24 tons/hectare dry milled grain. The weight production of harvested dry grain and dry milled grain in rice plants reflects the accumulation of organic compounds that plants have successfully synthesized from organic compounds, water and carbon dioxide which will contribute to plant dry weight (Alvarez *et al*, 2021). Active division of meristem cells in roots will spur good root development, so that active microbial activity in the root area and energy generated from the respiration process becomes more and supports plant canopy growth, thereby increasing plant dry weight (Olenska *et al*, 2020).

## 4 Conclusion

Granular organic fertilizer application 20 tons/hectare can improve the quality of bacteria from 10.1<sup>3</sup> to 10.1<sup>7</sup> cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is a treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare, when compared to production in the organic fertilizer treatment Granul 20 ton/hectare, as well as the recommended NPK fertilizer treatment.



---

## Compliance with ethical standards

### *Acknowledgments*

Acknowledgments must be inserted here.

### *Thank-you note*

The author would like to thank the Agriculture Office of Bone Bolango Regency, Gorontalo Province, Indonesia, for facilitating the implementation of this research.

### *Disclosure of conflict of interest*

The author declares no conflict of interest.

---

## References

- [1] Milosevic, T., Milosevic, N. and Mladenovic, J. The influence of organic, organo-mineral and mineral fertilizers on tree growth, yielding, fruit quality and leaf nutrient composition of apple cv. 'Golden Delicious Renders'. *Journal of Scientia Horticulture* 2022. 297:110 - 120.<https://doi.org/10.1016/j.scienta.2022.110978>.
- [2] Weerahewa, J. and Dayananda, D. Land use changes and economic effects of alternative fertilizer policies: A simulation analysis with a bio-economic model for a Tank Village of Sri Lanka. *Journal of Agricultural Systems* 2023. 205:563 - 514.<https://doi.org/10.1016/j.agsy.2022.103563>.
- [3] Puech, T. and Stark, F. Diversification of an integrated crop-livestock system: Agroecological and food production assessment at farm scale. *Journal of Agriculture, Ecosystems & Environment* 2023. 344: 108 - 118.<https://doi.org/10.1016/j.agee.2022.108300>.
- [4] Chen, D., Wang, X., Carrion, VJ, Yin, S., Yue, Z., Liao, Y., Dong, Y. and Li, X. Acidic amelioration of soil amendments improves soil health by impacting rhizosphere microbial assemblies. *Journal of Soil Biology and Biochemistry* 2022. 167: 599 - 609.<https://doi.org/10.1016/j.soilbio.2022.108599>.
- [5] Hui, K., Tan, W. and Song, Q. Long-term application of nitrogen fertilizer alters the properties of dissolved soil organic matter and increases the accumulation of polycyclic aromatic hydrocarbons. *Journal of Environmental Research* 2022. 215: 114 - 125.<https://doi.org/10.1016/j.envres.2022.114267>.
- [6] Gao, Y., Shao, G., Yang, Z., Zhang, K., Lu, J., Wang, Z., Wu, S. and Xu, D. Influences of soil and biochar properties and amount of biochar and fertilizer on the performance of biochar in improving plant photosynthetic rate: A meta-analysis. *European Journal of Agronomy* 2021. 130: 126 - 138.<https://doi.org/10.1016/j.eja.2021.126345>.
- [7] Hameed, A., Chen, YP, Shen, FT, Lin, SY, Huang, HI, Lin, YW and Young, CC Evaluation of a subtropical maize-rice rotation system maintained under sustainable long-term fertilizer inputs for intensification of agriculture. *Journal of Applied Soil Ecology* 2023. 184: 104 - 114.<https://doi.org/10.1016/j.apsoil.2022.104772>.
- [8] Negi, R., Kaur, T., Devi, R., Kour, D. and Yadav, AN Assessment of nitrogen-fixing endophytic and mineral solubilizing rhizospheric bacteria as a multifunctional microbial consortium for growth promotion of wheat and wild wheat relative *Aegilops kotschy*. *Journal Heliyon* 2022. 8:125 - 135.<https://doi.org/10.1016/j.heliyon.2022.e12579>.
- [9] Allouzi, MMA, Allouzi, SMA, Keng, ZX, Supramaniam, CV, Singh, A. and Chong, S. Liquid biofertilizers as a sustainable solution for agriculture. *Journal Heliyon* 2022. 8:609 - 619.<https://doi.org/10.1016/j.heliyon.2022.e12609>.
- [10] Sahab, S., Suhani, I., Srivastava, V., Chauhan, PS, Singh, RP and Prasad, V. Potential risk assessment of soil salinity to agroecosystem sustainability: Current status and management strategies. *Journal of Science of The Total Environment* 2021. 764:144 - 154.<https://doi.org/10.1016/j.scitotenv.2020.144164>.
- [11] Afzal, S. and Singh, NK Effect of zinc and iron oxide nanoparticles on plant physiology, seed quality and microbial community structure in a rice-soil-microbial ecosystem. *Journal of Environmental Pollution* 2022. 314: 120 - 131.<https://doi.org/10.1016/j.envpol.2022.120224>.
- [12] Swetha, RK, Dasgupta, S., Chakraborty, S., Li, B., Weindorf, DC, Mancini, M., Silva, SHG, Ribeiro, BT, Curi, N. and Ray, DP Using Nix color sensor and Munsell soil color variables to classify contrasting soil types and predict soil

- organic carbon in Eastern India. *Journal of Computers and Electronics in Agriculture* 2022. 199: 107 - 117.<https://doi.org/10.1016/j.compag.2022.107192>.
- [13] Jiang, L., Wu, L., Wang, Y., Xu, Q., Xu, Z. and Chen, W. Research progress on the divergence and genetic basis of agronomic traits in Xian and Gang rice. *The Crop Journal* 2022. 10:924 - 931.<https://doi.org/10.1016/j.cj.2022.02.006>.
- [14] Brambilla, S., Stritzler, M., Soto, G. and Ayub, N. A synthesis of functional contributions of rhizobacteria to growth promotion in diverse crops. *Journal of the Rhizosphere* 2022. 24: 100 - 110.<https://doi.org/10.1016/j.rhisph.2022.100611>.
- [15] Jiang, Z., Li, Y., Cao, B., Li, J., Ren, Z., Qu, J. and Zhang, Y. Noval bio-organic fertilizer containing *Arthrobacter* sp. DNS10 alleviates atrazine-induced growth inhibition on soybean by improving atrazine removal and nitrogen accumulation. *Chemosphere Journal* 2023. 313: 137 - 147.<https://doi.org/10.1016/j.chemosphere.2022.137575>.
- [16] Cheng, Y. and Wan, W. Strong linkage between nutrient-cycling functional gene diversity and ecosystem multifunctionality during winter composting with pig manure and fallen leaves. *Journal Science of The Total Environment* 2023. 867:161 - 172.<https://doi.org/10.1016/j.scitotenv.2023.161529>.
- [17] Harindintwali, JD, Wang, F., Yang, W., Zhou, J., Muhoza, B., Mugabowindekwe, M. and Yu, X. Harnessing the power of cellulolytic nitrogen-fixing bacteria for biovalorization of lignocellulosic biomass. *Journal of Industrial Crops and Products* 2022. 186: 115 - 125.<https://doi.org/10.1016/j.indcrop.2022.115235>.
- [18] Zainuddin, N., Keni, MF, Ibrahim, SAS and Masri, MMM Effect of integrated biofertilizers with chemical fertilizers on oil palm growth and soil microbial diversity. *Journal of Biocatalysis and Agricultural Biotechnology* 2022. 39: 102 - 112.<https://doi.org/10.1016/j.bcab.2021.102237>.
- [19] Alvarez, AL, Weyers, SL, Goemann, HM, Peyton, BM and Gardner, RD Microalgae, soil and plants: A critical review of microalgae as a renewable resource for agriculture. *Journal of Algal Research* 2021. 54:102 - 112.<https://doi.org/10.1016/j.algal.2021.102200>.
- [20] Olenska, E., Malek, W., Wojcik, M., Swiecicka, I., Thijs, S. and Vangronsveld, J. Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. *Journal of Science of The Total Environment* 2020. 743:140 - 150.<https://doi.org/10.1016/j.scitotenv.2020.140682>



#### 4. Bukti Konfirmasi Submit Revisi Pertama, Respon kepada Reviewer, dan Artikel Resubmit, Tanggal 30 Januari 2023

File Edit View History Bookmarks Tools Help

Results of article improvement | X +

https://mail.google.com/mail/u/0/#search/editor-ijlsra%40sciresjournals.com/KtbxLxGSpzZTmWPmZzRbfxn 90%

Getting Started Other Bookmarks

Gmail editor-ijlsra@sciresjournals.com X 90% Aktif ?

99+ Mail Chat Spaces Meet

3 dari 15 < >

**Mohamad Ikbal Bahua** <mohamad.bahua@ung.ac.id> kepada International 30 Jan 2023, 11:15 ☆ ↶ ⋮

Good afternoon  
Dear  
Managing Editor  
International Journal of Life Science Research Archive (IJLSRA)

Here We convey the results of the improvement of the article Manuscript number (ref no. IJLSRA-2023-0017) and a statement certificate about the article, according to the instructions of the editor.

Thanks  
Greeting  
Correspondent Author

Mohamad Ikbal Bahua  
[mohamad.bahua@ung.ac.id](mailto:mohamad.bahua@ung.ac.id)

5 Lampiran • Dipindai dengan Gmail

IJLSRA-2023-001... IJLSRA Authors D... Author Queries.d... cover letter IJLS... IJLSRA Authors n...

Activate Windows  
Go to Settings to activate Windows.

9:22 AM  
6/6/2023

## Organic fertilizer application to improve bacterial quality and rice production

Zulzain Ilahude, Hayatiningsih Gubali, Mohamad Lihawa and Mohamad Ikbah Bahua \*

*Department of Agrotechnology, Faculty of Agriculture, Universitas Negeri Gorontalo, Indonesia.*

International Journal of Life Science Research Archive, 2023, XX(XX), XXX-XXX

Publication history: Received on 10 December 2022; revised on 25 January 2023; accepted on 27 January 2023

Article DOI: xx

### Abstract

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil. This study aims to determine the increase in the quality of bacteria and production of rice plants (*Oryza sativa*). The research was carried out in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The research using Completely Randomized Block Design with 6 treatments 3 replications. The plant used is rice Cakrabuana variety. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo. The plot size used is (10 x 5) m. The bacterial population was calculated using the SANICHECK kit bacterial test. The dose of Granul organic fertilizer that uses 10 tons and 20 tons/hectare. The research results show that Granule organic fertilizer 20 tons/hectare can improve the quality of bacteria from 10.13 to 10.17 cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is the best treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare.

**Keywords:** *Bacterial quality; Organic fertilizers; Rice plant growth; Rice production*

### 1 Introduction

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil (Milosevic *et al.*, 2022). According to Weerahewa and Dayananda (2023), organic fertilizers can reduce environmental pollution and improve land quality in a sustainable manner. In principle, the organic farming system is a farming system that adheres to natural principles in building agro-ecosystem balance, so that it is beneficial to soil, water, air, plants and all living things, and is able to provide healthy food for human needs (Puech and Stark, 2023).

The addition of organic fertilizers to the soil, especially in soils with low organic matter, is an effort to ameliorate the soil so that the provision of plant nutrients can be more effective (Chen *et al.*, 2022). In general, the application of organic fertilizers to the soil will improve the physical, chemical and biological properties of the soil (Hui *et al.*, 2022). According to Gao *et al.* (2021), in soils that lack organic matter and degraded soils, organic fertilizer is the main requirement for soil amelioration, so that the provision of nutrient inputs is more efficient and effective.

Technology that can increase fertilization efficiency and the sustainability of rice production systems is microbial fertilizer technology contained in organic fertilizers (Hameed *et al.*, 2023). Supply of some of the N, P, K nutrients needed by rice plants can be carried out by rhizosphere bacteria, endophytic, green manure which have the ability to fix N<sub>2</sub> from the air, Phosphate solubilizing microbes can help mine Phosphate in the soil, so that it becomes available P for plants (Negi *et al.*, 2022). Microbial groups as biological agents that are widely used in organic fertilizers include *Azotobacter* sp, *Lactobacillus* sp, *Alkaligenes* sp, *Rhizobium* sp, *Pseudomonas* sp and Endophytic Bacteria such as *Ovhrbactrum pseudogignonense*, *Azospirillum* sp, and *Penicillium* sp (Allouzi *et al.*, 2022).

\* Corresponding author: Mohamad Ikbah Bahua

According to Sahab *et al* (2021), in paddy fields waterlogging greatly affects the microbial metabolic processes of rice plants, such as; eukaryotic, aquatic weeds, heterotrophic microflora and fauna including roots and rhizosphere which is a place of microbial activity. Microbial activity in paddy soil can cause changes in soil biochemical functions such as; dissolution, binding, mineralization, immobilization, oxidation and reduction (Afzal and Singh, 2022).

Based on the explanation of the various theoretical concepts above, the purpose of this study was to determine the increase in the quality of bacteria and the production of rice plants.

## 2 Material and methods

### 2.1 Place and Time of Implementation

The research was conducted in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The types of rice and organic fertilizers used were Cakrabuana varieties and Granul fertilizers. This research was carried out on alluvial soil types which have a high organic matter content, brown to gray in color with slightly loose soil structure. According to Swetha *et al* (2022), alluvial soils fall into the Vertic Endoaquepts classification, with a sandy loam texture class (15% clay and 43% silt on the treated layer, 23% clay and 40% silt on the bottom layer). The soil in the tilled layer (0-20 cm) has a slightly acidic reaction (pH-H<sub>2</sub>O 5.6). Low total N content (0.3 mg/g), low C-organic (3.2 mg/g), low Bray 1 extracted P (5.06 ppm P), low CEC (6.96 cmol(+)/kg), cations K, Na, Ca, Low exchangeable Mg 0.12 each; 0.24; 3.05; and 0.61 cmol(+)/kg.

### 2.2 Research Implementation

The study was arranged in a randomized block design with three replications and six treatments which included:

- No fertilizer
- NPK recommendation
- Granular Fertilizer 10 tons/hectare
- Granular Fertilizer 10 tons/hectare + recommended NPK
- Granular Fertilizer 20 tons/hectare
- Granular Fertilizer 20 tons/hectare + recommended NPK

The recommended dose of inorganic fertilizer for alluvial rice field soil in Gorontalo is 112.5 kg N; 45 kg P<sub>2</sub>O<sub>5</sub>; 60 kg K<sub>2</sub>O/hectare. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo.

The first stage of fertilizer application was carried out 10 days after planting for P fertilizer at once, 1/3 dose N, and 1/2 dose K. The second stage fertilizer application was carried out 21 days after planting for 1/3 dose N and 1/2 dose K, and the second stage was given the third is done 42 days after planting for 1/3 dose of N. Granular organic fertilizer is given simultaneously with soil processing.

### 2.3 Observed variable


- Bacterial quality, calculated using a kid bacterial test
- Plant height, measured from 12 clump samples until just before harvest
- The number of tillers, measured from 12 samples of clumps until just before harvest
- Production of dry harvested grain and dry milled grain, measured by the size of the tiles (3m x 3m) m.

## 3 Results and Discussion

### 3.1 Rice Plant Growth


Plant growth depends on genetic characteristics and the ability to adapt to environmental conditions of cultivation (Jiang *et al*, 2022). Observations of plant height during the rice growth period were observed from plants aged 14 days after planting until harvest. The growth of rice plant height from the age of 14 days after planting increased, the rice plant age increased, but after the plant was 94 days after planting, the rice plant height began to decrease, because the growth of rice grains had begun to bend and turn yellow. The best growth in rice plant height was in the treatment of granulated organic fertilizer 20 tons/hectare + NPK.

The increase in plant height growth is due to the addition of organic matter to the soil. The higher the dose of organic fertilizer used, the higher the plant growth, when compared to the control treatment. The ability of organic fertilizers to bind water and increase soil porosity will improve soil respiration, so that it can support root growth to help plant height growth. The high growth of rice plants is described in Figure 1.



**Figure 1** Growth of rice plant height at various plant ages and all treatments

The increase in rice plant height growth was in line with the growth in the number of tillers at various plant ages. The highest number of tillers was in the treatment of using granulated organic fertilizer 20 tons/ hectare + NPK, followed by the treatment using granulated organic fertilizer 20 tons/hectare. Organic fertilizers contain lots of humic acid and folic acid as well as growth regulators which can accelerate the growth of rice tillers (Brambilla *et al*, 2022). According to Jiang *et al* (2023), the application of organic fertilizers is beneficial in providing and absorbing nutrients, as well as activating soil microorganisms, so that the soil structure becomes crumbly which causes an expansion of root reach and increases the number of rice tillers. The growth of the number of rice tillers is described in Figure 2.



**Figure 2** The growth of the number of rice tillers at various ages of observation and all treatments

Provision of organic fertilizers can improve the quality of bacteria in the soil. Granul organic fertilizer dosage has a significant effect on improving the quality of bacteria in the soil. Soils that are not given organic matter of bacterial quality tend to be low, because the availability and activity of bacteria is needed in the process of absorbing and providing nutrients for plants. Organic fertilizers play an important role in the mineral cycle, especially the nitrogen, phosphorus, sulfur and carbon cycles (Cheng and Wan, 2023). Bacteria that play a role in the nitrogen cycle include Azotobacter and Azospirillum, these bacteria are non-symbiotic in nature that are able to bind free N<sub>2</sub> which has an influence on the physical and chemical properties of the soil so as to increase soil fertility (Harindintwali *et al*, 2022).


The quality of the bacteria at the beginning of the treatment ranged from  $10^2$  to  $10^3$  and the quality of the bacteria continued to increase after being given treatment. The treatment of using granular organic fertilizer 20 tons hectare<sup>-1</sup> can improve the quality of bacteria from  $10.1^3$  cfu/ml to  $10.1^7$  cfu/ml. In this study it was seen that the more doses of organic fertilizer given to the soil, the higher the quality of the bacteria produced. According to Zainuddin *et al* (2022), the application of organic fertilizers in the long term increases the amount of nutrients such as available nitrogen, available phosphorus and total zinc in the soil and will increase the number and diversity of bacteria in the soil for plant growth and production processes. The increase in the quality of bacteria at the beginning and end of the study is described in Table 1.

**Table 1** Bacterial quality at the beginning and end of the study

No	Treatment	Initial bacterial quality (cfu/ml)	Final bacterial quality (cfu/ml)
1	No fertilizer/control	$10.2^2$	$10.2^2$
2	NPK recommendations	$10.2^2$	$10.2^2$
3	Granular Fertilizer 10 tons/hectare	$10.1^3$	$10.1^6$
4	Granular Fertilizer 10 tons/hectare + NPK recommendation	$10.1^3$	$10.1^6$
5	Granular Fertilizer 20 tons/hectare	$10.1^3$	$10.1^7$
6	Granular Fertilizer 20 tons/hectare + NPK recommendation	$10.1^3$	$10.1^7$

### 3.2 Rice Production

The use of organic fertilizers can restore and increase soil fertility in addition to increasing crop production. Rice plant production through grain yield indicators is the role of organic fertilizer application which can facilitate the physiological processes of plant growth and production. The production results of rice plants are described in Figure 3.



**Figure 3** Production of harvested dry unhulled rice and dry milled unhusked rice (tonnes/hectare)

Figure 3 explains that the highest rice production was found in the treatment of granular organic fertilizer 20 tons/hectare + NPK with a production of 7.4 tons/hectare harvested dry grain, and 4.24 tons/hectare dry milled grain. The weight production of harvested dry grain and dry milled grain in rice plants reflects the accumulation of organic compounds that plants have successfully synthesized from organic compounds, water and carbon dioxide which will contribute to plant dry weight (Alvarez *et al*, 2021). Active division of meristem cells in roots will spur good root development, so that active microbial activity in the root area and energy generated from the respiration process becomes more and supports plant canopy growth, thereby increasing plant dry weight (Olenska *et al*, 2020).

## 4 Conclusion

Granular organic fertilizer application 20 tons/hectare can improve the quality of bacteria from  $10.1^3$  to  $10.1^7$  cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is a treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare., when compared to production in the organic fertilizer treatment Granul 20 ton/hectare, as well as the recommended NPK fertilizer treatment.

## Compliance with ethical standards

### *Acknowledgments*

The author would like to thank the Agriculture Office of Bone Bolango Regency, Gorontalo Province, Indonesia, for facilitating the implementation of this research.

### *Disclosure of conflict of interest*

The author declares no conflict of interest.

## References

- [1] Milosevic, T., Milosevic, N. and Mladenovic, J. The influence of organic, organo-mineral and mineral fertilizers on tree growth, yielding, fruit quality and leaf nutrient composition of apple cv. 'Golden Delicious Renders'. *Journal of Scientia Horticulture* 2022. 297:110 - 120.<https://doi.org/10.1016/j.scienta.2022.110978>.
- [2] Weerahewa, J. and Dayananda, D. Land use changes and economic effects of alternative fertilizer policies: A simulation analysis with a bio-economic model for a Tank Village of Sri Lanka. *Journal of Agricultural Systems* 2023. 205:563 - 514.<https://doi.org/10.1016/j.agsy.2022.103563>.
- [3] Puech, T. and Stark, F. Diversification of an integrated crop-livestock system: Agroecological and food production assessment at farm scale. *Journal of Agriculture, Ecosystems & Environment* 2023. 344: 108 - 118.<https://doi.org/10.1016/j.agee.2022.108300>.
- [4] Chen, D., Wang, X., Carrion, VJ, Yin, S., Yue, Z., Liao, Y., Dong, Y. and Li, X. Acidic amelioration of soil amendments improves soil health by impacting rhizosphere microbial assemblies. *Journal of Soil Biology and Biochemistry* 2022. 167: 599 - 609.<https://doi.org/10.1016/j.soilbio.2022.108599>.
- [5] Hui, K., Tan, W. and Song, Q. Long-term application of nitrogen fertilizer alters the properties of dissolved soil organic matter and increases the accumulation of polycyclic aromatic hydrocarbons. *Journal of Environmental Research* 2022. 215: 114 - 125.<https://doi.org/10.1016/j.envres.2022.114267>.
- [6] Gao, Y., Shao, G., Yang, Z., Zhang, K., Lu, J., Wang, Z., Wu, S. and Xu, D. Influences of soil and biochar properties and amount of biochar and fertilizer on the performance of biochar in improving plant photosynthetic rate: A meta-analysis. *European Journal of Agronomy* 2021. 130: 126 - 138.<https://doi.org/10.1016/j.eja.2021.126345>.
- [7] Hameed, A., Chen, YP, Shen, FT, Lin, SY, Huang, HI, Lin, YW and Young, CC Evaluation of a subtropical maize-rice rotation system maintained under sustainable long-term fertilizer inputs for intensification of agriculture. *Journal of Applied Soil Ecology* 2023. 184: 104 - 114.<https://doi.org/10.1016/j.apsoil.2022.104772>.
- [8] Negi, R., Kaur, T., Devi, R., Kour, D. and Yadav, AN Assessment of nitrogen-fixing endophytic and mineral solubilizing rhizospheric bacteria as a multifunctional microbial consortium for growth promotion of wheat and wild wheat relative *Aegilops kotschy*. *Journal Heliyon* 2022. 8:125 - 135.<https://doi.org/10.1016/j.heliyon.2022.e12579>.
- [9] Allouzi, MMA, Allouzi, SMA, Keng, ZX, Supramaniam, CV, Singh, A. and Chong, S. Liquid biofertilizers as a sustainable solution for agriculture. *Journal Heliyon* 2022. 8:609 - 619.<https://doi.org/10.1016/j.heliyon.2022.e12609>.
- [10] Sahab, S., Suhani, I., Srivastava, V., Chauhan, PS, Singh, RP and Prasad, V. Potential risk assessment of soil salinity to agroecosystem sustainability: Current status and management strategies. *Journal of Science of The Total Environment* 2021. 764:144 - 154.<https://doi.org/10.1016/j.scitotenv.2020.144164>.

- [11] Afzal, S. and Singh, NK Effect of zinc and iron oxide nanoparticles on plant physiology, seed quality and microbial community structure in a rice-soil-microbial ecosystem. *Journal of Environmental Pollution* 2022. 314: 120 - 131.<https://doi.org/10.1016/j.envpol.2022.120224>.
- [12] Swetha, RK, Dasgupta, S., Chakraborty, S., Li, B., Weindorf, DC, Mancini, M., Silva, SHG, Ribeiro, BT, Curi, N. and Ray, DP Using Nix color sensor and Munsell soil color variables to classify contrasting soil types and predict soil organic carbon in Eastern India. *Journal of Computers and Electronics in Agriculture* 2022. 199: 107 - 117.<https://doi.org/10.1016/j.compag.2022.107192>.
- [13] Jiang, L., Wu, L., Wang, Y., Xu, Q., Xu, Z. and Chen, W. Research progress on the divergence and genetic basis of agronomic traits in Xian and Gang rice. *The Crop Journal* 2022. 10:924 - 931.<https://doi.org/10.1016/j.cj.2022.02.006>.
- [14] Brambilla, S., Stritzler, M., Soto, G. and Ayub, N. A synthesis of functional contributions of rhizobacteria to growth promotion in diverse crops. *Journal of the Rhizosphere* 2022. 24: 100 - 110.<https://doi.org/10.1016/j.rhisph.2022.100611>.
- [15] Jiang, Z., Li, Y., Cao, B., Li, J., Ren, Z., Qu, J. and Zhang, Y. Noval bio-organic fertilizer containing *Arthrobacter* sp. DNS10 alleviates atrazine-induced growth inhibition on soybean by improving atrazine removal and nitrogen accumulation. *Chemosphere Journal* 2023. 313: 137 - 147.<https://doi.org/10.1016/j.chemosphere.2022.137575>.
- [16] Cheng, Y. and Wan, W. Strong linkage between nutrient-cycling functional gene diversity and ecosystem multifunctionality during winter composting with pig manure and fallen leaves. *Journal Science of The Total Environment* 2023. 867:161 - 172.<https://doi.org/10.1016/j.scitotenv.2023.161529>.
- [17] Harindintwali, JD, Wang, F., Yang, W., Zhou, J., Muhoza, B., Mugabowindekwe, M. and Yu, X. Harnessing the power of cellulolytic nitrogen-fixing bacteria for biovalorization of lignocellulosic biomass. *Journal of Industrial Crops and Products* 2022. 186: 115 - 125.<https://doi.org/10.1016/j.indcrop.2022.115235>.
- [18] Zainuddin, N., Keni, MF, Ibrahim, SAS and Masri, MMM Effect of integrated biofertilizers with chemical fertilizers on oil palm growth and soil microbial diversity. *Journal of Biocatalysis and Agricultural Biotechnology* 2022. 39: 102 - 112.<https://doi.org/10.1016/j.bcab.2021.102237>.
- [19] Alvarez, AL, Weyers, SL, Goemann, HM, Peyton, BM and Gardner, RD Microalgae, soil and plants: A critical review of microalgae as a renewable resource for agriculture. *Journal of Algal Research* 2021. 54:102 - 112.<https://doi.org/10.1016/j.algal.2021.102200>.
- [20] Olenska, E., Malek, W., Wojcik, M., Swiecicka, I., Thijs, S. and Vangronsveld, J. Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. *Journal of Science of The Total Environment* 2020. 743:140 - 150.<https://doi.org/10.1016/j.scitotenv.2020.140682>

Author's Declaration Form

**Title of article:** Organic Fertilizer Application to Improve Bacterial Quality and Rice Production

**Corresponding Author's name:** Mohamad Ikbah Bahua

**Names of co-authors:** Zulzain Ilahude (Author), Hayatiningsih Gubali (Co-Author),  
Mohamad Lihawa (Co-Author), Mohamad Ikbah Bahua (Co-Author).

**E-mail of corresponding author.** [mohamad.bahua@ung.ac.id](mailto:mohamad.bahua@ung.ac.id)

**Manuscript number:** IJLSRA-2023-0017

The author (s) acknowledged that:

1. All authors have read, understand and agreed to the submission guidelines, policies and submission declaration of the journal.
2. The manuscript is the authors' original work.
3. The manuscript has not been published and is not being submitted or considered for publication elsewhere;
4. All authors have seen and approved the manuscript as submitted;
5. All authors participated in the work in a substantive way and are prepared to take public responsibility for the work.
6. All authors of the manuscript have no conflict of interests to declare.
7. The manuscript submitted to the journal is not copied or plagiarized version of some other published work. All the data taken from other sources is written in authors own language and properly cited.
8. The text, illustrations, and any other materials included in the manuscript do not infringe upon any existing copyright or other rights of anyone.
9. Submission of false or incorrect information/undertaking would invite appropriate penal actions as per norms/rules of the journal.
10. All authors shall not submit the paper for publication in any other Journal or Magazine till the decision is made by journal editors.
11. All authors confirm that upon acceptance of the paper, they will publish the paper by paying Article Processing Charges (APC), or withdraw it by paying Withdrawal Fee.



Signature

**Date:** Januari 30, 2023

**Place:** Gorontalo, Indonesia

(Mohamad Ikbah Bahua)  
Corresponding Author



## Author Queries

Manuscript Number: IJLSRA-2023-0017

Queries	Details Required	Author's Response
AQ1	Please check and confirm the type of the article.	√
AQ2	Please check and confirm the title of the article.	√
AQ3	Please check and confirm the spellings of all the author names, their sequence, affiliations, corresponding author is correctly indicated	√
AQ4	Please check and confirm that all the keywords given by author are mentioned and correctly spelled.	√
AQ5	Please check whether all heading and subheading levels are okay.	√
AQ6	Please check and confirm that all spellings and grammar in the entire text are correctly used.	√
AQ7	Please check and confirm that all the punctuation marks are appropriately given.	√
AQ8	Please check and confirm that all the scientific nomenclature, units, symbols and abbreviations are correctly presented	√
AQ9	Please check and confirm that all Tables and Figures are included in the manuscript.	√
AQ10	Please check and confirm that all the caption of Tables and Figures are correctly presented.	√
AQ11	Please check and confirm that all footnotes of tables are okay.	√
AQ12	Please check and confirm that all the formulae (if present) are correctly presented.	√
AQ13	Please check and confirm that statements about 'Compliance with Ethical Standards' and correctly presented before reference section. These include (i) Acknowledgement (compulsory), (ii) Disclosure of conflict of interest (compulsory), (iii) Statement of ethical approval (if any) and (iv) Informed consent (if any).	√
AQ14	Please check and confirm that all the References cited in the text are given in the reference list and vice versa. Please provide details in the list or delete the citation from the text if applicable	√

Author should check all above queries. If okay then mention "Checked and Confirmed" in the Author's Response column; otherwise, do the necessary changes in the galley proof (in blue color text) and mention "Checked and Corrected" in the Author's Response column.

## COVER LETTER

Date: January 30, 2023

To,  
Editor-in-Chief,  
International Journal of Life Science Research Archive (IJLSRA)  
Subject: submission of new manuscript for evaluation


Dear Sir/Madam,

I am enclosing herewith a manuscript entitled **“Organic Fertilizer Application to Improve Bacterial Quality and Rice Production”** for possible evaluation as a **(Research article/)** in International Journal of Life Science Research Archive (IJLSRA).

**Research is useful for scientific information in the field of agriculture, especially food crops in improving knowledge and skills for the public and practitioners of agricultural researchers.**

All of the authors declare that they have all participated in the design, execution, and analysis of the paper, and that they have approved the final version. Additionally, there are no conflicts of interest in connection with this paper, and the material described is not under publication or consideration for publication elsewhere.

Thank you,



Sincerely,

Mohamad Ikbah Bahua

Mailing address

E-mail: [mohamad.bahua@ung.ac.id](mailto:mohamad.bahua@ung.ac.id)

Contact phone number: +6285240795645

Fax number



# International Journal of Life Science Research Archive (IJLSRA)

## Author's name details

Please provide below details


**Manuscript ref. No.: IJLSRA-2023-0017**

**Title of the manuscript: "Organic Fertilizer Application to Improve Bacterial Quality and Rice Production"**

### All author's name in detail

Sr. No.	Initial/First name	Middle name	Surname/Last name	Email-Id
Mr. 1	ZI/Zulzain	Ilahude	Zulzain Ilahude	<a href="mailto:zulzainilahude@ung.ac.id">zulzainilahude@ung.ac.id</a>
Ms. 2	HG/Hayatiningsih	Gubali	Hayatiningsih Gubali	<a href="mailto:hayatiningsihgubali@ung.ac.id">hayatiningsihgubali@ung.ac.id</a>
Mr. 3	ML/Mohamad	Lihawa	Mohamad Lihawa	<a href="mailto:mohamadlihawa@ung.ac.id">mohamadlihawa@ung.ac.id</a>
Mr. 4	MIB/Mohamad	Ikbal Bahua	Mohamad Ikbal Bahua	<a href="mailto:mohamad.bahua@ung.ac.id">mohamad.bahua@ung.ac.id</a>

## 5. Bukti Konfirmasi Konfirmasi Review Tanggal 30 Januari 2023



## 6. Bukti Konfirmasi Artikel Published Online Tanggal 2 Februari 2023

The screenshot shows a Gmail interface on a Windows 10 desktop. The email is from 'International Journal of Life Science Research Archive' to 'editor-ijlsra@sciresjournals.com'. The subject line is 'Your Manuscript (IJLSRA-2023-0017) has been published in Volume 4 - Issue 1 (January - March 2023) of International Journal of Life Science Research Archive (IJLSRA)'. The email body contains the following text:

Dear Author,  
Greetings for the day...

We are pleased to inform you that your manuscript entitled **"Organic Fertilizer Application to Improve Bacterial Quality and Rice Production"** has been published in **Volume 4 - Issue 1 (January - March 2023)** of International Journal of Life Science Research Archive (IJLSRA).

**You can view or download your article from the below links**

<https://sciresjournals.com/ijlsra/content/organic-fertilizer-application...>  
<https://sciresjournals.com/ijlsra/sites/default/files/IJLSRA-2023-0017.pdf>

Crossref DOI and CrossMark metadata information will be updated very soon. We will inform you regarding this.

I hope you are satisfied with our service and have good experience while publishing with us. We will be happy to review and publish your next manuscript in this journal. You may also consider our other allied journals...

**International Journal of Biology and Pharmacy Research Archive (IJBPR)**  
**International Journal of Science and Technology Research Archive (IJSTRA)**  
**International Journal of Biological and Pharmaceutical Sciences Archive (IJBPSA)**  
**International Journal of Chemistry and Pharmacy Research Archive (IJCPRA)**

At the bottom right of the email, there is a watermark that says 'Activate Windows Go to Settings to activate Windows.'

## Organic fertilizer application to improve bacterial quality and rice production

Zulzain Ilahude, Hayatiningsih Gubali, Mohamad Lihawa and Mohamad Ikbah Bahua \*

*Department of Agrotechnology, Faculty of Agriculture, Universitas Negeri Gorontalo, Indonesia.*

International Journal of Life Science Research Archive, 2023, 04(01), 117–122

Publication history: Received on 10 December 2022; revised on 25 January 2023; accepted on 27 January 2023

Article DOI: <https://doi.org/10.53771/ijlsra.2023.4.1.0017>

### Abstract

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil. This study aims to determine the increase in the quality of bacteria and production of rice plants (*Oryza sativa*). The research was carried out in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The research using Completely Randomized Block Design with 6 treatments 3 replications. The plant used is rice Cakrabuana variety. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo. The plot size used is (10 x 5) m. The bacterial population was calculated using the SANICHECK kit bacterial test. The dose of Granul organic fertilizer that uses 10 tons and 20 tons/hectare. The research results show that Granule organic fertilizer 20 tons/hectare can improve the quality of bacteria from 10.13 to 10.17 cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is the best treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare.

**Keywords:** Bacterial quality; Organic fertilizers; Rice plant growth; Rice production

### 1 Introduction

Organic fertilizers are very useful for increasing the nutrient content needed by plants, increasing plant productivity, stimulating the growth of plant organs and fertilizing the soil (Milosevic *et al.*, 2022). According to Weerahewa and Dayananda (2023), organic fertilizers can reduce environmental pollution and improve land quality in a sustainable manner. In principle, the organic farming system is a farming system that adheres to natural principles in building agro-ecosystem balance, so that it is beneficial to soil, water, air, plants and all living things, and is able to provide healthy food for human needs (Puech and Stark, 2023).

The addition of organic fertilizers to the soil, especially in soils with low organic matter, is an effort to ameliorate the soil so that the provision of plant nutrients can be more effective (Chen *et al.*, 2022). In general, the application of organic fertilizers to the soil will improve the physical, chemical and biological properties of the soil (Hui *et al.*, 2022). According to Gao *et al.* (2021), in soils that lack organic matter and degraded soils, organic fertilizer is the main requirement for soil amelioration, so that the provision of nutrient inputs is more efficient and effective.

Technology that can increase fertilization efficiency and the sustainability of rice production systems is microbial fertilizer technology contained in organic fertilizers (Hameed *et al.*, 2023). Supply of some of the N, P, K nutrients needed by rice plants can be carried out by rhizosphere bacteria, endophytic, green manure which have the ability to fix N<sub>2</sub> from the air, Phosphate solubilizing microbes can help mine Phosphate in the soil, so that it becomes available P for plants (Negi *et al.*, 2022). Microbial groups as biological agents that are widely used in organic fertilizers include *Azotobacter* sp, *Lactobacillus* sp, *Alkaligenes* sp, *Rhizobium* sp, *Pseudomonas* sp and Endophytic Bacteria such as *Ovrobactrum pseudogignonense*, *Azospirillum* sp, and *Penicillium* sp (Allouzi *et al.*, 2022).

\* Corresponding author: Mohamad Ikbah Bahua

According to Sahab *et al* (2021), in paddy fields waterlogging greatly affects the microbial metabolic processes of rice plants, such as; eukaryotic, aquatic weeds, heterotrophic microflora and fauna including roots and rhizosphere which is a place of microbial activity. Microbial activity in paddy soil can cause changes in soil biochemical functions such as; dissolution, binding, mineralization, immobilization, oxidation and reduction (Afzal and Singh, 2022).

Based on the explanation of the various theoretical concepts above, the purpose of this study was to determine the increase in the quality of bacteria and the production of rice plants.

## 2 Material and methods

### 2.1 Place and Time of Implementation

The research was conducted in paddy fields belonging to the Agriculture Service of Bone Bolango Regency, Gorontalo Province, Indonesia, from September to December 2022. The types of rice and organic fertilizers used were Cakrabuana varieties and Granul fertilizers. This research was carried out on alluvial soil types which have a high organic matter content, brown to gray in color with slightly loose soil structure. According to Swetha *et al* (2022), alluvial soils fall into the Vertic Endoaquepts classification, with a sandy loam texture class (15% clay and 43% silt on the treated layer, 23% clay and 40% silt on the bottom layer). The soil in the tilled layer (0-20 cm) has a slightly acidic reaction (pH-H<sub>2</sub>O 5.6). Low total N content (0.3 mg/g), low C-organic (3.2 mg/g), low Bray 1 extracted P (5.06 ppm P), low CEC (6.96 cmol(+)/kg), cations K, Na, Ca, Low exchangeable Mg 0.12 each; 0.24; 3.05; and 0.61 cmol(+)/kg.

### 2.2 Research Implementation

The study was arranged in a randomized block design with three replications and six treatments which included:

- No fertilizer
- NPK recommendation
- Granular Fertilizer 10 tons/hectare
- Granular Fertilizer 10 tons/hectare + recommended NPK
- Granular Fertilizer 20 tons/hectare
- Granular Fertilizer 20 tons/hectare + recommended NPK

The recommended dose of inorganic fertilizer for alluvial rice field soil in Gorontalo is 112.5 kg N; 45 kg P<sub>2</sub>O<sub>5</sub>; 60 kg K<sub>2</sub>O/hectare. Plant spacing using the Jajar Legowo system 4: 1 with a spacing of 15 cm x 22 cm and a row width of 40 cm Legowo.

The first stage of fertilizer application was carried out 10 days after planting for P fertilizer at once, 1/3 dose N, and 1/2 dose K. The second stage fertilizer application was carried out 21 days after planting for 1/3 dose N and 1/2 dose K, and the second stage was given the third is done 42 days after planting for 1/3 dose of N. Granular organic fertilizer is given simultaneously with soil processing.

### 2.3 Observed variable


- Bacterial quality, calculated using a kid bacterial test
- Plant height, measured from 12 clump samples until just before harvest
- The number of tillers, measured from 12 samples of clumps until just before harvest
- Production of dry harvested grain and dry milled grain, measured by the size of the tiles (3m x 3m) m.

## 3 Results and Discussion

### 3.1 Rice Plant Growth


Plant growth depends on genetic characteristics and the ability to adapt to environmental conditions of cultivation (Jiang *et al*, 2022). Observations of plant height during the rice growth period were observed from plants aged 14 days after planting until harvest. The growth of rice plant height from the age of 14 days after planting increased, the rice plant age increased, but after the plant was 94 days after planting, the rice plant height began to decrease, because the growth of rice grains had begun to bend and turn yellow. The best growth in rice plant height was in the treatment of granulated organic fertilizer 20 tons/hectare + NPK.

The increase in plant height growth is due to the addition of organic matter to the soil. The higher the dose of organic fertilizer used, the higher the plant growth, when compared to the control treatment. The ability of organic fertilizers to bind water and increase soil porosity will improve soil respiration, so that it can support root growth to help plant height growth. The high growth of rice plants is described in Figure 1.



**Figure 1** Growth of rice plant height at various plant ages and all treatments

The increase in rice plant height growth was in line with the growth in the number of tillers at various plant ages. The highest number of tillers was in the treatment of using granulated organic fertilizer 20 tons/ hectare + NPK, followed by the treatment using granulated organic fertilizer 20 tons/hectare. Organic fertilizers contain lots of humic acid and folic acid as well as growth regulators which can accelerate the growth of rice tillers (Brambilla *et al*, 2022). According to Jiang *et al* (2023), the application of organic fertilizers is beneficial in providing and absorbing nutrients, as well as activating soil microorganisms, so that the soil structure becomes crumbly which causes an expansion of root reach and increases the number of rice tillers. The growth of the number of rice tillers is described in Figure 2.



**Figure 2** The growth of the number of rice tillers at various ages of observation and all treatments

Provision of organic fertilizers can improve the quality of bacteria in the soil. Granul organic fertilizer dosage has a significant effect on improving the quality of bacteria in the soil. Soils that are not given organic matter of bacterial quality tend to be low, because the availability and activity of bacteria is needed in the process of absorbing and providing nutrients for plants. Organic fertilizers play an important role in the mineral cycle, especially the nitrogen, phosphorus, sulfur and carbon cycles (Cheng and Wan, 2023). Bacteria that play a role in the nitrogen cycle include *Azotobacter* and *Azospirillum*, these bacteria are non-symbiotic in nature that are able to bind free  $N_2$  which has an influence on the physical and chemical properties of the soil so as to increase soil fertility (Harindintwali *et al*, 2022).




The quality of the bacteria at the beginning of the treatment ranged from  $10^2$  to  $10^3$  and the quality of the bacteria continued to increase after being given treatment. The treatment of using granular organic fertilizer 20 tons hectare<sup>-1</sup> can improve the quality of bacteria from  $10.1^3$  cfu/ml to  $10.1^7$  cfu/ml. In this study it was seen that the more doses of organic fertilizer given to the soil, the higher the quality of the bacteria produced. According to Zainuddin *et al* (2022), the application of organic fertilizers in the long term increases the amount of nutrients such as available nitrogen, available phosphorus and total zinc in the soil and will increase the number and diversity of bacteria in the soil for plant growth and production processes. The increase in the quality of bacteria at the beginning and end of the study is described in Table 1.

**Table 1** Bacterial quality at the beginning and end of the study

No	Treatment	Initial bacterial quality (cfu/ml)	Final bacterial quality (cfu/ml)
1	No fertilizer/control	$10.2^2$	$10.2^2$
2	NPK recommendations	$10.2^2$	$10.2^2$
3	Granular Fertilizer 10 tons/hectare	$10.1^3$	$10.1^6$
4	Granular Fertilizer 10 tons/hectare + NPK recommendation	$10.1^3$	$10.1^6$
5	Granular Fertilizer 20 tons/hectare	$10.1^3$	$10.1^7$
6	Granular Fertilizer 20 tons/hectare + NPK recommendation	$10.1^3$	$10.1^7$

### 3.2 Rice Production

The use of organic fertilizers can restore and increase soil fertility in addition to increasing crop production. Rice plant production through grain yield indicators is the role of organic fertilizer application which can facilitate the physiological processes of plant growth and production. The production results of rice plants are described in Figure 3.



**Figure 3** Production of harvested dry unhulled rice and dry milled unhusked rice (tonnes/hectare)

Figure 3 explains that the highest rice production was found in the treatment of granular organic fertilizer 20 tons/hectare + NPK with a production of 7.4 tons/hectare harvested dry grain, and 4.24 tons/hectare dry milled grain. The weight production of harvested dry grain and dry milled grain in rice plants reflects the accumulation of organic compounds that plants have successfully synthesized from organic compounds, water and carbon dioxide which will contribute to plant dry weight (Alvarez *et al*, 2021). Active division of meristem cells in roots will spur good root development, so that active microbial activity in the root area and energy generated from the respiration process becomes more and supports plant canopy growth, thereby increasing plant dry weight (Olenska *et al*, 2020).

## 4 Conclusion

Granular organic fertilizer application 20 tons/hectare can improve the quality of bacteria from  $10.1^3$  to  $10.1^7$  cfu/ml compared to other treatments. Granular organic fertilizer 20 tons/hectare and recommended NPK is a treatment that can increase the production of harvested dry grain by 7.4 tons/hectare and dry milled grain by 4.24 tons/hectare., when compared to production in the organic fertilizer treatment Granul 20 ton/hectare, as well as the recommended NPK fertilizer treatment.

## Compliance with ethical standards

### *Acknowledgments*

The author would like to thank the Agriculture Office of Bone Bolango Regency, Gorontalo Province, Indonesia, for facilitating the implementation of this research.

### *Disclosure of conflict of interest*

The author declares no conflict of interest.

## References

- [1] Milosevic, T., Milosevic, N. and Mladenovic, J. The influence of organic, organo-mineral and mineral fertilizers on tree growth, yielding, fruit quality and leaf nutrient composition of apple cv. 'Golden Delicious Renders'. *Journal of Scientia Horticulture* 2022. 297:110 - 120.<https://doi.org/10.1016/j.scienta.2022.110978>.
- [2] Weerahewa, J. and Dayananda, D. Land use changes and economic effects of alternative fertilizer policies: A simulation analysis with a bio-economic model for a Tank Village of Sri Lanka. *Journal of Agricultural Systems* 2023. 205:563 - 514.<https://doi.org/10.1016/j.agsy.2022.103563>.
- [3] Puech, T. and Stark, F. Diversification of an integrated crop-livestock system: Agroecological and food production assessment at farm scale. *Journal of Agriculture, Ecosystems & Environment* 2023. 344: 108 - 118.<https://doi.org/10.1016/j.agee.2022.108300>.
- [4] Chen, D., Wang, X., Carrion, VJ, Yin, S., Yue, Z., Liao, Y., Dong, Y. and Li, X. Acidic amelioration of soil amendments improves soil health by impacting rhizosphere microbial assemblies. *Journal of Soil Biology and Biochemistry* 2022. 167: 599 - 609.<https://doi.org/10.1016/j.soilbio.2022.108599>.
- [5] Hui, K., Tan, W. and Song, Q. Long-term application of nitrogen fertilizer alters the properties of dissolved soil organic matter and increases the accumulation of polycyclic aromatic hydrocarbons. *Journal of Environmental Research* 2022. 215: 114 - 125.<https://doi.org/10.1016/j.envres.2022.114267>.
- [6] Gao, Y., Shao, G., Yang, Z., Zhang, K., Lu, J., Wang, Z., Wu, S. and Xu, D. Influences of soil and biochar properties and amount of biochar and fertilizer on the performance of biochar in improving plant photosynthetic rate: A meta-analysis. *European Journal of Agronomy* 2021. 130: 126 - 138.<https://doi.org/10.1016/j.eja.2021.126345>.
- [7] Hameed, A., Chen, YP, Shen, FT, Lin, SY, Huang, HI, Lin, YW and Young, CC Evaluation of a subtropical maize-rice rotation system maintained under sustainable long-term fertilizer inputs for intensification of agriculture. *Journal of Applied Soil Ecology* 2023. 184: 104 - 114.<https://doi.org/10.1016/j.apsoil.2022.104772>.
- [8] Negi, R., Kaur, T., Devi, R., Kour, D. and Yadav, AN Assessment of nitrogen-fixing endophytic and mineral solubilizing rhizospheric bacteria as a multifunctional microbial consortium for growth promotion of wheat and wild wheat relative *Aegilops kotschy*. *Journal Heliyon* 2022. 8:125 - 135.<https://doi.org/10.1016/j.heliyon.2022.e12579>.
- [9] Allouzi, MMA, Allouzi, SMA, Keng, ZX, Supramaniam, CV, Singh, A. and Chong, S. Liquid biofertilizers as a sustainable solution for agriculture. *Journal Heliyon* 2022. 8:609 - 619.<https://doi.org/10.1016/j.heliyon.2022.e12609>.
- [10] Sahab, S., Suhani, I., Srivastava, V., Chauhan, PS, Singh, RP and Prasad, V. Potential risk assessment of soil salinity to agroecosystem sustainability: Current status and management strategies. *Journal of Science of The Total Environment* 2021. 764:144 - 154.<https://doi.org/10.1016/j.scitotenv.2020.144164>.

- [11] Afzal, S. and Singh, NK Effect of zinc and iron oxide nanoparticles on plant physiology, seed quality and microbial community structure in a rice-soil-microbial ecosystem. *Journal of Environmental Pollution* 2022. 314: 120 - 131.<https://doi.org/10.1016/j.envpol.2022.120224>.
- [12] Swetha, RK, Dasgupta, S., Chakraborty, S., Li, B., Weindorf, DC, Mancini, M., Silva, SHG, Ribeiro, BT, Curi, N. and Ray, DP Using Nix color sensor and Munsell soil color variables to classify contrasting soil types and predict soil organic carbon in Eastern India. *Journal of Computers and Electronics in Agriculture* 2022. 199: 107 - 117.<https://doi.org/10.1016/j.compag.2022.107192>.
- [13] Jiang, L., Wu, L., Wang, Y., Xu, Q., Xu, Z. and Chen, W. Research progress on the divergence and genetic basis of agronomic traits in Xian and Gang rice. *The Crop Journal* 2022. 10:924 - 931.<https://doi.org/10.1016/j.cj.2022.02.006>.
- [14] Brambilla, S., Stritzler, M., Soto, G. and Ayub, N. A synthesis of functional contributions of rhizobacteria to growth promotion in diverse crops. *Journal of the Rhizosphere* 2022. 24: 100 - 110.<https://doi.org/10.1016/j.rhisph.2022.100611>.
- [15] Jiang, Z., Li, Y., Cao, B., Li, J., Ren, Z., Qu, J. and Zhang, Y. Noval bio-organic fertilizer containing *Arthrobacter* sp. DNS10 alleviates atrazine-induced growth inhibition on soybean by improving atrazine removal and nitrogen accumulation. *Chemosphere Journal* 2023. 313: 137 - 147.<https://doi.org/10.1016/j.chemosphere.2022.137575>.
- [16] Cheng, Y. and Wan, W. Strong linkage between nutrient-cycling functional gene diversity and ecosystem multifunctionality during winter composting with pig manure and fallen leaves. *Journal Science of The Total Environment* 2023. 867:161 - 172.<https://doi.org/10.1016/j.scitotenv.2023.161529>.
- [17] Harindintwali, JD, Wang, F., Yang, W., Zhou, J., Muhoza, B., Mugabowindekwe, M. and Yu, X. Harnessing the power of cellulolytic nitrogen-fixing bacteria for biovalorization of lignocellulosic biomass. *Journal of Industrial Crops and Products* 2022. 186: 115 - 125.<https://doi.org/10.1016/j.indcrop.2022.115235>.
- [18] Zainuddin, N., Keni, MF, Ibrahim, SAS and Masri, MMM Effect of integrated biofertilizers with chemical fertilizers on oil palm growth and soil microbial diversity. *Journal of Biocatalysis and Agricultural Biotechnology* 2022. 39: 102 - 112.<https://doi.org/10.1016/j.bcab.2021.102237>.
- [19] Alvarez, AL, Weyers, SL, Goemann, HM, Peyton, BM and Gardner, RD Microalgae, soil and plants: A critical review of microalgae as a renewable resource for agriculture. *Journal of Algal Research* 2021. 54:102 - 112.<https://doi.org/10.1016/j.algal.2021.102200>.
- [20] Olenska, E., Malek, W., Wojcik, M., Swiecicka, I., Thijs, S. and Vangronsveld, J. Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. *Journal of Science of The Total Environment* 2020. 743:140 - 150.<https://doi.org/10.1016/j.scitotenv.2020.140682>