Bukti Korespondensi

The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)"

| | | × 荘 | O Perg | • 🤉 🕸 | Ⅲ | UN | |
|--|--|--|---|---------------------|-----------|-----------|---------|
| • • • • • | 0 64 28 D : | | | 3 | dari 3 | < > | / |
| Manuscript recept | ion confirmation (Eksternal) 🤉 Kotak Ma | esuk x | | | | × | 9 |
| muntean.gc@gmail.com kepada saya 👻 | | | | Sen, 14 Sep 20. | 20, 19.06 | ☆ | ¢ |
| 🛪 Inggris 🗸 🖒 Indone | esia 🔹 Terjemahkan pesan | | | No | naktifkan | untuk: In | ggris ; |
| | | | | | | | |
| Dear Dr. Sri N. Hamzah, This is a message of confirma Regency)", submitted to our jou I must inform you that you did n | tion regarding the reception of your manuscript "The sumal "Aquaculture, Aquarium, Conservation & Legislation", ot provide us with a submission letter redacted according | uccess of coral rehabilitation through trans I will contact you with further information re to our format. Please redact one and e-mail | plantation using spider modules (ca garding your manuscript. I it directly to me. | se study: Botutonuo | marine ar | rea, Bon | e Bola |
| Dear Dr. Sn N. Hamzah, This is a message of confirma Regencyl [*] , submitted to our jou I must inform you that you did n For any questions, please conta Please try to keep one message | tion regarding the reception of your manuscript "The s mai "Aquacuture, Aquanum, Conservation & Legislation". at provide us with a submission letter redacted according ct me. thread (respond/ask with "Reply") for an easier message | uccess of coral rehabilitation through trans I will contact you with further information re to our format. Please redact one and e-mail management. | plantation using spider modules (ca garding your manuscript. I it directly to me. | se study: Botutonuo | marine ar | rea, Bon | e Bola |
| Dear Dr. Sn N. Hamzah, This is a message of confirma Regency?, submitted to our jou I must inform you that you did n For any questions, please conta Please try to keep one message Thank you for considering our jo | tion regarding the reception of your manuscript "The s mai "Aquaculture, Aquarium, Conservation & Legislation". ot provide us with a submission letter redacted according ct me. thread (respond/ask with "Reply") for an easier message sumal. | uccess of coral rehabilitation through trans I will contact you with further information re to our format. Please redact one and e-mail management. | plantation using spider modules (ca garding your manuscript. it directly to me. | se study: Botutonuo | marine a | rea, Bon | e Bola |



Submission letter

Article title:

The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)

Name of the authors: Sri Nuryatin Hamzah, Sitti Nursinar

Hereby I would like to submit the manuscript entitled "The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)" to Aquaculture, Aquarium, Conservation & Legislation -International Journal of the Bioflux Society.

This manuscript was not submitted or published to any other journal. The authors declare that the manuscript is an original paper and contain no plagiarised text. All authors declare that they are not currently affiliated or sponsored by any organization with a direct economic interest in subject of the article. My co-authors have all contributed to this manuscript and approve of this submission.

Corresponding author

Name : Sri Nuryatin Hamzah

Signature : WAGE

Date : 14 September 2020

| M Gmail | | Q muntean X ⅔ O Pergi ▼ ⑦ ፡፡ | |
|------------------------------------|-------|--|--------------------------|
| 🖉 Tulis | | | ari 3 < ゝ 🎢 🕶 |
| 🖵 🛛 Kotak Masuk | 1.018 | Sri Nuryatin Hamzah <sri.nuryatin@ung.ac.id> @ 20 Agu 2021</sri.nuryatin@ung.ac.id> | 1, 20.05 🛧 🕤 🚦 |
| 🕁 Berbintang | | Dear Dr. George Muntean | |
| 🕚 Ditunda | | Editor of AACL Bioflux Journal, | |
| D Penting | | I hope you are well. I would be grateful if you could let me know whether there has been any further progress on the review of our manusci | ript with the title "The |
| ▶ Terkirim | | Success of Coral Rehabilitation through Transplantation Using Spider Modules (Case Study: Botutonuo Marine Area, Bone Bolango Reger | ncy). |
| 🗋 Draf | 2 | As a reminder, here I am sending the certificate of preliminary accepted for publication that we have received since October 2, 2020. And a | also, the payment of our |
| - 🕞 Kategori | | article has been received on October 12, 2020. I do wish that our research article is adequate to meet all the requirements of Bioflux AACI | L. |
| ළ, Sosial | 573 | I am looking forward to hearing from you soon. Thank you very much. | |
| (i) Update | 198 | Kind regards, | |
| 📮 Forum | 2 | Sri Nuryatin Hamzah | |
| 🟷 Promosi | 155 | | |
| ✓ Selengkapnya | | Satu lampiran • Dipindai dengan Gmail 🛈 | ۵. |
| Label | + | VOL JANE YAN | |
| Prioritas | | Bin Register Strategy | |
| Serbaneka | | | |



Bioflux (publishing house) 54 Ceahlau Street, Cluj-Napoca 400488, Romania, European Union

Certificate

This certificate shows that the manuscript entitled "The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)", authored by Sri N. Hamzah and Sitti Nursinar, submitted on 09.12.2020, to our journal Aquaculture, Aquarium, Conservation & Legislation -International Journal of the Bioflux Society (AACL Bioflux), has been preliminary accepted for publication.

Thank you for your participation!

Cordially yours,

Editor, George-Cătălin Muntean

Senior Researcher Ioan Valentin Petrescu-Mag, PhD Editor-in-chief



| M | Gmail | | Q muntean | × C Pergi |
|--------------------|--------------|-------|---|--|
| 1 | Tulis | | | 1 dari 3 < > > |
| V | Kotak Masuk | 1.018 | kepada saya 🔹 | |
| ☆ | Berbintang | | XA Inggris • > Indonesia • Terjemahkan pesan | ivonaktirkan untuk. Inggits X |
| 0 | Ditunda | | Dear Dr. Sri N. Hamzah, I have compiled all the necessary modifications indicated by the reviewers in the | e comments made in this version of your manuscript |
| Σ | Penting | | Address all the issues an solve them. Make sure that Track Changes is activated | d (see attachment) so you can see the comments and where I asked you to use the |
| ⊳ | Terkirim | | Reply function from the comment (see attachment), do so. Work ONLY in this document, because this is a protected version of your manusc | cript that allows me to easily track your changes and to final edit your manuscript. |
| D | Draf | 2 | Make sure that all the authors information is correct, both at the beginning of the r the required information. | manuscript and at the end, in the information cassette, in which you need to provide |
| • 🗅 | Kategori | | Let me know if you have any questions. | |
| De | Sosial | 573 | After you are finished, send the manuscript back to me and I will proceed to the p | publishing process with it and send you a final preview. |
| (j | Update | 198 | | |
| 믹 | Forum | 2 | 3 Lampiran • Dipindai dengan Gmail 🗊 | ± @ |
| \bigtriangledown | Promosi | 155 | ung Citatin Muntaan Giday | |
| ~ | Selengkapnya | | w that you removed some author is reply. | In a rehabilitation through many space model (close strafe; and, space feature galaxy (space) a model of the strategies |
| Labe | el | + | E Reply Reply with the replacement of the logs with the | .doc |
| | Prioritas | | too some firstened with white private same | |



The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)

¹Sri N. Hamzah, ¹Sitti Nursinar

¹ Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Gorontalo City, Indonesia. Corresponding author: S. N. Hamzah, sri.nuryatin@ung.ac.id

Abstract. Botutonuo Waters, Bone Bolango Regency, is an area that has coral reef potential for marine tourism practices. However, the high human activity in this water area causes the coral reef ecosystem to be stressed, and it is in the moderate to damaged category. One of the alternatives that can be implemented to protect and conserve coral reefs is through coral transplantation. This study was conducted to provide information about the success of transplantation using the spider module. The experiment was conducted for two months by placing ten spider modules in the water. A total of 60 coral fragments were used where six coral fragments were tied to each module. Coral growth monitoring was carried out every two weeks for four times. The results revealed that the absolute growth of corals for two months of observation showed positive results, namely 1.52 cm. The growth rate of the transplanted corals was 0.19 cm/week or 0.03 cm/day. The survival rate of transplanted corals was 100% and is included in the category of high transplant success rate. **Key Words**: Botutonuo waters, coral reef, growth, spider, survival rate.

Introduction. Coral reefs are one of the ecosystems that are crucial for the sustainability of existing resources in coastal areas and oceans. The existence of this ecosystem provides many benefits not only for the biota association but also for humans. Rusandi et al (2016) stated that ecologically, coral reefs function as a habitat for various reef fish caught by fishermen, as a coast protector from waves, a source of germplasm, and economically as a source of income through marine tourism and ornamental coral trades. Currently, the rate of utilization of coral reefs in coastal areas. Giyanto et al (2017) reported that the current condition of coral reefs in coastal areas. Giyanto et al (2017) reported that the current condition of coral reefs in Indonesia was 36.18% in the bad category, 34.3% in the moderate category, 22.96% in the good category, and only 6.56% in the very good category. In general, the trend of the condition of live coral cover is decreasing, which deserves serious attention due to the huge role of coral reefs in the coastal environment. Even some research results indicate that coastal areas with damaged coral reefs have the potential to receive bigger waves and are threatened with coastal abrasion.

Botutonuo waters are one of the waters in the Tomini Bay area of Gorontalo. Currently, the Botutonuo water area is used for various activities such as fishing and marine tourism. Information obtained from the coastal community of Botutonuo, that in 1980–1990 destructive fishing practices were taking place. The amount of rubble was found to be an indicator of destructive fishing in the past. If no measures are taken to restore this condition, the damage to coral reefs can get worse. Restoration of coral reef ecosystems can occur naturally. However, the time required is quite long, so interventions are needed to restore the condition of coral reefs in numerous locations in a short time.

Several methods have been implemented to restore coral reef conditions in a shorter

Commented [GCM1]: Rephrase. Explain better.

Commented [GCM2]: Rephrase. The connection between utilization and regeneration is not clear.

Commented [GCM3]: Rephrase and explain better. You obtained the information directly? If the information comes from a published source, it needs to be cited and the appropriate reference provided. Use Reply.

Commented [GCM4]: If the information comes from a published source, it needs to be cited and the appropriate reference provided. Use Reply.

time, including artificial reef methods, mineral accretion (bio rock), and coral transplantation. Coral transplantation is a method of coral cultivation by cutting a portion of a particular coral colony to be planted in a new place (Prameliasari et al 2012). Transplantation through asexual fragmentation is essential as an effort to reproduce corals for recovery, trade, and marine tourism (Nedimyer et al 2011; Suharsono et al 2013). Transplantation using spider modules has never been tested in Gorontalo Province since this method is the newest and not widely used in Indonesia. It was identified that transplanting with the spider module was only found in South Sulawesi and Bali. Therefore, there is a lack of information about coral growth using the spider module.

This study aims to provide information about coral growth and survival rates using the spider module as well as a reference in selecting a transplant model that is cheap and easy to apply in the field.

Material and Method

Study site. This research was conducted in May-July 2020 at Botutonuo waters, Bone Bolango Regency. The research location is depicted in Figure 1.

LEGEND Area of Tatteret (AOT) The transplantation the transplantation Map Scale 1 cm = 185 m

Figure 1. Location of coral transplantation using spider modules.

Preparation and module construction. The placement of the spider module was determined in the initial observation, namely through the manta tow survey, to see the condition of the coral reefs around the Botutonuo waters. The spider module was placed on the bottom of the water where damaged coral reef conditions were observed. The spider construction can be seen in Figure 2.



Figure 2. Spider module construction (A); illustration of coral fragments tied to spider modules (B).

Commented [GCM5]: If the information comes from a published source, it needs to be cited and the appropriate reference provided. Or is this part of the information from Prameliasari et al 2012? Use Reply.

Commented [GCM6]: Reference needed. Cite and add reference to reference list.

The spider construction was made of steel with a diameter of 12 mm, which was shaped to resemble a spider's web and legs (Figure 2). The frame height is \pm 50 cm, with the bottom diameter of \pm 120 cm, and the distance among the legs is \pm 60 cm. Each spider frame was coated with sand, which is glued using resin and serves to prevent rust from the iron when the module was lowered into the water.

Seed procurement, fragmentation, and fragment binding. Coral seedlings were obtained from broodstock nearby the transplant site. The seedlings were taken by cutting the parent corals from the site using pliers and scissors into small colonies (fragments) with a size range of 3–12 cm for branching corals (Rani et al 2017; Suharsono et al 2013). The coral fragments were then put in baskets and soaked with seawater to keep the coral fragments from stress. The coral taken was fragments of *Acropora* sp. as the species generally grows faster. The coral fragments collected were then tied using cable ties to the spider frame, as many as 6 fragments for each module, so that a total of 60 fragments were tied to 10 spider modules.

Measurement of growth and survival rates of coral fragments. Coral fragment growth measurements were carried out every two weeks for two months. At the start of rearing, transplanted corals were measured as initial length using a caliper. Furthermore, every two weeks, the increase in branch length in the waters was measured manually. Also, the number of dead or missing fragments in each experimental unit was counted to determine the survival rate. During the measurement of coral growth, cleaning of the module area, which was overgrown with algae, was also carried out to prevent the death of coral fragments due to algal dominance.

Measurement of water quality parameters. Water quality parameters measured in this study were the temperature, salinity, pH, dissolved oxygen (DO), current, and transparency. Measurement of the parameters was performed in situ along with the measurements of coral growth.

Data analysis. The results of coral reef measurements were tabulated and analyzed. Some of the analysis parameters are the following:

1. Absolute growth

Coral growth over a period was calculated using Effendie (1979) formula:

$$\beta = Lt - Lo$$

where: β = absolute growth (cm); Lt = the average branch length after the tth observation; Lo = average branch length at the beginning of research.

2. Coral growth rate Coral Growth Rate was calculated using Effendie (1979) formula:

$$P = \frac{Lt - L0}{t}$$

where: P = increase in length/height of transplanted coral fragments; Li+1 = the mean length/height of the fragments after the tth observation; L0 = the mean length/height of the initial fragments; t = observation time.

3. Survival rate

The formula (Effendi 1979) used for the coral survival rate is:

Commented [GCM7]: Mention the devices used to determine these parameters.

$$S = \frac{Nt}{N_0} \times 100\%$$

where: S = survival rate (%); Nt = Number of coral fragments at the end of the observation; and No = Number of coral pieces at the beginning of observation.

Results and Discussion

The range of coral lengths at the beginning of the study was 8.67-10.08 cm, with an average of 9.36 cm. The measurement results showed that there was coral growth during the two months of observation (Figure 3).



Figure 3. Coral growth using spider modules

Figure 3 shows that there was an increase in coral growth at the research location every two weeks of measurement for two months of field observation. The average size of the initial coral fragments was 9.36 cm and at the end of the study, in the fourth observation, the length was 10.88 cm. The results designated that the absolute growth of the transplanted corals was 1.52 cm. The absolute growth of *Acropora* sp., which was transplanted using this spider, was higher than that of Subhan (2019) and Rani et al (2017), which were 0.59 cm and 0.1-0.2 cm, respectively. The results showed that the coral growth observation experiment in the Botutonuo waters shows better performance than previous studies using the same transplant module. Furthermore, absolute growth in this study is better than the results obtained by Alim (2019) in the same research location with the concrete beam transplant module. Alim (2019) reported that the absolute growth of coral during two months of observation was 0.6 cm. Based on visual observation, the transplanted coral began to stick to the spider module, and some coral fragments had started to wrap the ties (Figure 4).





Observation I (Week 0)

Observation II (Week 2)



Observation III (Week 4)



Figure 4. Observation of growth and morphological changes in transplanted corals using spider modules.



Figure 5. Coral fragments attached to observations II and III (week 2 and week 4). Figure 4 shows changes in the morphology of the transplanted corals, both changes in the height increase of corals and the adding of new coral branches. The life of coral fragments in the transplant module can be observed through the tie cables and the attachment of elements (Figure 5) to the substrate. Coral fragments are categorized as living on the transplant media when they are attached or when the binding cables have been wrapped by new coral tissue at the end of the transplant. According to Edward and Gomez (2007), corals that have stuck to and covered the substrate are less likely to escape.

Growth rate. In general, the growth rate of corals transplanted using spiders was 0.19 cm/week or 0.03 cm/day. The results of measuring coral growth rate can be seen in Figure 6.



Figure 6. Coral growth rate using the spider module.

Figure 6 shows that the transplanted *Acropora* sp. growth rates were 0.17 cm/week or 0.02 cm/day, 0.25 cm/week or 0.04 cm/day, and 0.34 cm/week or 0.05 cm/day on the second, third, and fourth observations, consecutively. The results showed that the coral growth rate

of *Acropora* sp., transplanted using the spider method, was faster than Subhan (2019) results in Bali using the same coral genus, method, and depth. Where the growth rate of corals obtained was 0.09 cm/week in observation II, and 0.14 cm/week in observation III and IV.

The use of the spider transplant modules in this study also showed a better growth rate performance compared to the use of the concrete block transplant module in Botutonuo waters by Alim (2019), where the growth rates on the blocks were 0.01 cm/day in the second and third week of observation, and 0.03 cm/day at the fourth week of observation. The research informs that applying the spider transplant module is more effective in rehabilitating coral reefs and generates growth faster than the concrete beam transplant. Although to see to what extent its effectiveness and development still needs to be investigated through further observation and continuous monitoring.

Coral survival rate using spider modules. The analysis of coral survival rates using the spider module can be seen in Figure 7.



Figure 7. The survival rate of *Acropora* sp. for two months of observation.

Figure 7 shows that during the two months of observation, there were no dead coral fragments or coral fragments that had escaped from the spider module. The total number of coral fragments observed at the beginning and the end of the observation amounted to 60 fragments. The results of this study indicate that the use of spider modules for coral transplantation is successful. Harriot and Fisk (1988) in Mompala et al (2017) stated that transplantation of corals was considered successful when the survival rate was in the range of 50–100%. The analysis of the coral survival rate in this study was higher than the results of Subhan's research (2019). There was a decrease in the survival rate of corals using the spider method for a depth of 6 meters, which was only 82% at the end of the study.

Similarly, the results of research conducted by Rani et al (2017) with the same method showed a coral survival rate of 95.56% at the end of the study. The study results indicate that *Acropora* sp. can grow well in Botutonuo waters. Williams et al (2019) reported that transplanting using the spider method succeeded in restoring the condition of coral reefs in the rehabilitation area.

The main factor causing the high survival rate in this study lies in the initial transplantation treatment, starting from finding and tying seedlings to mobilizing spider modules to the research location, being carried out carefully, and trying not to be exposed to open air, thereby minimizing stress levels on corals.

Water quality parameters. Water quality parameters are one of the essential factors that play a role in the survival of aquatic organisms, especially coral reefs. It causes the

Commented [GCM8]: Rephrase. Is this from Subhan (2019) research? If so, rephrase so that the citation is appropriate. Use Reply.

Commented [GCM9]: Although this is cited in Mompala et al (2017), you need to add this to the reference list also. Please add.

Commented [GCM10]: Rephrase. Again, not clear if this is from Subhan (2019) research. Use Reply and answer me.

distribution of coral reefs in water areas to be uneven. The in-situ measurements of water quality parameters can be seen in Table 1.

Table 1

Results of measurement of water quality parameters

| | | Obser | vation | Seawater Quality Standards for Coral | |
|------------------|------|-------|--------|---|--|
| Parameters | Ι | II | III | IV | Reefs (Minister of Environment Decree No. 51/2004) |
| Temperature (°C) | 29 | 29 | 29 | 29.3 | 28-30 |
| Salinity (‰) | 29 | 29 | 25.5 | 27.7 | 33-34 |
| рН | 7.98 | 8.01 | 7.89 | 7.57 | 7-8.5 |
| DO (mg/l) | 2.5 | 6.09 | 3.99 | 5.92 | >5 |
| Current (m/s) | 0.02 | 0.015 | 0.02 | 0.03 | - |
| Transparency (m) | 6 | 6 | 6 | 6 | > 5 m |

Table 1 shows that the temperature values in Botutonuo waters are in the range of $29-29.3^{\circ}$ C. Supriharyono (2000) stated that a suitable temperature for coral growth ranges from 25-29°C. However, coral animals can tolerate temperatures with a minimum of 16°C and a maximum of 36°C (Kinsman 1964). Salinity ranges between 25.5-29 ‰. Low salinity value is thought to be influenced by rainfall due to rain during the research. In addition, low salinity is also predicted to be affected by the supply of fresh water from the rivers around the study site. However, it is a rainfed river, not a continuous stream. The results of pH measurements at the study sites were in the range of 7.89–8.01. This value is in the normal pH range in Indonesian waters and the range of pH in Indonesia varies from 7.7 to 8.2.

The dissolved oxygen (DO) measurements at the research location were in the range 2.5-6.05 mg/L. The range of DO for the first and second observations is below the standard quality for coral reefs based on Minister of Environment Decree No. 51/2004. However, according to Swingle (1968) in Salmin (2005), the minimum DO of 2 mg/L is sufficient to support the life of organisms.

The current velocity obtained at the research location ranging from 0.015 to 0.02 m/s is in a low category when referring to Yusuf et al (2012), which is below 0.5 m/s. The slow movement of the current is due to the morphology of the bay-shaped location so that the current velocity is low. The results of measuring the transparency of Botutonuo waters obtained of 100% or 6 m, as the spider module was placed at a depth of 6 m. Based on the seawater quality standards for coral reefs, the water transparency in Botutonuo is still in the range of quality standard values.

Conclusions. To sum up, absolute coral growth for two months of observation showed a positive result of 1.52 cm. The growth rate of the transplanted corals during the study was 0.19 cm/week or 0.03 cm/day. The 100% survival rate is included in the transplant category with a high success rate. The quality of Botutonuo waters as a transplantation location using spider modules still supports the growth of transplanted corals.

Acknowledgments. We express our gratitude to LP2M UNG for the reliance given to us to conduct this research through PNBP UNG funding. Our thanks also go to Hidayat Bachtiar, S. Kel., Alinton Pisuna, Dadang S. Alim, S.Pi., Jufriyanto Umar, S.Pi., Ronal Lahmutu for their assistance during data collection and Nuralim Pasisingi, S.Pi., M.Si., for her assistance in writing.

Commented [GCM11]: Just like the reference situation before, you need to add this also to the reference list.

References

- Alim D. S., 2019 [Analysis of growth rate and survival of transplanted corals in Botutonuo Beach waters, Bone Bolango Regency, Gorontalo Province]. Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Gorontalo State University, Gorontalo [in Indonesian].
- Edward A. J., Gomez E. D., 2007 Reef restoration concepts and guidelines: making sensible management choices in the face of uncertainty. The Coral Reef Targeted Research & Capacity Building for Management Programme: St Lucia, Australia, 38 p.
- Effendie M. I., 1979 [Fisheries biological methods]. Yayasan Dewi Sri, Bogor, 112p [in Indonesian].
- Giyanto, Abrar M., Hadi T. A., Budiyanto A., Hafizt M., Salatalohy A., Iswari M. Y., 2017 [Status of Indonesia's coral reef 2017]. Oceanography Research Center - Indonesian Institute of Sciences [in Indonesian].
- Kinsman J. J., 1964 Reef coral tolerance to high temperatures and salinities. Nature 202:1280-1282.
- Minister of Environment, 2004 [Decree of the Minister of Environment No. 51 year 2004 about water quality standard for coastal waters]. Minister of Environment of the Republic of Indonesia, Jakarta [in Indonesian].
- Mompala K., Rondonuwu A. B., Rembet U. N. W. J., 2017 [The growth rate of Acropora sp. transplanted on artificial reefs in Kareko Waters of North Lembeh Sub-District of Bitung City]. Jurnal Ilmiah Platax 5(2):234–242 [in Indonesian].
- Nedimyer K., Gaines K., Roach S., 2011 Coral Tree Nursery©: An innovative approach to growing corals in an ocean-based field nursery. AACL Bioflux 4(4):442-446.
- Prameliasari R. T. A., Munasik, Wijayanti D. P., 2012 [The effect of fragment size differences and transplantation methods on the growth of *Pocillopora damicornis* corals in Awur Bay, Jepara]. Journal of Marine Research 1(1):159-168 [in Indonesian].
- Rani C., Tahir A., Jompa J., Faisal A., Yusuf S., Werorilangi S., Arniati, 2017 [The success of rehabilitation of coral reefs due to the 2016 bleaching incident with the transplant technique]. Spermonde 3(1):13–19 [in Indonesian].
- Rizki T. Y., Tito C. K., Setiawan A., 2011 [pH variations in Indonesian waters]. https://www.researchgate.net/profile/Camellia_Tito/publication/262673263_Variasi_p H_di_Perairan_Indonesia/links/5645a1d908aef646e6cca454/Variasi-pH-di-Perairan-Indonesia.pdf [in Indonesian].
- Rusandi A., Suharsono, Wiadnyana N., Yusri S., 2016 [National action plan for coral conservation period 1: 2017-2021]. Directorate of Conservation and Marine Biodiversity, Directorate General of Marine Spatial Management, Ministry of Marine Affairs and Fisheries [in Indonesian].
- Salmin, 2005 [Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD) as indicators for determining water quality]. Oseana XXX(3):21–26 [in Indonesian].
- Subhan M. A., 2019 [Growth rate of Acropora loripes coral reef using transplantation method of spider skeleton module in the waters of Les Village, Buleleng Regency, Bali]. Universitas Islam Negeri Syarif Hidayatullah [in Indonesian].
- Suharsono, Siringoringo R. M., Hadi T. A., Giyanto, Tuti Y., Budiyanto A., Sulha S., 2013 [Development of coral transplant techniques in Indonesia]. Oceanography Research Center - Indonesian Institute of Sciences, Jakarta [in Indonesian].
- Supriharyono, 2000 [Coral reef ecosystem management]. Djambatan, Jakarta, 108 p [in Indonesian].
- Williams S. L., Sur C., Janetski N., Hollarsmith J. A., Rapi S., Barron L., Heatwole S. J., Yusuf A. M., Yusuf S., Jompa J., Mars F., 2019 Large-scale coral reef rehabilitation after blast fishing in Indonesia: Coral reef rehabilitation. Restoration Ecology, 27(2):447–456.

Yusuf M., Handoyo G., Wulandari S. Y., 2012 [Characteristics of flow patterns in relation to water quality conditions and abundance of phytoplankton in the waters of the Karimun Jawa Marine National Park]. Buletin Oseanografi Marina 1:63-74 [in Indonesian].

Received: 12 September 2020. Accepted: 02 October 2020. Published online: xxxx 2021.

Authors: [author full name, author affiliation, e-mail:] [author full name, author affiliation, e-mail:]

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: [in order of importance, authors full last name and abbreviated first names] 2021 The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency). AACL Bioflux 14(X):xxxx-xxxx.

Commented [GCM12]: Fill in according to the instructions in the parentheses.

Commented [GCM13]: Fill in according to the instructions in the parentheses.



Sri Nuryatin Hamzah <sri.nuryatin@ung.ac.id> kepada muntean.gc • 📼 17 Sep 2021, 20.20 🔥 🕤 🚦

Dear Dr. George Muntean Editor, Bioflux SRL

I have read and highlighted some corrections on my manuscript (please find attached to this email) and I put Identify of the authors as well. I look forward to your next email. Thank you very much

Best regards

Sri Nuryatin Hamzah

Satu lampiran • Dipindai dengan Gmail 🛈





The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency)

¹Sri N. Hamzah, ¹Sitti Nursinar

¹ Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Gorontalo City, Indonesia. Corresponding author: S. N. Hamzah, sri.nuryatin@ung.ac.id

Abstract. Botutonuo Waters, Bone Bolango Regency, is an area that has coral reef potential for marine tourism practices. However, the high human activity in this water area causes the coral reef ecosystem to be stressed, and it is in the moderate to damaged category. However, the high human activity such as destructive fishing and increasing population in coastal area causes the coral reef ecosystem to be stressed and damaged. One of the alternatives that can be implemented to protect and conserve coral reefs is through coral transplantation. This study was conducted to provide information about the success of transplantation using the spider module. The experiment was conducted for two months by placing ten spider modules in the water. A total of 60 coral fragments were used where six coral fragments were tied to each module. Coral growth monitoring was carried out every two weeks for four times. The results revealed that the absolute growth of corals for two months of observation showed positive results, namely 1.52 cm. The growth rate of the transplanted corals was 0.19 cm/week or 0.03 cm/day. The survival rate of transplanted corals was 100% and is included in the category of high transplant success rate. **Key Words**: Botutonuo waters, coral reef, growth, spider, survival rate.

Introduction. Coral reefs are one of the ecosystems that are crucial for the sustainability of existing resources in coastal areas and oceans. The existence of this ecosystem provides many benefits not only for the biota association but also for humans. Rusandi et al (2016) stated that ecologically, coral reefs function as a habitat for various reef fish caught by fishermen, as a coast protector from waves, a source of germplasm, and economically as a source of income through marine tourism and ornamental coral trades. Currently, the rate of utilization of coral reefs is not as fast as its regeneration rate, so that it has an impact on the damage to coral reefs in coastal areas. Currently, the rate of exploitation of coral reefs is not as fast as the rate of regeneration, so that it has an impact on damage to coral reefs in coastal areas. Giyanto et al (2017) reported that the current condition of coral reefs in Indonesia was 36.18% in the bad category, 34.3% in the moderate category, 22.96% in the good category, and only 6.56% in the very good category. In general, the trend of the condition of live coral cover is decreasing, which deserves serious attention due to the huge role of coral reefs in the coastal environment. Even some research results indicate that coastal areas with damaged coral reefs have the potential to receive bigger waves and are threatened with coastal abrasion.

Botutonuo waters are one of the waters in the Tomini Bay area of Gorontalo. Currently, the Botutonuo water area is used for various activities such as fishing and marine tourism. Information obtained from the coastal community of Botutonuo, that in 1980-1990 destructive fishing practices were taking place. Information obtained from the coastal community of Botutonuo that in 1980-1990 fishing activities by destructive methods mainly were carried out in this water area. The amount of rubble was found to be an indicator of destructive fishing in the past. If no measures are taken to restore this condition, the damage to coral reefs can get worse. Restoration of coral reef ecosystems can occur naturally. However, the time required is quite long, so interventions are needed to restore the condition of coral reefs in numerous locations in a short time.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

2445

Commented [GCM14]: Rephrase. Explain better.

Commented [SNH15R14]: I have highlighted the corrected sentence with yellow colour

Commented [GCM16]: Rephrase. The connection between utilization and regeneration is not clear.

Commented [SNH17R16]: I have highlighted the corrected sentence with yellow colour

Commented [GCM18]: Rephrase and explain better. You obtained the information directly? If the information comes from a published source, it needs to be cited and the appropriate reference provided. Use Reply.

Commented [SNH19R18]: Yes, I obtain the information directly through interviews with the coastal community of Botutonuo

Commented [GCM20]: If the information comes from a published source, it needs to be cited and the appropriate reference provided. Use Reply.

Commented [SNH21R20]: Information is written based on interviews and direct observations, where rubble has been found in a large area, which indicates that this area has experienced destructive fishing.

Several methods have been implemented to restore coral reef conditions in a shorter time, including artificial reef methods, mineral accretion (bio rock), and coral transplantation (Suharsono et al 2013). Coral transplantation is a method of coral cultivation by cutting a portion of a particular coral colony to be planted in a new place (Prameliasari et al 2012). Transplantation through asexual fragmentation is essential as an effort to reproduce corals for recovery, trade, and marine tourism (Nedimyer et al 2011; Suharsono et al 2013). Transplantation using spider modules has never been tested in Gorontalo Province since this method is the newest and not widely used in Indonesia. It was identified that transplanting with the spider module was only found in South Sulawesi and Bali (William et al 2019; Subhan 2019). Therefore, there is a lack of information about coral growth using the spider module.

This study aims to provide information about coral growth and survival rates using the spider module as well as a reference in selecting a transplant model that is cheap and easy to apply in the field.

Material and Method

Study site. This research was conducted in May-July 2020 at Botutonuo waters, Bone Bolango Regency. The research location is depicted in Figure 1.



Figure 1. Location of coral transplantation using spider modules.

Preparation and module construction. The placement of the spider module was determined in the initial observation, namely through the manta tow survey, to see the condition of the coral reefs around the Botutonuo waters. The spider module was placed on the bottom of the water where damaged coral reef conditions were observed. The spider construction can be seen in Figure 2.



Figure 2. Spider module construction (A); illustration of coral fragments tied to spider modules (B).

The spider construction was made of steel with a diameter of 12 mm, which was shaped to resemble a spider's web and legs (Figure 2). The frame height is \pm 50 cm, with the bottom diameter of \pm 120 cm, and the distance among the legs is \pm 60 cm. Each spider

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

2446

Commented [GCM22]: If the information comes from a published source, it needs to be cited and the appropriate reference provided. Or is this part of the information from Prameliasari et al 2012? Use Reply.

Commented [SNH23R22]: I've added the published source

Commented [GCM24]: Reference needed. Cite and add reference to reference list. Commented [SNH25R24]: I've added the published source frame was coated with sand, which is glued using resin and serves to prevent rust from the iron when the module was lowered into the water.

Seed procurement, fragmentation, and fragment binding. Coral seedlings were obtained from broodstock nearby the transplant site. The seedlings were taken by cutting the parent corals from the site using pliers and scissors into small colonies (fragments) with a size range of 3-12 cm for branching corals (Rani et al 2017; Suharsono et al 2013). The coral fragments were then put in baskets and soaked with seawater to keep the coral fragments from stress. The coral taken was fragments of Acropora sp. as the species generally grows faster. The coral fragments collected were then tied using cable ties to the spider frame, as many as 6 fragments for each module, so that a total of 60 fragments were tied to 10 spider modules.

Measurement of growth and survival rates of coral fragments. Coral fragment growth measurements were carried out every two weeks for two months. At the start of rearing, transplanted corals were measured as initial length using a caliper. Furthermore, every two weeks, the increase in branch length in the waters was measured manually. Also, the number of dead or missing fragments in each experimental unit was counted to determine the survival rate. During the measurement of coral growth, cleaning of the module area, which was overgrown with algae, was also carried out to prevent the death of coral fragments due to algal dominance.

Measurement of water quality parameters. Water quality parameters measured in this study were the temperature, salinity, and dissolved oxygen (DO) using a Lutron DO-5510, pH using pH Meter AS-218, current using Flowatch Meter FL-03, and transparency using Secchi Disk. Measurement of the parameters was performed in situ along with the measurements of coral growth.

Data analysis. The results of coral reef measurements were tabulated and analyzed. Some of the analysis parameters are the following:

4. Absolute growth Coral growth over a period was calculated using Effendie (1979) formula:

 $\beta = Lt - Lo$

where: β = absolute growth (cm); Lt = the average branch length after the tth observation; Lo = average branch length at the beginning of research.

5. Coral growth rate Coral Growth Rate was calculated using Effendie (1979) formula:

$$P = \frac{Lt - L0}{t}$$

where:

P = increase in length/height of transplanted coral fragments; Li+1 = the mean length/height of the fragments after the t^{th} observation; L0 = the mean length/height of the initial fragments; t = observation time.

Survival rate The formula (Effendi 1979) used for the coral survival rate is:

$$S = \frac{Nt}{N_0} \times 100\%$$

S = survival rate (%); Nt = Number of coral fragments at the end of the where: observation; and No = Number of coral pieces at the beginning of observation.

2447

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

Commented [GCM26]: Mention the devices used to determine these parameter

Commented [SNH27R26]: I have highlighted the devices used in water quality measurement

Results and Discussion

The range of coral lengths at the beginning of the study was 8.67-10.08 cm, with an average of 9.36 cm. The measurement results showed that there was coral growth during the two months of observation (Figure 3).



Figure 3. Coral growth using spider modules

Figure 3 shows that there was an increase in coral growth at the research location every two weeks of measurement for two months of field observation. The average size of the initial coral fragments was 9.36 cm and at the end of the study, in the fourth observation, the length was 10.88 cm. The results designated that the absolute growth of the transplanted corals was 1.52 cm. The absolute growth of *Acropora* sp., which was transplanted using this spider, was higher than that of Subhan (2019) and Rani et al (2017), which were 0.59 cm and 0.1-0.2 cm, respectively. The results showed that the coral growth observation experiment in the Botutonuo waters shows better performance than previous studies using the same transplant module. Furthermore, absolute growth in this study is better than the results obtained by Alim (2019) in the same research location with the coral during two months of observation was 0.6 cm. Based on visual observation, the transplanted coral began to stick to the spider module, and some coral fragments had started to wrap the ties (Figure 4).

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl





Observation I (Week 0)

Observation II (Week 2)



Figure 4. Observation of growth and morphological changes in transplanted corals using spider modules.



Figure 5. Coral fragments attached to observations II and III (week 2 and week 4).

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

Figure 4 shows changes in the morphology of the transplanted corals, both changes in the height increase of corals and the adding of new coral branches. The life of coral fragments in the transplant module can be observed through the tie cables and the attachment of elements (Figure 5) to the substrate. Coral fragments are categorized as living on the transplant media when they are attached or when the binding cables have been wrapped by new coral tissue at the end of the transplant. According to Edward and Gomez (2007), corals that have stuck to and covered the substrate are less likely to escape.

Growth rate. In general, the growth rate of corals transplanted using spiders was 0.19 cm/week or 0.03 cm/day. The results of measuring coral growth rate can be seen in Figure 6.



Figure 6 shows that the transplanted Acropora sp. growth rates were 0.17 cm/week or 0.02 cm/day, 0.25 cm/week or 0.04 cm/day, and 0.34 cm/week or 0.05 cm/day on the second, third, and fourth observations, consecutively. The results showed that the coral growth rate of Acropora sp., transplanted using the spider method, was faster than Subhan (2019) results in Bali using the same coral genus, method, and depth. Where the growth rate of corals obtained was 0.09 cm/week in observation II, and 0.14 cm/week in observation III and IV. The results showed that the coral growth rate of Acropora sp., using the spider method in Botutonuo waters, was faster than Subhan (2019) result in Bali, which also uses the same method, genus and depth, where the coral growth rate obtained by Subhan (2019) was 0.09 cm/week in observation II and 0.14 cm/week in observations III and IV.

The use of the spider transplant modules in this study also showed a better growth rate performance compared to the use of the concrete block transplant module in Botutonuo waters by Alim (2019), where the growth rates on the blocks were 0.01 cm/day in the second and third week of observation, and 0.03 cm/day at the fourth week of observation. The research informs that applying the spider transplant module is more effective in rehabilitating coral reefs and generates growth faster than the concrete beam transplant. Although to see to what extent its effectiveness and development still needs to be investigated through further observation and continuous monitoring.

Coral survival rate using spider modules. The analysis of coral survival rates using the spider module can be seen in Figure 7.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

2450

Commented [GCM28]: Rephrase. Is this from Subhan (2019) research? If so, rephrase so that the citation is appropriate. Use Reply

Commented [SNH29R28]: I have corrected the sentence and added the source



Figure 7. The survival rate of Acropora sp. for two months of observation.

Figure 7 shows that during the two months of observation, there were no dead coral fragments or coral fragments that had escaped from the spider module. Figure 7 shows that no dead coral fragments or coral fragments had escaped from the spider module during the two months of observation. The total number of coral fragments observed at the beginning and the end of the observation amounted to 60 fragments. The results of this study indicate that the use of spider modules for coral transplantation is successful. Harriot and Fisk (1988) in Mompala et al (2017) stated that transplantation of corals was considered successful when the survival rate was in the range of 50–100%. The analysis of the coral survival rate in this study was higher than the results of Subhan's research (2019). There was a decrease in the survival rate of corals using the spider method for a depth of 6 meters, which was only 82% at the end of the study. In this study, the survival rate of transplanted corals in Botutonuo waters was 100%, and the results of this study were higher than the results of Subhan's (2019) study, which was only 82%.

Similarly, the results of research conducted by Rani et al (2017) with the same method showed a coral survival rate of 95.56% at the end of the study. The study results indicate that *Acropora* sp. can grow well in Botutonuo waters. Williams et al (2019) reported that transplanting using the spider method succeeded in restoring the condition of coral reefs in the rehabilitation area.

The main factor causing the high survival rate in this study lies in the initial transplantation treatment, starting from finding and tying seedlings to mobilizing spider modules to the research location, being carried out carefully, and trying not to be exposed to open air, thereby minimizing stress levels on corals.

Water quality parameters. Water quality parameters are one of the essential factors that play a role in the survival of aquatic organisms, especially coral reefs. It causes the distribution of coral reefs in water areas to be uneven. The in-situ measurements of water quality parameters can be seen in Table 1.

Results of measurement of water quality parameters

| Та | ble | 1 |
|----|-----|---|
| | | _ |

sentence

| | | Obser | vation | Seawater Quality Standards for Coral | |
|------------------|------|-------|--------|---|--|
| Parameters | Ι | II | III | IV | Reefs (Minister of Environment Decree No. 51/2004) |
| Temperature (°C) | 29 | 29 | 29 | 29.3 | 28-30 |
| Salinity (‰) | 29 | 29 | 25.5 | 27.7 | 33-34 |
| рН | 7.98 | 8.01 | 7.89 | 7.57 | 7-8.5 |

2451

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl Commented [GCM30]: Although this is cited in Mompala et al (2017), you need to add this to the reference list also. Please add. Commented [SNH31R30]: I've added to reference

Commented [GCM32]: Rephrase. Again, not clear if this is from Subhan (2019) research. Use Reply and answer me. Commented [SNH33R32]: I have highlighted the corrected

| DO (mg/l) | 2.5 | 6.09 | 3.99 | 5.92 | >5 |
|------------------|------|-------|------|------|-------|
| Current (m/s) | 0.02 | 0.015 | 0.02 | 0.03 | - |
| Transparency (m) | 6 | 6 | 6 | 6 | > 5 m |

Table 1 shows that the temperature values in Botutonuo waters are in the range of 29–29.3°C. Supriharyono (2000) stated that a suitable temperature for coral growth ranges from 25–29°C. However, coral animals can tolerate temperatures with a minimum of 16°C and a maximum of 36°C (Kinsman 1964). Salinity ranges between 25.5–29 ‰. Low salinity value is thought to be influenced by rainfall due to rain during the research. In addition, low salinity is also predicted to be affected by the supply of fresh water from the rivers around the study site. However, it is a rainfed river, not a continuous stream. The results of pH measurements at the study sites were in the range of 7.89–8.01. This value is in the normal pH range in Indonesian waters and the range of seawater quality standards for coral reefs. Rizki et al (2011) found that the annual pattern of pH in Indonesia varies from 7.7 to 8.2.

The dissolved oxygen (DO) measurements at the research location were in the range 2.5-6.05 mg/L. The range of DO for the first and second observations is below the standard quality for coral reefs based on Minister of Environment Decree No. 51/2004. However, according to Swingle (1968) in Salmin (2005), the minimum DO of 2 mg/L is sufficient to support the life of organisms.

The current velocity obtained at the research location ranging from 0.015 to 0.02 m/s is in a low category when referring to Yusuf et al (2012), which is below 0.5 m/s. The slow movement of the current is due to the morphology of the bay-shaped location so that the current velocity is low. The results of measuring the transparency of Botutonuo waters obtained of 100% or 6 m, as the spider module was placed at a depth of 6 m. Based on the seawater quality standards for coral reefs, the water transparency in Botutonuo is still in the range of quality standard values.

Conclusions. To sum up, absolute coral growth for two months of observation showed a positive result of 1.52 cm. The growth rate of the transplanted corals during the study was 0.19 cm/week or 0.03 cm/day. The 100% survival rate is included in the transplant category with a high success rate. The quality of Botutonuo waters as a transplantation location using spider modules still supports the growth of transplanted corals.

Acknowledgments. We express our gratitude to LP2M UNG for the reliance given to us to conduct this research through PNBP UNG funding. Our thanks also go to Hidayat Bachtiar, S. Kel., Alinton Pisuna, Dadang S. Alim, S.Pi., Jufriyanto Umar, S.Pi., Ronal Lahmutu for their assistance during data collection and Nuralim Pasisingi, S.Pi., M.Si., for her assistance in writing.

References

- Alim D. S., 2019 [Analysis of growth rate and survival of transplanted corals in Botutonuo Beach waters, Bone Bolango Regency, Gorontalo Province]. Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Gorontalo State University, Gorontalo [in Indonesian].
- Edward A. J., Gomez E. D., 2007 Reef restoration concepts and guidelines: making sensible management choices in the face of uncertainty. The Coral Reef Targeted Research & Capacity Building for Management Programme: St Lucia, Australia, 38 p.
- Effendie M. I., 1979 [Fisheries biological methods]. Yayasan Dewi Sri, Bogor, 112p [in Indonesian].
- Giyanto, Abrar M., Hadi T. A., Budiyanto A., Hafizt M., Salatalohy A., Iswari M. Y., 2017 [Status of Indonesia's coral reef 2017]. Oceanography Research Center -Indonesian Institute of Sciences [in Indonesian].

Harriot, V. J., Fisk, D. A., 1988 Coral transplantation as reef management option. Proceedings of the 6th International Coral Reef Symposium 2:375-379.

Kinsman J. J., 1964 Reef coral tolerance to high temperatures and salinities. Nature 202:1280-1282.

2452

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl Commented [GCM34]: Just like the reference situation before, you need to add this also to the reference list. Commented [SNH35R34]: I've added to reference

- Minister of Environment, 2004 [Decree of the Minister of Environment No. 51 year 2004 about water quality standard for coastal waters]. Minister of Environment of the Republic of Indonesia, Jakarta [in Indonesian].
- Mompala K., Rondonuwu A. B., Rembet U. N. W. J., 2017 [The growth rate of *Acropora* sp. transplanted on artificial reefs in Kareko Waters of North Lembeh Sub-District of Bitung City]. Jurnal Ilmiah Platax 5(2):234–242 [in Indonesian].
- Nedimyer K., Gaines K., Roach S., 2011 Coral Tree Nursery©: An innovative approach to growing corals in an ocean-based field nursery. AACL Bioflux 4(4):442-446.
- Prameliasari R. T. A., Munasik, Wijayanti D. P., 2012 [The effect of fragment size differences and transplantation methods on the growth of *Pocillopora damicornis* corals in Awur Bay, Jepara]. Journal of Marine Research 1(1):159-168 [in Indonesian].
- Rani C., Tahir A., Jompa J., Faisal A., Yusuf S., Werorilangi S., Arniati, 2017 [The success of rehabilitation of coral reefs due to the 2016 bleaching incident with the transplant technique]. Spermonde 3(1):13–19 [in Indonesian].
- Rizki T. Y., Tito C. K., Setiawan A., 2011 [pH variations in Indonesian waters]. https://www.researchgate.net/profile/Camellia_Tito/publication/262673263_Variasi _pH_di_Perairan_Indonesia/links/5645a1d908aef646e6cca454/Variasi-pH-di-Perairan-Indonesia.pdf [in Indonesian].
- Rusandi A., Suharsono, Wiadnyana N., Yusri S., 2016 [National action plan for coral conservation period 1: 2017-2021]. Directorate of Conservation and Marine Biodiversity, Directorate General of Marine Spatial Management, Ministry of Marine Affairs and Fisheries [in Indonesian].
- Salmin, 2005 [Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD) as indicators for determining water quality]. Oseana XXX(3):21–26 [in Indonesian].
- Subhan M. A., 2019 [Growth rate of *Acropora loripes* coral reef using transplantation method of spider skeleton module in the waters of Les Village, Buleleng Regency, Bali]. Universitas Islam Negeri Syarif Hidayatullah [in Indonesian].
- Suharsono, Siringoringo R. M., Hadi T. A., Giyanto, Tuti Y., Budiyanto A., Sulha S., 2013 [Development of coral transplant techniques in Indonesia]. Oceanography Research Center - Indonesian Institute of Sciences, Jakarta [in Indonesian].
- Supriharyono, 2000 [Coral reef ecosystem management]. Djambatan, Jakarta, 108 p [in Indonesian].

Swingle, H. S., 1968 Standardization of chemical analysis for water and pond muds. FAO Fish, Rep. 44, 4: 379-406.

- Williams S. L., Sur C., Janetski N., Hollarsmith J. A., Rapi S., Barron L., Heatwole S. J., Yusuf A. M., Yusuf S., Jompa J., Mars F., 2019 Large-scale coral reef rehabilitation after blast fishing in Indonesia: Coral reef rehabilitation. Restoration Ecology, 27(2):447–456.
- Yusuf M., Handoyo G., Wulandari S. Y., 2012 [Characteristics of flow patterns in relation to water quality conditions and abundance of phytoplankton in the waters of the Karimun Jawa Marine National Park]. Buletin Oseanografi Marina 1:63-74 [in Indonesian].

Received: 12 September 2020. Accepted: 02 October 2020. Published online: xxxx 2021. Authors:

[Sri Nuryatin Hamzah, Department of Aquatic Resources Management, Gorontalo State University, Jl. Jendral Sudirman, No. 6, 96128, Gorontalo City, Indonesia, e-mail: sri.nuryatin@ung.ac.id]

[Sitti Nursinar, Department of Aquatic Resources Management, Gorontalo State University, Jl. Jendral Sudirman, No. 6, 96128, Gorontalo City, Indonesia, e-mail: sitti.nursinar@ung.ac.id]

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. How to cite this article:

[Hamzah, S. N., Nursinar, N.,] 2021 The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency). AACL Bioflux 14(X):xxxx-xxxx.

Commented [GCM36]: Fill in according to the instructions in the parentheses.

Commented [GCM37]: Fill in according to the instructions in the parentheses.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl



Final Review



The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency) ¹Sri N. Hamzah, ¹Sitti Nursinar

¹ Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Gorontalo City, Indonesia. Corresponding author: S. N. Hamzah, sri.nuryatin@ung.ac.id

Abstract. Botutonuo Waters, Bone Bolango Regency, is an area that has coral reef potential for marine tourism practices. However, high human activity such as destructive fishing practices and increasing population in coastal area causes the coral reef ecosystem to be stressed and damaged. One of the alternatives that can be implemented to protect and conserve coral reefs is through coral transplantation. This study was conducted to provide information about the success of transplantation using the spider module. The experiment was conducted for two months by placing ten spider modules in the water. A total of 60 coral fragments were used where six coral fragments were tide to each module. Coral growth monitoring was carried out every two weeks for four times. The results revealed that the absolute growth of corals for two months of observation showed positive results, namely 1.52 cm. The growth rate of the transplanted corals was 0.19 cm/week or 0.03 cm/day. The survival rate of transplanted corals was 100% and is included in the category of high transplant success rate. **Key Words**: Botutonuo waters, coral reef, growth, spider, survival rate.

Introduction. Coral reefs are one of the ecosystems that are crucial for the sustainability of existing resources in coastal areas and oceans. The existence of this ecosystem provides many benefits not only for the biota association but also for humans. Rusandi et al (2016) stated that ecologically, coral reefs function as a habitat for various reef fish caught by fishermen, as a coast protector from waves, a source of germplasm, and economically as a source of income through marine tourism and ornamental coral trades. Currently, the rate of exploitation of coral reefs in not as fast as the rate of regeneration, so that has an impact on damage to coral reefs in coastal areas. Giyanto et al (2017) reported that the current condition of coral reefs in Indonesia was 36.18% in the bad category, 34.3% in the moderate category, 22.96% in the good category, and only 6.55% in the yery good category. In general, the trend of the condition of locear leves in the coastal environment. Even some research results indicate that coastal areas with damaged coral reefs have the potential to receive bigger waves and are threatened with coastal abraision.

damaged coral reefs have the potential to receive bigger waves and are threatened with coastal abrasion. Botutonuo waters are one of the waters in the Tomini Bay area of Goronalo. Currently, the Botutonuo water area is used for various activities such as fishing and marine tourism. Information obtained from the coastal community of Botutonuo shows that in 1980-1990 fishing activities were carried out by destructive methods in this area. The amount of rubble was found to be an indicator of destructive fishing in the past. If no measures are taken to restore this condition, the damage to coral reefs can get worse. Restoration of coral reef ecosystems can occur naturally. However, the time required is quite long, so interventions are needed to restore the condition of coral reefs in numerous locations in a short time. Several methods have been implemented to restore coral reef conditions in a shorter time, including artificial reef methods, mineral accretion (bio rock), and coral

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

transplantation (Suharsono et al 2013). Coral transplantation is a method of coral cultivation by cutting a portion of a particular coral colony to be planted in a new place (Prameliasari et al 2012). Transplantation through asexual fragmentation is essential as an effort to reproduce corals for recovery, trade, and marine tourism (Nedimyer et al 2011; Suharsono et al 2013). Transplantation using spider modules has never been tested in Gorontalo Province since this method is the newest and not widely used in Indonesia. It was identified that transplanting with the spider module was only found in South Sulawesi and Bali (Williams et al 2019; Subhan 2019). Therefore, there is a lack of information about coral growth using the spider module. This study aims to provide information about coral growth and survival rates using

This study aims to provide information about coral growth and survival rates using the spider module as well as a reference in selecting a transplant model that is cheap and easy to apply in the field.

Material and Method

Study site. This research was conducted in May-July 2020 at Botutonuo waters, Bone Bolango Regency. The research location is depicted in Figure 1.



Figure 1. Location of coral transplantation using spider modules.

Preparation and module construction. The placement of the spider module was determined in the initial observation, namely through the manta tow survey, to see the condition of the coral reefs around the Botutonuo waters. The spider module was placed on the bottom of the water where damaged coral reef conditions were observed. The spider construction can be seen in Figure 2.



Figure 2. Spider module construction (A); illustration of coral fragments tied to spider modules (B).

The spider construction was made of steel with a diameter of 12 mm, which was shaped to resemble a spider's web and legs (Figure 2). The frame height is \pm 50 cm, with the bottom diameter of \pm 120 cm, and the distance among the legs is \pm 60 cm. Each spider frame was coated with sand, which is glued using resin and serves to prevent rust from the iron when the module was lowered into the water.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

Seed procurement, fragmentation, and fragment binding. Coral seedlings were obtained from broodstock nearby the transplant site. The seedlings were taken by cutting the parent corals from the site using pliers and scissors into small colonies (fragments) with a size range of 3-12 cm for branching corals (Rani et al 2017; Suharsono et al 2013). The coral fragments were then put in baskets and soaked with seawater to keep the coral fragments from stress. The coral taken was fragments collected were then tied using cable ties to the spider frame, as many as 6 fragments for each module, so that a total of 60 fragments.

Measurement of growth and survival rates of coral fragments. Coral fragment growth measurements were carried out every two weeks for two months. At the start of rearing, transplanted corals were measured as initial length using a caliper. Furthermore, every two weeks, the increase in branch length in the waters was measured manually. Also, the number of dead or missing fragments in each experimental unit was counted to determine the survival rate. During the measurement of coral growth, cleaning of the module area, which was overgrown with algae, was also carried out to prevent the death of coral fragments due to algal dominance.

Measurement of water quality parameters. Water quality parameters measured in this study were temperature, salinity, and dissolved oxygen (DO) using a Lutron DO-5510, pH using pH Meter AS-218, current using Flowatch Meter FL-03, and transparency using Secchi Disk. Measurement of the parameters was performed in situ along with the measurements of coral growth.

Data analysis. The results of coral reef measurements were tabulated and analyzed. Some of the analysis parameters are the following:

1. Absolute growth Coral growth over a period was calculated using Effendie (1979) formula:

 $\beta = Lt - Lo$

where: β = absolute growth (cm); Lt = the average branch length after the tth observation; Lo = average branch length at the beginning of research.

2. Coral growth rate

Coral Growth Rate was calculated using Effendie (1979) formula:

 $\mathbf{P} = \frac{Lt - L0}{t}$

where: P = increase in length/height of transplanted coral fragments; Li+1 = the mean length/height of the fragments after the tth observation; L0 = the mean length/height of the initial fragments; t = observation time.

Survival rate

The formula (Effendi 1979) used for the coral survival rate is:

$S = \frac{Nt}{N_0} \ge 100\%$

where: S = survival rate (%); Nt = Number of coral fragments at the end of the observation; and No = Number of coral pieces at the beginning of observation.

AACL Bioflux, 2021, Volume 14, Issue X. 2447 http://www.bioflux.com.ro/aacl

Results and Discussion

The range of coral lengths at the beginning of the study was 8.67-10.08 cm, with an average of 9.36 cm. The measurement results showed that there was coral growth during the two months of observation (Figure 3).



Figure 3. Coral growth using spider modules

Figure 3 shows that there was an increase in coral growth at the research location every two weeks of measurement for two months of field observation. The average size of the initial coral fragments was 9.36 cm and at the end of the study, in the fourth observation, the length was 10.88 cm. The results designated that the absolute growth of the transplanted corals was 1.52 cm. The absolute growth of *Acropora* sp., which was transplanted using this spider, was higher than that of Subhan (2019) and Rani et al (2017), which were 0.59 cm and 0.1-0.2 cm, respectively. The results showed that the coral growth observation experiment in the Botutonuo waters shows better performance than previous studies using the same transplant module. Furthermore, absolute growth in this study is better than the results obtained by Alim (2019) in the same research location with the concrete beam transplant module. Alim (2019) reported that the absolute growth of cral during two months of observation was 0.6 cm. Based on visual robservation, the transplanted coral began to stick to the spider module, and some coral fragments had started to wrap the ties (Figure 4).

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl



Observation I (Week 0)

Observation II (Week 2)



Figure 4. Observation of growth and morphological changes in transplanted corals using spider modules.



Figure 5. Coral fragments attached to observations II and III (week 2 and week 4).

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

Figure 4 shows changes in the morphology of the transplanted corals, both changes in the height increase of corals and the adding of new coral branches. The life of coral fragments in the transplant module can be observed through the tie cables and the attachment of elements (Figure 5) to the substrate. Coral fragments are categorized as living on the transplant media when they are attached or when the binding cables have been wrapped by new coral tissue at the end of the transplant. According to Edward and Gomez (2007), corals that have stuck to and covered the substrate are less likely to escape.

Growth rate. In general, the growth rate of corals transplanted using spiders was 0.19 cm/week or 0.03 cm/day. The results of measuring coral growth rate can be seen in Figure 6.



Figure 6 shows that the transplanted *Acropora* sp. growth rates were 0.17 cm/week or 0.02 cm/day, 0.25 cm/week or 0.04 cm/day, and 0.34 cm/week or 0.05 cm/day on the second, third, and fourth observations, consecutively. The results showed that the coral growth rate of *Acropora* sp., using the spider method in Botutonuo waters, was faster than Subhan (2019) result in Bali, which also uses the same method, genus, and depth, where the coral growth care obtained by Subhan (2019) was 0.09 cm/wask in

where the coral growth rate obtained by Subhan (2019) was 0.09 cm/week in observation II and 0.14 cm/week in observations III and IV. The use of the spider transplant modules in this study also showed a better growth rate performance compared to the use of the concrete block transplant module in Botutonuo waters by Alim (2019), where the growth rates on the blocks were 0.01 cm/day in the second and third week of observation, and 0.03 cm/day at the fourth week of observation. The research informs that applying the spider transplant module is more effective in rehabilitating coral reefs and generates growth faster than the concrete beam transplant. Although to see to what extent its effectiveness and development still needs to be investigated through further observation and continuous monitoring.

Coral survival rate using spider modules. The analysis of coral survival rates using the spider module can be seen in Figure 7.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl



Figure 7. The survival rate of Acropora sp. for two months of observation

Figure 7 shows that no dead coral fragments or coral fragments had escaped from the spider module during the two months of observation. The total number of coral fragments observed at the beginning and the end of the observation amounted to 60 fragments. The results of this study indicate that the use of spider modules for coral transplantation is successful. Harriot and Fisk (1988) in Mompala et al (2017) stated that transplantation of corals was considered successful when the survival rate was in the range of 50-100%. In this study, the survival rate of transplanted corals in Botutonuo waters was 100%, and the results of this study were higher than the results of Subhan's (2019) study, which was only 82%.

(2019) study, which was only 82%. Similarly, the results of research conducted by Rani et al (2017) with the same method showed a coral survival rate of 95.56% at the end of the study. The study results indicate that *Acropora* sp. can grow well in Botutonuo waters. Williams et al (2019) reported that transplanting using the spider method succeeded in restoring the condition of coral reefs in the rehabilitation area.

The main factor causing the high survival rate in this study lies in the initial transplantation treatment, starting from finding and tying seedlings to mobilizing spider modules to the research location, being carried out carefully, and trying not to be exposed to open air, thereby minimizing stress levels on corals.

Water quality parameters. Water quality parameters are one of the essential factors that play a role in the survival of aquatic organisms, especially coral reefs. It causes the distribution of coral reefs in water areas to be uneven. The in-situ measurements of water quality parameters can be seen in Table 1.

Table 1

| Results of | measurement | of | water | quality | parameters |
|------------|-------------|----|--------------|---------|------------|
| results of | medaaremene | | HACCO | quanty | parameters |

| | Observation | | | | Seawater Quality | | |
|------------------|-------------|-------|------|------|--|--|--|
| Parameters | I | Ш | III | IV | Reefs (Minister of Environment Decree No. 51/2004) | | |
| Temperature (°C) | 29 | 29 | 29 | 29.3 | 28-30 | | |
| Salinity (‰) | 29 | 29 | 25.5 | 27.7 | 33-34 | | |
| pH | 7.98 | 8.01 | 7.89 | 7.57 | 7-8.5 | | |
| DO (mg/l) | 2.5 | 6.09 | 3.99 | 5.92 | >5 | | |
| Current (m/s) | 0.02 | 0.015 | 0.02 | 0.03 | - | | |
| Transparency (m) | 6 | 6 | 6 | 6 | > 5 m | | |

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl

Table 1 shows that the temperature values in Botutonuo waters are in the range of 29–29.3°C. Supriharyono (2000) stated that a suitable temperature for coral growth ranges from 25–29°C. However, coral animals can tolerate temperatures with a minimum of 16°C and a maximum of 36°C (Kinsman 1964). Salinity ranges between 25.5–29 ‰. Low salinity value is thought to be influenced by rainfall due to rain during the research. In addition, low salinity is also predicted to be affected by the supply of fresh water from the rivers around the study site. However, it is a rainfed river, not a continuous stream. The results of pH measurements at the study sites were in the range of 7.89–8.01. This value is in the normal pH range in Indonesian waters and the range of seawater quality standards for coral reefs. Rizki et al (2011) found that the annual pattern of pH in Indonesia varies from 7.7 to 8.2.

The dissolved oxygen (DO) measurements at the research location were in the range 2.5-6.05 mg/L. The range of DO for the first and second observations is below the standard quality for coral reefs based on Minister of Environment Decree No. 51/2004. However, according to Swingle (1968) in Salmin (2005), the minimum DO of 2 mg/L is sufficient to support the life of organisms.

The current velocity obtained at the research location ranging from 0.015 to 0.02 m/s is in a low category when referring to Yusuf et al (2012), which is below 0.5 m/s. The slow movement of the current is due to the morphology of the bay-shaped location so that the current velocity is low. The results of measuring the transparency of Botutonuo waters obtained of 100% or 6 m, as the spider module was placed at a depth of 6 m. Based on the seawater quality standards for coral reefs, the water transparency in Botutonuo is still in the range of quality standard values.

Conclusions. To sum up, absolute coral growth for two months of observation showed a positive result of 1.52 cm. The growth rate of the transplanted corals during the study was 0.19 cm/week or 0.03 cm/day. The 100% survival rate is included in the transplant category with a high success rate. The quality of Botutonuo waters as a transplantation location using spider modules still supports the growth of transplanted corals.

Acknowledgments. We express our gratitude to LP2M UNG for the reliance given to us to conduct this research through PNBP UNG funding. Our thanks also go to Hidayat Bachtiar, S. Kel., Alinton Pisuna, Dadang S. Alim, S.Pi., Jufriyanto Umar, S.Pi., Ronal Lahmutu for their assistance during data collection and Nuralim Pasisingi, S.Pi., M.Si., for her assistance in writing.

References

- Alim D. S., 2019 [Analysis of growth rate and survival of transplanted corals in Botutonuo Beach waters, Bone Bolango Regency, Gorontalo Province]. Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Gorontalo State University, Gorontalo [in Indonesian].
- Edward A. J., Gomez E. D., 2007 Reef restoration concepts and guidelines: making sensible management choices in the face of uncertainty. The Coral Reef Targeted Research & Capacity Building for Management Programme: St Lucia, Australia, 38

Effendie M. I., 1979 [Fisheries biological methods]. Yayasan Dewi Sri, Bogor, 112p [in Indonesian].

Giyanto, Abrar M., Hadi T. A., Budiyanto A., Hafizt M., Salatalohy A., Iswari M. Y., 2017 [Status of Indonesia's coral reef 2017]. Oceanography Research Center -Indonesian Institute of Sciences [in Indonesian].

Harriot V. J., Fisk D. A., 1988 Coral transplantation as reef management option. Proceedings of the 6th International Coral Reef Symposium 2:375-379.

Kinsman J. J., 1964 Reef coral tolerance to high temperatures and salinities. Nature 202:1280-1282.

AACL Bioflux, 2021, Volume 14, Issue X. 2452 http://www.bioflux.com.ro/aacl

- Minister of Environment, 2004 [Decree of the Minister of Environment No. 51 year 2004 about water quality standard for coastal waters]. Minister of Environment of the Republic of Indonesia, Jakarta [in Indonesian].
- Mompala K., Rondonuwu A. B., Rembet U. N. W. J., 2017 [The growth rate of Acropora sp. transplanted on artificial reefs in Kareko Waters of North Lembeh Sub-District of Bitung City]. Jurnal Ilmiah Platax 5(2):234–242 [in Indonesian]. Nedimyer K., Gaines K., Roach S., 2011 Coral Tree Nursery©: An innovative approach to
- growing corals in an ocean-based field nursery. AACL Bioflux 4(4):442-446. Prameliasari R. T. A., Munasik, Wijayanti D. P., 2012 [The effect of fragment size differences and transplantation methods on the growth of *Pocillopora damicornis* corals in Awur Bay, Jepara]. Journal of Marine Research 1(1):159-168 [in Indonesian].
- Rani C., Tahir A., Jompa J., Faisal A., Yusuf S., Werorilangi S., Arniati, 2017 [The success of rehabilitation of coral reefs due to the 2016 bleaching incident with the transplant technique]. Spermonde 3(1):13–19 [in Indonesian].
 Rizki T. Y., Tito C. K., Setiawan A., 2011 [pH variations in Indonesian waters].
- https://www.researchgate.net/profile/Camellia_Tito/publication/262673263_Variasi _pH_di_Perairan_Indonesia/links/5645a1d908aef646e6cca454/Variasi-pH-di-Perairan-Indonesia.pdf [in Indonesian].
- Rusandi A., Suharsono, Wiadnyana N., Yusri S., 2016 [National action plan for coral conservation period 1: 2017-2021]. Directorate of Conservation and Marine Biodiversity, Directorate General of Marine Spatial Management, Ministry of Marine Affairs and Fisheries [in Indonesian].
- Salmin, 2005 [Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD) as indicators for determining water quality]. Oseana XXX(3):21-26 [in Indonesian].
 Subhan M. A., 2019 [Growth rate of Acropora loripes coral reef using transplantation
- method of spider skeleton module in the waters of Les Village, Buleleng Regency, Bali]. Universitas Islam Negeri Syarif Hidayatullah [in Indonesian].
- Suharsono, Siringoringo R. M., Hadi T. A., Giyanto, Tuti Y., Budiyanto A., Sulha S., 2013 [Development of coral transplant techniques in Indonesia]. Oceanography Research Center - Indonesian Institute of Sciences, Jakarta [in Indonesian].
- Supriharyono, 2000 [Coral reef ecosystem management]. Djambatan, Jakarta, 108 p [in Indonesian].
- Swingle H. S., 1968 Standardization of chemical analysis for water and pond muds. FAO Fish, Rep. 44, 4: 379-406.
- Williams S. L., Sur C., Janetski N., Hollarsmith J. A., Rapi S., Barron L., Heatwole S. J., Yusuf A. M., Yusuf S., Jompa J., Mars F., 2019 Large-scale coral reef rehabilitation after blast fishing in Indonesia: Coral reef rehabilitation. Restoration Ecology, 27(2):447-456.
- Yusuf M., Handoyo G., Wulandari S. Y., 2012 [Characteristics of flow patterns in relation to water quality conditions and abundance of phytoplankton in the waters of the Karimun Jawa Marine National Park]. Buletin Oseanografi Marina 1:63-74 [in Indonesian].

Received: 12 September 2020. Accepted: 02 October 2020. Published online: xxxx 2021. Authors: Sri Nuryatin Hamzah, Department of Aquatic Resources Management, Gorontalo State University, JL Jendral Sudirman, No. 6, 96128,

Gorontalo City, Indonesia, e-mail: sri.nuryatin@ung.ac.id Sitti Nursinar, Department of Aquatic Resources Management, Gorontalo State University, Jl. Jendral Sudirman, No. 6, 96128, Goro

Site Nurshar, Department of Aquate Resources Management, before the University, JL Jeneral Sourmain, No. 6, 99126, Gorontalo (Cty, Indonesis, e-mail: site).nurshar@ung.acid This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. How to cite this article:

Hamzah S. N., Nursinar N., 2021 The success of coral rehabilitation through transplantation using spider modules (case study: Botutonuo marine area, Bone Bolango Regency). AACL Bioflux 14(X):xxxx-xxxx.

AACL Bioflux, 2021, Volume 14, Issue X. http://www.bioflux.com.ro/aacl 2453



Publish:

Link http://www.bioflux.com.ro/docs/2021.3023-3031.pdf



Search Keywords Search

You are here > Home · Volume 14(5)/2021 Volume 14(5)/2021 AACL Bioflux First pages, 2021, AACL Bioflux 14(5):i-viii. Instructions to authors Submission letter Purwanto Y., Santoso H., Manohas J., Zaini M., Tumiwa J. H., Nugraha E., 2021 Analysis of the diversity index and dominance of bottom gillnet catches in Kulu Model of paper waters, North Minahasa Regency, Indonesia. AACL Bioflux 14(5):2639-2649. Reviewer information pack Tao C. T., Truyen P. M., Khanh L. V., Terahara T., Hai T. N., Hoa N. V., 2021 The Editorial Board Expanded influence of feeding reduction on the survival and growth of the larvae and postlarvae of freshwater prawn (Macrobrachium rosenbergii): an applied biofloc Coverage / databases technology study. AACL Bioflux 14(5):2650-2661. Volume 16(6)/2023 (December, 30) Volume 16(5)/2023 (October, 30) Tapilatu J. R., Siburian R. H. S., Tapilatu M. E., 2021 Species identification, density, and type of substrate of clam (Tridacnidae) in Kali Lemon coastal water -Volume 16(4)/2023 (August, 30) Kwatisore, Cenderawasih Bay, Papua, Indonesia. AACL Bioflux 14(5):2662-2671. Volume 2(1)/2009 (January, 30) Heriyati E., Rustadi, Isnansetyo A., Triyatmo B., Istiqomah I., 2021 Microbubble aerator test and harvest target prediction based on oxygen consumption of red Volume 1(2)/2008 (December, 30) tilapia (Oreochromis sp.). AACL Bioflux 14(5):3006-3022. Volume 1(1)/2008 (September, 30) Hamzah S. N., Nursinar N., 2021 The success of coral rehabilitation through Volume Pilot/2007 (December, 30) transplantation using spider modules (case study: Botutonuo marine area, Bone available printed only Bolango Regency). AACL Bioflux 14(5):3023-3031. Pontus Euxinus, Volume 1 (1980) -Mohammad A. S., Farrag M. M. S., Mustafa A. A., Osman A. G. M., 2021 Parent Journal Ultrastructure and morphometry of the sagittal otolith as confirmatory identification in three carangid species from the northern Red Sea, Egypt. AACL AACL Bioflux Bioflux 14(5):3032-3044 Ecology, Evolution, Behavior and Prawito P., Mulvasari G., 2021 Comparative social vulnerability of fishermen in the coastal area of Bengkulu and Central Java, Indonesia. AACL Bioflux 14(5):3045-Systematics 3054. SIR 2021 Dahlia, Ardiansyah, Indrayani, Jabbar F. B. A., 2021 The abundance and diversity 0.26 of phosphate-solubilizing bacteria in integrated recirculating aquaculture powered by scimagojr.com systems as a function of harvesting regime of duckweed (Lemna minor L.). AACL Bioflux 14(5):3055-3067.

> Danish-Daniel M., Saari N. A., Okomoda V. T., Musa N., Noor M. E. M., Pau T. M., lehata S., Fadaak S. N. E. S. J., Sung Y. Y., Usup G., 2021 Culture-dependent and 16S metagenomics analyses of blue-bell tunicate *Clavelina coerulea* microbiome. AACL Bioflux 14(5):3068-3091.