

Covid-19 forecast using Holt-Winters exponential smoothing

by Sri Endang Saleh

Submission date: 21-Jun-2023 07:43PM (UTC+0800)

Submission ID: 2120281010

File name: Djakaria_2021_J._Phys._Conf._Ser._1882_012033.pdf (809.18K)

Word count: 3050

Character count: 16406

PAPER • OPEN ACCESS

Covid-19 forecast using Holt-Winters exponential smoothing

To cite this article: I Djakaria and S E Saleh 2021 *J. Phys.: Conf. Ser.* **1882** 012033

View the [article online](#) for updates and enhancements.

You may also like

- Administered Machine Learning Models for Covid-19 Future Forecasting K Atchaya, M Darshinii, R Harini et al.
- The autoregressive integrated vector model approach for covid-19 data in nesia and Singapore R Fitriani, W D Revildy, E Marhamah et al.
- Weakly-supervised lesion analysis with a CNN-based framework for COVID-19 Kaichao Wu, Beth Jeffs, Xiangyuan Ma et al.

The advertisement features a dark blue background. On the left, a hand is shown reaching out to touch a glowing globe. The globe is surrounded by a network of white icons representing people and connections. To the right of the globe is the ECS logo, which consists of the letters 'ECS' inside a white circle. Further right, the text 'Connect with decision-makers at ECS' is written in a bold, yellow font. Below this, in a smaller white font, is the text 'Accelerate sales with ECS exhibits, sponsorships, and advertising!'. At the bottom right, a yellow play button icon is followed by the text 'Learn more and engage at the 244th ECS Meeting!' in a yellow font.

Connect with decision-makers at ECS

Accelerate sales with ECS exhibits, sponsorships, and advertising!

▶ Learn more and engage at the 244th ECS Meeting!

1 Covid-19 forecast using Holt-Winters exponential smoothing

I Djakaria¹ and S E Saleh²

¹ Statistics Department, Faculty of Mathematics and Natural Science, Gorontalo State University, Gorontalo – Indonesia

² Development Economics Department, Economics Faculty, Gorontalo State University, Gorontalo – Indonesia

Email: ¹ iskar@ung.ac.id; ² sriendang_s@ung.ac.id

Abstract. Covid-19 has spread throughout the world, including in Indonesia and it is known that the virus is reported to be highly contagious. Indonesia is the fourth-largest population for Covid-19 cases in Asia, while in the world the top three for covid-19 cases are United States, Brazil, and India, so it is inevitable, Indonesia will be greatly felt the spread of the pandemic coronavirus, even estimated to be freed from the coronavirus in a shorter period when compared to other countries with less exposure to COVID-19. In this paper, we study the covid-19 prediction model using Holt-Winters exponential smoothing, for a certain period. This study uses the Covid-19 pandemic data area of Gorontalo, Indonesia, from April 10 to October 13, 2020 (especially total cases). It was found that using Holt-Winters exponential smoothing, the best forecasting model is the one with smoothing parameters $\alpha = 0.1$ and $\gamma = \delta = 0.5$ for trend and seasonality respectively, which gives the smallest MAPE value of 6.14.

1. Introduction

The Covid-19, whose spread was declared by WHO to be a pandemic on March 11, 2020, when there was a spike in the number of cases reported by Italy, Iran, South Korea, and Japan [1]. All people's activities are severely restricted by this pandemic, such as economic activities, religious activities, teaching and learning activities, office activities, and others. National celebrations such as the Population Census (SP2020) must be carried out with the Covid-19 protocol.

Indonesia first confirmed two Covid-19 cases on March 2, 2020, and as of October 17, 2020, has reported more than 357,762 cases in total, or 1,327 cases per million of the population. In Gorontalo's situation, the first confirmed one Covid-19 case on April 10, 2020, and until October 17, 2020 has confirmed 2,964 cases in total, or 2,481 cases per million of the population of Gorontalo. Many studies have been carried out to predict the end of Covid-19. Predicting the turning point, duration, and attack rate of COVID-19 outbreaks in major Western countries has been carried out [3]. Meanwhile, [4] predicted the optimal lockdown period with a parametric approach using a three-phase maturation SIRD model for the COVID-19 pandemic. Time series data is a series of data collected based on time orders at the same intervals. Time series data are widely recorded in various fields such as agriculture, tourism, economy and business, health, and others, including the Covid-19 pandemic. Seasonal patterns can be analyzed by the Holt-Winters exponential smoothing method. The time series method that is often used in forecasting is the Holt-Winters exponential smoothing method. This paper describes the accuracy of the Holt-Winters exponential smoothing time series forecasting model in the Covid-19 pandemic.



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

² The purpose of this study is to find the best forecasting model with the appropriate smoothing parameters. Besides, this study provides predictive or results of forecasting Covid-19 data. It can be shown, either with visual graphs and the result of computation through the existing formula models.

²⁷ Literature review

The Covid-19 pandemic is the most significant global crisis affecting the lives of most of the world's population after the Second World War. Beyond the size and reach of the aftermath of World War, it has affected all the Nations on our planet. The health consequences of a pandemic are devastating. To date, the number of Covid-19 deaths has exceeded 7²¹00 and is sadly destined to grow exponentially in the future [1][5][6]. Thus, urgent action is needed to prevent the spread of the Covid-19 pandemic to the wider community. Currently, there is no vaccine or specific medication ²⁰ COVID-19. Though, several ongoing clinical trials are evaluating potential treatments. The recent spread of COVID-19 has revived the attention of the scientific and political community in the mathematical model of an epidemic. Many researchers are making efforts to suggest new, refined models for examining the current situation and predicting possible future scenarios. The SIR model, where SIR refers to Vulnerable, Infected and Removed, was proposed by Griffiths and Higham [2] which is based on a system of initial value problems of ordinary differential equations (ODE).

In this paper, we describe the methods used to predict the Covid-19 pandemic data using the multiplicative Holt-Winters exponential smoothing. This means the paper wants to answer the question that "what ²⁹ the Covid-19 pandemic data forecasting mode ⁵ using Holt-Winters exponential smoothing?" Holt-Winters exponential smoothing is appropriate if the data is only influenced by trend patterns. However, if the data is not only influenced by trend patterns, but also seasonal patterns, then Holt exponential smoothing is not appropriate for forecasting because it cannot detect any seasonal patterns. Therefore, the Holt-Winters exponential smoothing by adding one parameter to overcome the seasonal patterns in the data. This method, which contains trend and seasonal parameters, is called the Holt-Winters exponential smoothing method. The method used in this study is multiplicative.

2.1. Method of smoothing

In the technique of time series forecasting, it is common for data to show trend patterns, where data show a pattern of an increase or decrease trend. The Holt-Winters exponential smoothing is a method of forecasting using an exponential smoothing approach based on the results of the previous period forecasting. This method adds a parameter that handles seasonal data patterns. The model was chosen based on seasonal patterns ⁵ [8].

The smoothing method is used to forecast the time-series data containing trend patterns, seasonal patterns, or containing both simultaneously. Smoothing is taking the average value over several years to assess the ¹⁴ value of a particular year [9]. The smoothing method is categorized into two parts, i.e the smoothing method and the ¹⁴ exponential smoothing method [17]. Forecasting data that is influenced by seasonal or trend patterns is conducted using the exponential smoothing method by allocating different weights for past data and these weights have exponential decreasing characteristics [17].

2.2. Method of multiplicative Holt-Winters exponential smoothing

The Holt-Winters forecasting algorithm was developed by Charles Holt and Peter Winters. The algorithm smooths time series data and then uses it for forecasts other aspects in the data they concern [11] and [13]. Exponential smoothing is a method for smoothing a time series data that allocates exponentially decreasing weights and values for past data. There are three types of exponential smoothing. The first type is the single exponential smoothing time series forecasting for univariate data. This type is used when the time series data do not have a systematic structure. The data do not have trends and seasonality [10]. According to Panda M [11], this type of exponential smoothing only uses a single parameter α that lies between 0 and 1 as a smoothing factor. A smaller α value indicates slower learning that takes more past observations to estimate. On the other hand, a larger value indicates faster learning that requires more recent observations to make an estimate.

The next type is the double exponential smoothing where in addition to α , another smoothing parameter γ is used for change in trend. There are two types of trends namely additive trend which provides linear trend analysis and multiplication trend which provides exponential trend analysis. It was observed that during the multi-step forecast in the long term, the trend is not a feasible possibility. Therefore, dampening may be practical by reducing the trend size for the future forecast with a straight line (no trend).

Finally, the third type of exponential smoothing is the triple exponential smoothing method. This method is a smoothing technique used when a series shows seasonal variations, that is, the method allows seasonality. The triple exponential smoothing method depends on three parameters that are α , γ , and δ which values lies between zero and one namely $0 < \alpha, \gamma, \delta < 1$ [12].

The Holt-Winters triple exponential smoothing is the newest exponential smoothing method, named after its founders Charles Holt and Peter Winters, which is useful for finding patterns of changing levels, trends, and seasons over time by using additive or multiplicative seasons. In this paper, Holt-Winters triple exponential smoothing [12] is used for forecasting, with the multiplicative model, i.e

- Exponential smoothing of original data (at the time t), (see also [3][6][13][14][15])

$$L_t = \alpha \frac{Y_t}{S_{t-s}} + (1 - \alpha)(L_{t-1} + T_{t-1}) \quad (1)$$

- Trend patterns smoothing at the time t

$$T_t = \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1} \quad (2)$$

- Seasonal patterns smoothing (at the time t)

$$S_t = \delta \frac{Y_t}{L_t} + (1 - \delta)S_{t-s} \quad (3)$$

So, the p -period forecasting forward is

$$\hat{Y}_{t+p} = (L_t + pT_t)S_{t-s+p} \quad (4)$$

where $(0 < \alpha, \gamma, \delta < 1)$.

2.3. Evaluation of forecasting model

The data pattern to be analyzed in a forecasting method plays a significant role. The best forecasting method selection is based on the prediction error rate, if the method is considered a correct method [16]. It is understandable if a forecasting method cannot accurately predict the future state of the data. Therefore each forecasting method produces errors. The smaller the error rate generated by the forecasting model is, the closer the forecasting result will be to the actual result. To calculate the prediction errors, several statistics are used including mean squared deviation (MSD), mean absolute deviation (MAD), and mean absolute percentage error (MAPE). Forecasting using MSD and MAD as a measuring tool for accuracy can cause problems because this measure does not provide a comparison between different time series and different time intervals. MSD and MAD are absolute measures that are highly dependent on the scale of the time-series data. Since MSD and MAD involve the square of a series of values, it is not align with our intuitions. Thus, due to the limitations of MSD and MAD as a measuring tool for forecasting accuracy, we used MAPE as an alternative measure of the accuracy in the forecasting.

3. Methodology

The steps for the forecasting method described above will be clearly stated in this section.

3.1. Data collection technique

The research stages used in this study are as follows.

- Conduct to plot Covid-19 pandemic actual data collection

This study uses the Covid-19 pandemic data area of Gorontalo, Indonesia, between April 10 – October 13, 2020 (especially total cases) [18].

24

- Plotting Covid-19 pandemic time series data using Holt-Winters exponential smoothing.
- Select the smoothing parameter by trial and error which makes the accuracy of the predictions the better.

3.2. Data analysis technique

The analysis of the Covid-19 pandemic data was carried out with the help of *MINITAB 17 Statistical Software* and *Microsoft Excel*.

4. Result and discussion

Plotting of Covid-19 pandemic time series actual cases data, area of Gorontalo, April 10 – October 13, 2020 (especially total cases), Figure 1. This data plot shows that the data is influenced by both trend and increasing patterns. Holt-Winters exponential smoothing method, as stated above, uses three parameters, i.e the level parameter (α), the trend parameter (γ), and the seasonal parameter (δ). Therefore, several forecasting models will be obtained with different parameters. Decide on the best model for predicting the Covid-19 pandemic data using this method can use the MAPE value. The model taken is a model that has the smallest MAPE value. The following are some models for predicting the Covid-19 pandemic data using the multiplicative Holt-Winters exponential smoothing method, by forecasting three times trial and error, respectively, for the first experiment, $\alpha = \gamma = \delta = 0.2$, the MAPE accuracy is 6.48, see Figure 2, the second experiment, $\alpha = 0.1$, $\gamma = 0.2$, $\delta = 0.5$, MAPE is obtained equal to 6.56, like Figure 3, and the third experiment, $\alpha = 0.2$, $\gamma = \delta = 0.5$, with a MAPE value of 6.14 (Figure 4), as below:




Figure 1. Plotting of Covid-19 pandemic time series actual data.




Figure 2. Holt-Winters exponential smoothing for time series of Covid-19 pandemic data. using $\alpha = \gamma = \delta = 0.2$.




Figure 3. Holt-Winters exponential smoothing for time series of Covid-19 pandemic data using $\alpha = 0.1, \gamma = 0.2, \delta = 0.5$.




Figure 4. Holt-Winters exponential smoothing for time series of Covid-19 pandemic data using $\alpha = 0.1$, $\gamma = \delta = 0.5$.

Based on the time series data forecasting plot above (Figure 4), the forecasting model is obtained using the Holt-Winters exponential smoothing method, according to Equations (1), (2), (3), and (4), respectively, which gives the smallest MAPE value (6.14) as follows.

- Exponential smoothing of original data (at the time t),

$$L_t = 0.1 \frac{Y_t}{S_{t-12}} + 0.9(L_{t-1} + T_{t-1}) \quad (5)$$

- Smoothing trend patterns (at the time t)

$$T_t = 0.5(L_t - L_{t-1}) + 0.5T_{t-1} \quad (6)$$

- Smoothing seasonal patterns (at the time t)

$$S_t = 0.5 \frac{Y_t}{L_t} + 0.5S_{t-14} \quad (7)$$

So, the p -period forecasting forward is

$$\begin{aligned} \hat{Y}_{t+p} &= (L_t + pT_t)S_{t+p} \\ &= \left[\left(0.1 \frac{Y_t}{S_{t-14}} + 0.9(L_{t-1} + T_{t-1}) \right) + p(0.2(L_t - L_{t-1}) + 0.8T_{t-1}) \right] \\ &\quad \times \left[0.5 \frac{Y_{t-s+p}}{L_{t-s+p}} + 0.5S_{t-14+p} \right] \end{aligned} \quad (8)$$

5. Conclusion

According to the analysis and discussion of Covid-19 pandemic data forecasting area of Gorontalo from April 10, 2020 to October 13, 2020 (especially total cases), it can be concluded that:

1. The forecasting model using Holt-Winters exponential smoothing is best with smoothing parameters for level $\alpha = 0.1$, trend, and seasonality ($\gamma = \delta = 0.5$), respectively.
2. The trend of confirmed cases of Covid-19 pandemic prediction, in Gorontalo Province, using the Holt-Winters exponential smoothing forecasting data model won't be over any time soon. Except

that, there is government intervention such as procuring vaccines to give up the spread of Covid-19. This can be seen in the visualization of Figure 4, where the model does not show a decreasing trend.

6. References

- [1] Zhang X, Ma R and Wang L 2020 Predicting turning point, duration, and attack rate of COVID-19 outbreaks in major Western countries *Chaos, Solitons & Fractals* 109829
- [2] Griffiths D F and Higham D J 2010 *Numerical methods for ordinary differential equations: initial value problems* (Springer Science & Business Media)
- [3] Gupta R and Pal S K 2020 Trend Analysis and Forecasting of COVID-19 outbreak in India *medRxiv*
- [4] Lalwani S, Sahni G, Mewara B and Kumar R 2020 Predicting optimal lockdown period with parametric approach using three-phase maturation SIRD model for COVID-19 pandemic *Chaos, Solitons & Fractals* 109939
- [5] Malavika B, Marimuthu S, Joy M, Nadaraj A, Asirvatham E S and Jeyaseelan L 2020 Forecasting COVID-19 epidemic in India and high incidence states using SIR and logistic growth models *Clinical Epidemiology and Global Health* 9 26-33
- [6] Kurita J, Sugawara T, & Ohkusa Y 2020 Forecast of the COVID-19 outbreak and effects of self-restraint in going out in Tokyo Japan *medRxiv*.
- [7] Kalekar P S 2004 Time series forecasting using holt-winters exponential smoothing *Kanwal Rekhi School of Information Technology* 4329008 1-13
- [8] Pongdatu G A N and Putra Y H 2018 Seasonal time series forecasting using SARIMA and Holt Winter's exponential smoothing *In IOP Conference Series: Materials Science and Engineering* 407 1 012153
- [9] Abd Jalil N A, Ahmad M H and Mohamed N 2013 Electricity load demand forecasting using exponential smoothing methods *World Applied Sciences Journal* 22 11 1540-1543
- [10] Djauhari M A, Asrah N M, Li L S and Djakaria I 2020 Forecasting model of electricity consumption in Malaysia: A geometric Brownian motion approach *Solid State Technology* 63 3 40-46
- [11] Panda M 2020 Application of ARIMA and Holt-Winters forecasting model to predict the spreading of COVID-19 for India and its states *medRxiv*
- [12] Bezerra A K L and Santos É M C 2020 Prediction the daily number of confirmed cases of COVID-19 in Sudan with ARIMA and Holt Winter exponential smoothing *International Journal of Development Research* 10 08 39408-39413
- [13] Makatjane K and Moroke N 2016 Comparative study of holt-winters triple exponential smoothing and seasonal Arima: forecasting short term seasonal car sales in South Africa *Risk governance & control: financial markets & institutions* 6 1 Winter 2016
- [14] Djakaria I 2019 Djalaluddin Gorontalo airport passenger data forecasting with Holt's-Winters' exponential smoothing multiplicative event-based method *IOP Conference Series: Journal of Physics: Conference Series* 1320 1 012051
- [15] Jere S and Siyanga M 2016 Forecasting inflation rate of Zambia using Holt's exponential smoothing *Open Journal of Statistics* 6 2 363-372
- [16] Sumitra I D and Basri I 2020 Forecasting the number of outpatient patient visits using the ARIMA, SES and Holt-Winters methods at XYZ Community Health Center *IOP Conference Series: Materials Science and Engineering* 879 1 012060
- [17] Makridakis S, Wheelwright S C, Hyndman R J and McGee V E 1999 *Forecasting Methods and Application* (Hoboken: John Wiley & Sons, Inc.)
- [18] Worldometer 2020 Coronavirus cases data <https://www.worldometers.info/coronavirus/#countries> accessed on September 19, 2020

Covid-19 forecast using Holt-Winters exponential smoothing

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	archive.conscientiabeam.com Internet Source	1 %
2	journal.universitasbumigora.ac.id Internet Source	1 %
3	journals.plos.org Internet Source	1 %
4	Wei Du, Xiaohua Zhao, Huiming Hou, Jinping Fu, Jianxin Su. "Analogy of scaling law for heterogeneous interfacial fracture to explain the Gutenberg–Richter formula for earthquakes", Journal of Statistical Mechanics: Theory and Experiment, 2022 Publication	1 %
5	pdfs.semanticscholar.org Internet Source	1 %
6	Submitted to Sabanci Universitesi Student Paper	1 %
7	Abdelhafid Zeroual, Fouzi Harrou, Abdelkader Dairi, Ying Sun. "Deep Learning Methods for Forecasting COVID-19 Time-Series Data: A	1 %

Comparative Study", Chaos, Solitons & Fractals, 2020

Publication

8	Submitted to CSU, San Francisco State University Student Paper	1 %
9	www.doria.fi Internet Source	1 %
10	id.123dok.com Internet Source	1 %
11	Submitted to Erasmus University of Rotterdam Student Paper	<1 %
12	e-journal.unair.ac.id Internet Source	<1 %
13	ojs3.unpatti.ac.id Internet Source	<1 %
14	"Study on Production Planning Demand Model Based on SPSS Statistical Analysis", Academic Journal of Computing & Information Science, 2023 Publication	<1 %
15	Huber Nieto-Chaupis. "Exploring Geographical Topologies and Diffusion of Monkeypox Infections at the Beginning Pandemic", 2022 IEEE 22nd International Conference on	<1 %

Bioinformatics and Bioengineering (BIBE), 2022

Publication

16

www.ijert.org

Internet Source

<1 %

17

"Thermal Analisis of the Increase in Ambient Temperature Due to Motor Vehicle Activities", International Journal of Engineering and Advanced Technology, 2019

Publication

<1 %

18

jglobal.jst.go.jp

Internet Source

<1 %

19

repository.untar.ac.id

Internet Source

<1 %

20

www.ihs.nl

Internet Source

<1 %

21

"Artificial Intelligence for COVID-19", Springer Science and Business Media LLC, 2021

Publication

<1 %

22

Fachrur Rozi, Agung Budi Santoso, I Gusti Ayu Putu Mahendri, Ronald Timbul Pardamean Hutapea et al. "Indonesian market demand patterns for food commodity sources of carbohydrates in facing the global food crisis", Heliyon, 2023

Publication

<1 %

23 Rediat Takele Figa. "Stochastic modelling for predicting COVID-19 prevalence in East Africa Countries", Infectious Disease Modelling, 2020
Publication <1 %

24 publications.afropolitanjournals.com
Internet Source <1 %

25 www.ripublication.com
Internet Source <1 %

26 www.scribd.com
Internet Source <1 %

27 www.tisc.rs
Internet Source <1 %

28 Payal Rana, Sudipa Chauhan, Anuj Mubayi. "Burden of cytokines storm on prognosis of SARS-CoV-2 infection through immune response: dynamic analysis and optimal control with immunomodulatory therapy", The European Physical Journal Special Topics, 2022
Publication <1 %

29 www.mdpi.com
Internet Source <1 %
