



Universitas Perikanan dan Ilmu Kelautan
Universitas Negeri Gorontalo
Indonesia

ISBN No : 978-602-72985-0-7




PROCEEDING

International

SEMINAR 2015

in accordance with Sail Tomini and
Festival of Boalemo 2015



Fishery and Marine Science

Supported By :



Proceeding

ISBN : 978-602-72985-0-7

International Seminar Fishery and Marine Sciences in Accordate with Sail Tomini and Festival of Boalemo. Fakultas Perikanan dan Ilmu Kelautan Universitas Negeri Gorontalo. 2015

Pelindung

Rektor Universitas Negeri Gorontalo
(Prof. Dr. H. Syamsu Qamar Badu, MP)

Penanggung Jawab

Dekan Fakultas Perikanan dan Ilmu Kelautan
(Dr. Abdul Hafidz Olii, S.Pi, M.Si)

Editors

Basri Amin, S.Sos, MA
Prof. Dr. Ir. Mahludin Baruwadi, MP
Dr. Ir. Syamsuddin, MP
Dr. Aziz Salam, ST, M.Agr
Dr. Rieny Sulistijowati, S.Pi, M.Si
Dr. Juliana, S.Pi, MP
Faizal Kasim, S.IK, M.Si

Redaktur Pelaksana

Mulis, S.Pi, M.Sc
ZC. Fachrusyiah, S.St.Pi, M.Si
Wawan K. Tolinggi, SP, M.Si
Ir. Rully Tuiyo, MP


Desain Grafis

ZC. Fachrusyiah, S.St.Pi, M.Si

Penerbit:

Fakultas Perikanan dan Ilmu Kelautan
Universitas Negeri Gorontalo
Jalan Jendral Sudirman No. 6 Kota Gorontalo
Gorontalo-Indonesia

Foto Cover : bpmptsp.gorontaloprov.go.id, ekokusnur.com
Copyright © September 2015



International Seminar Fishery and Marine Sciences in Accordance with Sail Tomini and Festival of Boalemo. Fakultas Perikanan dan Ilmu Kelautan Universitas Negeri Gorontalo

BN : 978-602-72985-0-7

Hak Cipta dilindungi Undang-undang

rights reserved

Editors : Basri Amin, Mahludin Baruwadi, Syamsuddin, Aziz Salam, Rienny Sulistijowati, Juliana, Faisal
sim

Terbitkan Oleh : Fakultas Perikanan dan Ilmu Kelautan Universitas Negeri Gorontalo

Dilarang memperbanyak sebagian dan atau seluruh isi buku ini tanpa seizin dari penyunting



Proceeding

ISBN : 978-602-72985-0-7

International Seminar Fishery and Marine Sciences in Accordance with Sail Tomini and Festival of Boalemo. Fakultas Perikanan dan Ilmu Kelautan Universitas Negeri Gorontalo. 2015

- Assessment Of Coral Reefs Condition In Molotabu Waters Bone Bolango District**
Femy M Sahami, Sri Nuryatin Hamzah, Sitti Nursinar..... 1-11
- Effectiveness And Productivity Hand Lines With Different Branch Line Around Portable Fish Aggregating Device**
Roza Yusfiandayani, Diniyah dan Wawan Dedi Ariawan..... 12-21
- The Impact Of Moratorium To The Catch Of Fishermen With Boat Size Under 30 Gt (Case Study: Bitung, Manado Cities And Minahasa Utara, Minahasa Tenggara Regencies)**
Samuel Hamel, Hetty M.P. Ondang, Daniel H. Ndahawali, Khairul Jamil, Lexy Hosang, Karyanto..... 21-29
- Prospective Capture Fisheries In Northern Gorontalo District**
Lis Melissa Yapanto, Syamsuddin 30-36
- Vegetation Structure Of Seagrass In Saronde Island, Ponelo Archipelago Subdistrict, North Gorontalo District, Gorontalo Province**
Siti Rahmi A.R. Nusi, Abdul Hafidz Oliy, Syamsuddin 37-48
- Chemical Characteristics Of Fishbones Gelatin That Was Extracted From Tuna (*Thunnus* Sp) By Using Of Palm Vinegar Acid**
Asri Silvana Naisu, Nikmawatusanti Yusuf, Mohammad Zulkifly..... 48-59
- The Seedling Of Flying Fish (*Hirundichthys Oxycephalus*) High Economic Value Of Aquatic Biota As An Effort Of Germplasm Preservation In Indonesian Eastern Water Areas**
Sugeng Raharjo, Dasep Hasbullah, Endah Soetanti, Sadat, Hamita Agusanty, Muhammad Amri Tiro..... 60-75
- Lawi-Lawi Cultivation (*Caulerpa*, Sp) Innovative Technology Of Fishery Agriculture Sector Environmental Friendly And High Profitability**
Dasep Hasbullah, Sugeng Raharjo, Endah Soetanti, Jumriadi, Imran Lapong, Hamita Agusanty Mike Rimmer..... 76-90
- Isolation And Identification Of Bioactive Compound Of Kopasanda (*Chromolaena Odorata*) Leaf To Fight *Vibrio Harveyi* On Post-Larval Tiger Prawn (*Penaeus Monodon* Fabricius)**
Harlina, Syahrul, Jayadi, Rustam Zaenal Abidin 91-101
- The Design And Specification Of Deep Sea Trawl (Midwater Trawl) Operated In Okhotsk Sea – Russia**
Aris Widagdo, Nyoman Suyasa, Danu Sudrajat, Eko Rahmanto, Rachmat Juanda..... 102-112
- Wangurer Fishermen's Socio-economic After Moratorium Enforcement. (Case Study : Under 30 GT Fishermen in Wangurer District, Bitung City)**
Novie Wijaya, Khairul Jamil 113-118
- Organoleptic Hedonic Quality And Chemical Analysis On Sticks Seaweed *Kappaphycus Alvarezii* With Fortification Flour Rebon Shrimp (*Mysis* Sp.)**



Rita Marsuci Harmain, Faiza Dali, Winawanti Amrulla.....	119-130
The Physics And Chemical Characteristics Of Sausage Catfish Subtitution By Algae (<i>Kappaphycus Alvarezii</i>) Rieny Sulistijowati, Rita Harmain, Nur Hidayat Rauf.....	131-140
Propeller Shaft Bearing of Fishing Vessel Kaminton Tambunan	141-148
Development of Grouper Aquaculture with Community-Based Partnership Fachrudin Zain Olilingo.....	149-153
Resistance Of The Three Species Of Seaweed (<i>Kappaphycus Alvarezii</i>, <i>Euचेuma Striatum</i>, And <i>Euचेuma Denticulatum</i>) To The Extreme Salinity In Waters Of Tomini Bay Petrus Rani Pong Masak, Wiwin Kusuma Perdana Sari, and Nova F. Simatupang.....	154-159
Diversity And Density Of Macroinvertebrates (Order: Diptera) In Umbulrejo River, District Dampit Malang Miftahul Khair Kadim.....	160-166
Demersal Fish Caught In Lembeh Strait (Case Study : Vertical Hand Line) Heru Santoso, Khairul Jamil, Yuli Purwanto.....	167-172
Valuasi Nilai Ekonomi Perikanan Danau Limboto Provinsi Gorontalo Hasim.....	168-178
Analisis Ramah Lingkungan (<i>Environment Friendly</i>)Perikanan Cakalang (<i>Katsuwonuspelamislinneus</i>) Berkelanjutan Di Perairan Teluk Tomini Propinsi Gorontalo Syamsuddin, Achmar Mallawa, Yuniarti Koniyo.....	179-194
Partisipasi Masyarakat Pesisir Dalam Pengelolaan Danau Limboto Provinsi Gorontalo Bebay S.D. Banteng	195-203
Adaptation and Mitigation against Rob Flood Based Socio-Economic Assessment in Coastal Communities Sriwulan Village, Sayung, Demak Nur Indah Septriani*, Suparjo Rasazli Carong, Falmuriati, Fitriawan Umar, Gigih Girriastowo.....	204-217
Studi Peran Perempuan Pesisir Dalam Menunjang Aktivitas Perikanan Di Desa Torosiaje Laut Kecamatan Popayato Kabupaten Pohuwato Sumrin, Abd. Hafidz Olli, Alfi Sahri Remi Baruadi,.....	218-229
The analysis of production factors of catching yellowfin tuna (<i>thunnus albacares</i>) the small-scale in seram sea waters Haruna Dewi Shinta Achmad.....	230-236
The Use Of Plant Tembelekan (<i>Lantana Camara</i> Linn) Extract To Prevent Ice-Ice Disease And Trigger Growth Rate Of Seaweed <i>Kappaphycus Alvarezii</i> Rahmat Sofyan Patadjai , Indriyani Nur and Syamsul Kamri.....	237-250



Onboard Observation Of Deep Sea Trawl (Midwater Trawl) Operated In Okhotsk Sea-Russia
Aris Widagdo, Nyoman Suyasa, Danu Sudrajat, Eko Rahmanto, Rachmat Juanda251-258

Rendement and chemical composition alteration During processing of dried abalone *haliotis asinina*
Maharani.....259-269

Analysis Of Suitability And Carrying Capacity Of Marine Tourism At Bandengan Waters In Jepara District
Citra Panigoro, Juliana.....270-276

Biodiversity of Coral Fish in Bontang Waters, East Kalimantan
Ristiana Eryati, Muchlis Efendi dan Irwan Ramadhan R278-287

Heavy Metals Concentrations In Fish Cultivated In Floating Cage In Kutai Kartanegara
Ghitarina, Sulistyawati, Henny Pagoray, Abdunnur, Ahmad Rafii288-294

The assessment of the sea coral reefs in the waters needs to be done in order to ensure the survival of a community. The aim of the research was to assess the current state of the coral reefs in Molokan waters, Molokan District, Gorontalo Province. The assessment was consisted of the coral reef's condition coverage percentage, life form categories, and site threat. Site location was divided 4 station. Data collection was conducted by using line transect method placed at depth of 3 meters and 10 meter respectively. The condition of the coral reefs was analyzed by referring to KGP-MENLH No. 42/191. Water quality was tested in which including salinity, temperature, pH, dissolved oxygen and water brightness. Results showed that the Coral Massive was dominant form at depth of 3 meters, whereas at 10 meters significant different between life form by it. The highest coverage shown by coral reefs at 10 meters depth of 41.1%. The percentage gives the figure about the bad condition of the coral reefs in Molokan waters.

Keywords: Assessment, Coral Reefs, Molokan Waters

1. INTRODUCTION

Coral reefs are one of the under marine ecosystems that support marine life's productivity and high economic value. Coral reefs function as a place to spawn, feed, marine life habitat, and as a source of protection as well as the headquarters of fishing communities through various activities as well as tourism activities. Furthermore coral reefs are also a source of bioactive substance raw materials which is used in pharmacy, and medicine, and another important function is as coastal protection from degradation and erosion, but currently coral reefs in Indonesia has undergone many damage degradation (CORREMA, 2019).

Coral reef ecosystem is one of the aquatic ecosystems with high level of productivity. The existence of coral ecosystems can be found in almost all Indonesian waters, including in the Gulf of Tomini which environment is rich (Ministry of Environment, 2009). Sustainability of the reef is largely determined by environmental conditions. Coral reefs can grow well when environmental conditions conducive to its growth. Molokan Waters (Molokan) is part of the Gulf of Tomini.

Sonanti & Hamzan (2022), the condition of coral reefs in the waters around Molokan Village waters in Molokan is "seriously damaged" with low coral coverage of 40%. The extent of damage will increase if no treatment or management is taken and this may affect the life of associated organisms in it and support the sustainability of the industry occurred in the future.



THE PHYSICS AND CHEMICAL CHARACTERISTICS OF SAUSAGE CATFISH SUBSTITUTION BY ALGAE (*Kappaphycus alvarezii*)

Rieny Sulistijowati, Rita Harmain, Nur Hidayat Rauf

Faculty of Marine and Fisheries Gorontalo State University

PO BOX 5, Zip Code 96126 Indonesia

rinyulistijowati@gmail.com

Abstract

This research was conducted to know the physics and chemical characteristics of sausage catfish substitution by algae (*Kappaphycus alvarezii*). This research was an experimentally research conducted in laboratory of Agency Analise Agricultural Product Bogor used completely randomized design with repeated two times. The range of concentration substitution *Kappaphycus alvarezii* used 0% (control), 11%, 15% and 18%. The physics parameters were springiness and cohesiveness, the chemical parameters were protein, fat, water, ash and fiber content. The results showed that treatment of concentration substitution 18 % of *Kappaphycus alvarezii* can improved physics characteristics sausage catfish; springiness 3.63 mm and cohesiveness 0.72. The treatment of concentration substitution 15 % of *Kappaphycus alvarezii* can improved chemical characteristics sausage catfish; protein content 13.59%, fat content 2.97%, water content 60.68%, ash content 0.2%, fiber content 7.83% and carbohydrate content 14.69%.

Keywords: Sausage catfish, substitution, *Kappaphycus alvarezii*

INTRODUCTION

Development of Gorontalo Province has potential as a commodity catfish freshwater. The production of catfish as a commodity cultivation in 2013 amounted to 41.95 tonnes (Dinas Perikanan dan Kelautan, 2013). Public interest to consume catfish are very low. This is due to the slippery catfish morphology and head shape so that the width of Gorontalo people assume morphology like a snake. In addition, the old catfish generally large so as not sold alive or fresh, but serve as a parent to death so as not consumed (Siregar et al, 2013).

According Nurimala et al (2009), catfish has the characteristic appearance of the meat is white with a soft fiber making it suitable as a raw material pulverized fish flesh or surimi (the basic material of fish gel based products one of which is a sausage catfish). In addition, the catfish has a savory taste and contains 17% protein and 0.95% low fat. *Kappaphycus alvarezii* algae is one type of seaweed as raw material for carrageenan and used in foodstuffs because it has the ability to alter the functional properties of the product desired gel properties (Afriwanty 2008). *K. alvarezii* contains some important chemical content of the dietary fiber and iodine are quite high. According Astawan et al (2004), the content of dietary fiber in seaweed *K. alvarezii* amounted to 78.94% and amounted to 282.93 mg iodine / g. According Dinas Perikanan dan Kelautan (2011) the production of algae *K. alvarezii* is 5,228 tonnes in 2011.

The object this research sausage catfish substitution by algae (*K.alvarezii*). The intent of this study was effect substitution algae (*K.alvarezii*) toward physics and chemical sausage catfish.



II. MATERIALS AND METHODS

2.1 Materials and equipment

The material for processing of sausages used catfish the size of 600-900 kg / head, bamboo as a sleeve (1 cm long and 1.5 cm in diameter), sugar, ice cubes, corn oil, tapioca starch, seaweed harvesting *K. alvarezii*, pepper, garlic, onion, salt (NaCl), red chillies. Curing material consists of sawdust, coconut shells, coconut husks. Materials chemical used distilled water, K₂SO₄, HgO, H₂SO₄, NaOH 30%, H₃BO₃, indicator methylred, 0.02 N HCl, chloroform, petroleum ether, hexane and ninhydrin.

Equipment used in this study were knives, cutting boards, containers, filter cloth, scales, grinder, processor, blender, stuffer, pan, thermometer, a cup of metal, cup porcelain, pestle and mortar, desiccator, analytical balance, an electric heater, Bunsen, steam bath, hot plate, furnace, distillation, burette, funnel, pipette, flask 50 mL, 125 mL erlenmeyer, Soxhlet flask, extractor soxhlet, texture analyzer.

The Processing Sausage Catfish substitution by algae (*K.alvarezii*)

The steps processing of sausage catfish substitution by algae (*K.alvarezii*) were the algae porridge. Digestion of catfish meat and ground spices. Mixing catfish, ground spices and concentration algae porridge 0, 11, 15 and 18% (L0, L1, L2, L3), charging sleeve, steaming, smoking and analyse data.

Data Analyzed

The physics parameters were springiness and cohesiveness, the chemical parameters were water, protein, fat, and fiber content. The measurement springiness and cohesiveness has been done according to method illustrated by Chen et al (2013). Water content used gravimetric method (ISO 01.2354.2-2006), ash content used gravimetric method (SNI 01-2354.1-2006), protein content used microkjeldahl (SNI 01-2354.4-2006), fat content used extraction method soxhlet, crude fiber by Sudarnadji (1997), and carbohydrate content difference (Winarno, 2002).

2.4. Research Design

The was carried out experimentally in the laboratory by using a completely randomized design non fact. The treatment were concentration substitution *K. alvarezii* 0, 11, 15 and 18% (control, L1, L2, L3). Each treatment was repeated 2 times, experiment total 8 units. Parameters were observed physical characteristic (springiness and cohesiveness) and chemical characteristics (water, ash, protein, fat fiber and carbohydrate content). The results were analyzed statistically by ANOVA analysis followed by Duncan's Multiple Range Test (DMRT) if significant different.

III. Result and Discussion

3.1 Physical Characteristics of Sausage Catfish

The physical properties of the sausage catfish c sausage texture will show the quality of the sausage because the sausage is a product -based gel (basic gel product). Sausage texture parameter measurement include springiness and cohesiveness. The results of the physical analysis of the texture of sausages catfish can be seen in Figure 1.



Phisics Characteristics Sausage Catfish




Figure 1. Phisics Characteristic of Sausage Catfish

Based on Figure 1, shows that the average value of springiness lowest for the treatment of L2 is 1.83 mm and the lowest value contained in the treatment cohesiveness L2 is 0.41. The highest springiness value and cohesiveness are on treatment L3 are 3.63 mm and 0.72. Results of analysis of variance that the substitution K. alvarezii significant effect ($P < 0.05$) against the springiness of sausage catfish. Duncan advanced test distance L3 that the springiness treatment (substitution K. alvarezii 18%) was significantly different to the treatment L1 and L2.

Results of analysis of variance that the substitution K. alvarezii significantly affect the value of the cohesiveness of the sausage catfish texture. Duncan analysis that K. alvarezii substitution of 18% was significantly differences to controls, L1 and L2, while among the controls, L1 and L2 undifference.

Differences in the springiness of the sausage catfish value influenced by the amount of starch and pulp K. alvarezii varied. The springiness and cohesiveness of the sausage on the treatment L3 higher than L1 and L2 sausage due to the amount of porridge K. alvarezii 18%.

The springiness and cohesiveness of the texture undifference in the control, L1 and L2 treatment. The use of starch alone produces less chewy texture objectively because the lack of ability of the starch to absorb water. Amylopectin starch contained in the group responsible for forming a chewy texture, but on L1 and L2, have been using a mixture of tapioca higher than K. alvarezii. Amount of K. alvarezii indirectly make carrageenan against water holding capacity reduced so that the gel formation mechanism sausage sausage reduced so that the springiness and cohesiveness of the texture considered undifference in value.

Carrageenan more Water Holding Capacity (WHC) is strong have a high degree of polarity that can form a strong matrix when binds with water, thus indirectly carrageenan will bind water will make the gel becomes stronger (Yulianti, 2003). Carrageenan types kappa contained in K. alvarezii a jelly that is rigid, hard and brittle compared to iota carrageenan is soft (Santoso and Nugraha, 2008). This research the substitution K. alvarezii high (18%) make the cohesiveness of sausage catfish increase.



3.2 Chemical Characteristic of Sausage Catfish

- Water Content

Knowledge of food product an water content can provide information about freshness , texture and shelf quality deterioration) a food products. Results of the analysis of water content catfish substituted sausage b *alvarezii* can be seen in Figure 2.




Figure 2. Water content of Sausage Catfish

The water content due to the hydrophilic properties between starch and pulp *K.alvarezii*. Tapioca contains starch granules which can bind with the aid of hot water and then it will turn into a gel. There are two in the starch fraction of the amylose and amylopectin. Based Figure 2 shows that the average water content ranges from 57.8 to 71.43% with the lowest water content (57.8%) contained in the control sausage and the high water content (71.43%) contained in the sausage L3 (substitution *K. alvarezii* 18%). The higher the substitution amount of tapioca and *K. alvarezii* against the increasing water content of sausages. Analysis of variance that the substitution *K. alvarezii* highly significant effect ($P < 0.01$) on increasing the water content of the sausage.

Duncan advanced test distance that the treatment of the water content of sausages treated L3 differ toward sausages control, L1 and L2 treatment. However, the treatment sausage on L1 and L2 and control undifferent. Increased improve the ability to bind more water from amylose.

K. alvarezii algae have properties that can bind water because it contains a functional group the karageenan that acts as hydrocolloid (Afriwanti, 2008). Carrageenan has sulfate groups that can bind water. content of sulfate groups that are on the negatively charged carrageenan along the polymer chain is hydrophilic and which can bind water or other hydroxyl group (Widodo, 2008).

Based on the ISO 3020-1995 quality sausages, sausages required moisture content is a maximum of 60%. treatment sausage L1, L2 and control, water content quality levels sausage. But sausage treatment L3 not eligible because the water content based SNI has exceeded the maximum limit 71.43%.

- Ash Content

The data of ash content sausage catfish substitution by algae (*K.alvarezii*) can be seen in Figure 3.






Figure 3. Ash content of Sausage Catfish

Figure 3 shows the average value of ash content ranged sausage between 0.18 to 0.25%. Low ash content is on the sausage control (without *K. alvarezii*) ie 0.18% and ash content of the highest catfish sausage treatment (substitution *K. alvarezii* 18%) which amounted to 0.25%. However, the findings of the analysis of variance that the substitution *K. alvarezii* insignificantly affect ash content of sausages catfish.

The describe of ash content depicting gross inorganic substances in the hearts of food products. Posted dried food organic material is 96% organic and water, the rest consisting of the mineral elements. Mineral elements inorganic substances known also inorganic or ash content from combustion process, materials organic materials will burned out, while the Inorganic Materials NO, which is called with ash (Winarno, 2002).

All treatments ash content meets the requirements set out ash that posted ISO 3020-1995 maximum of 3%. Levels of ash caused sausage *K. alvarezii* used laundering has experienced many times with fresh water so the ash content reduced. According Vishnu and Rachmawaty (2010), *K. alvarezii* washing by fresh water reduces ash content lower than the 15.13 washing by salty air that produces ash 35% higher.

Content

Fats in food have diverse functions as a source of energy, contributing to the formation of texture, heat transfer medium and as a solvent in vitamins A, D, E and K. The results of the analysis of fat sausage catfish substitution *K. alvarezii* can be seen in Figure 3.






Figure 4. Fat content of Sausage Catfish

Figure 4 shows a decrease in the levels of fat sausage along with increasing concentrations of *K. alvarezii* substituted on the sausage catfish. The highest fat content found in the control sausage (without *K. alvarezii*) which is 6.68% and the lowest fat content found in sausages treatment L3 (substitution *K. alvarezii* 18%) was 1.24%.

Analysis of variance that there is a highly significant effect ($P < 0.01$) the substitution *K. alvarezii* towards decreased levels of fat sausage. Duncan advanced test distance that the fat content of sausage control and L1 (substitution *K. alvarezii* 11%) highly significant different with sausage fat content L2 and L3 (substitution *K. alvarezii* 15% and 18%). *K. alvarezii* substitution of 15% or 18% has been able to lower the amount of fat in sausages significantly. This is because *K. alvarezii* has the ability to absorb some organic elements like grease. That is because the ability of *K. alvarezii* fiber. According Marsono (2004), that the fiber-containing foods such as *K. alvarezii* have organic absorption capability is the ability to absorb some organic matter such as fat and cholesterol. Sausage fat content catfish quality fat levels set by ISO 3020-1995 which is a maximum of 25%. Fat content in sausages were lower affected by the constituent materials that contain fats such as meat catfish and vegetable oils. Meat catfish fat very low at 0.29% (Nurilmala et al 2008).

-Protein Content

Protein is one of the main important nutrients because the human body needs for growth and for other metabolic functions. Results of the analysis protein content of sausages catfish substitution *K. alvarezii* can be seen in Figure 5.






Figure 5. Protein content of Sausage Catfish

Based on Figure 5 shows that the protein value catfish sausages increased with increasing concentrations of *K. alvarezii* substitution. The lowest protein content found in the control treatment (without *K. alvarezii*) 13.28% and the highest protein content found in sausages treatment L3 (substitution *K. alvarezii*) 13.75%. Results of analysis of variance that the substitution *K. alvarezii* significant effect ($P < 0.05$) increased levels of the protein. Results within Duncan test that the sausage L3 significantly differences treatment outcomes to the control sausage (without the addition of seaweed) and sausage L1 treatment (substitution *K. alvarezii* 11%). This is because *K. alvarezii* also contains proteins which amounted to 5.9% (Astawan et al 2004). Any increase in the concentration of *K. alvarezii* then an increase in the value of protein statistically sausage catfish.

Research results Widodo (2008) reported that the addition of the sausage karageenan of 0.5% would increase the amount of protein in fish sausage kurisi (*Nemipterus nematophorus*). The protein content of catfish throughout treatment sausage eligible protein based ISO 3020-1995 that at least 13%. The protein content affected by the component catfish meat, eggs and other protein-containing components. During preparation of the dough sausage or emulsions, meat protein has two functions: 1) emulsifying fat, and 2) a water binder (Kusnandar, 2011).

Fiber Content

Dietary fiber is a carbohydrate group nonpati, which is prepared on the class of polysaccharides such as pectin, lignin and gums. Foods that have high fiber value is one of the criteria for healthy foods because fiber has many benefits, especially in the human digestive health. Results of the analysis of the fiber in sausages treatment results can be seen in Figure 6.

Figure 6 shows an increase in the fiber content of sausages catfish line with increasing concentrations of *K. alvarezii* substitution. Sausage control (without *K. alvarezii*) contain low fiber content which is 1.29% while the sausage result L3 treatment (substitution *K. alvarezii* 18%) containing the highest fiber content which 8.83%. However, analysis of variance that insignificantly effect substitution *K. alvarezii* to increase fiber content.

The formula sausage catfish used tapioca control 20%, L1 30%, L2 25% and L3 20%. There all any fiber content. So fiber in control treatment from tapioca.






Figure 6. Fiber content of Sausage Catfish

Astawan et al (2004) stated that *K. alvarezii* very high in fiber which is 78.94%. The recommended fiber intake is 30 g / person / day. Based on the recommended amount of fiber is the amount that can be consumed L3 sausage is a sausage of 400 g / day for the fulfillment of the fiber content. Marsono (2004) describes the functions of the human digestive fibers that are not water-soluble dietary fiber (insoluble fiber) have a nature easily withstand water, causing the stool bulky / expands (bulky) and easily removed. The nature of hail also caused by the increasing age of bacteria in the feces that is rich in fiber, because fiber food is an excellent substrate for the growth of the microflora in the colon.

-Carbohydrate Content

Carbohydrates are one of the macromolecule that is used as a source of energy for the human body. Carbohydrate analysis results can be seen in Figure 7. Figure 7 shows the average value of carbohydrate decreased in line with increasing concentrations of *K. alvarezii* substitution. The highest carbohydrate contained in the control is 20.78% and the lowest for the treatment of carbohydrates L3 (substitution *K. alvarezii*) is 4.58%.

Results of analysis of variance that the substitution *K. alvarezii* insignificantly affect the amount of carbohydrates. Because all treatment used tapioca and *K. alvarezii* for springiness and cohesiveness of sausage. Carbohydrate contributed from both component.






Figure 7. Fiber content of Sausage Catfish

Carbohydrates calculated from the amount of 100 percent of the material components are reduced by the accumulation of other nutritional components, namely water, ash, protein, fat and fiber (Winarno, 2002). Thus that, the value of carbohydrates is influenced by the levels of other nutrients, carbohydrates value decreases due to nutrients such as water, protein, ash and fiber increases. Besides other cause, tapioca as a constituent of sausage is one source of carbohydrate in number decreases with increasing concentrations of *K. alvarezii* as treatment. Carbohydrate content catfish exception sausage sausage L3 (substitution *K. alvarezii* 18%) did not meet the carbohydrate levels set by ISO 3020-1995 is a maximum of 8%.

CONCLUSIONS

Based on the results and discussion of research results that have been obtained, it can be concluded:

1. Substitution of *K. alvarezii* by 18 % (L3) become springiness and cohesiveness better of the sausage catfish.
2. Substitution *K. alvarezii* results ash, fiber and carbohydrate content insignificant differences. Substitution *K. alvarezii* 15 % (L2) produces good water, fat and protein content of the sausage catfish.

REFERENCES

- Astawan, M., Koswara, S., dan Herdiani, F. 2004. Pemanfaatan Rumput Laut (*Eucheuma cottonii*) Untuk Meningkatkan Kadar Iodium dan Serat Pangan pada Selai Dan Dodol. *J Tekn. Dan Industri Pangan XV*(2): 61-69.
- Afriwenty, M.D. 2008. Mempelajari Pengaruh Penambahan Tepung Rumput Laut (*Kappaphycus alvarezii*) Terhadap Karakteristik Fisik Surimi Ikan Nila (*Oreochromis sp.*). [Skripsi]. Program Studi Teknologi Hasil Perikanan. Fakultas Perikanan dan Ilmu Kelautan. Institut Pertanian Bogor. Bogor.
- Atmadja, W, Kadi, Sulistijo, Rachmaniar. 1996. Pengenalan Jenis-jenis Rumput Laut Indonesia. Jakarta: Puslitbang Oseanologi-LIPI.



[BSN]. Badan Standarisasi Nasional. 1995. Standar mutu sosis. 3020-1995. Standar Nasional Indonesia. Jakarta: BSN.

_____. 2006. SNI 01-2346-2006-Petunjuk Pengujian Organoleptik dan atau Sensori. Standar Nasional Indonesia. Jakarta: BSN.

_____. 2006. Penentuan kadar air pada produk perikanan. SNI No. 01-2354.2-2006. Jakarta

_____. 2006. Penentuan kadar lemak pada produk perikanan. SNI No. 01-2354.3-2006. Jakarta.

_____. 2006. Penentuan kadar protein dengan metode total Nitrogen pada produk perikanan. SNI No. 01-2354.4-2006. Jakarta

_____. 2010. Penentuan kadar abu dan abu tak larut asam pada produk perikanan. SNI No. 2354.1-2010. Jakarta

Chen L, Opara UL. 2013. Texture measuerement approaches in fresh and processed food. J food research international 51 : 823 - 835

Marsono, Y. 2004. Serat Pangan Dalam Prospektif Ilmu Gizi. Pidato Pengukuhan Guru Besar di Fakultas Teknologi Pertanian. Universitas Gadjah Mada. Yogyakarta.

Nurilmala, M., Nurjanah, dan Utama. 2009. Kemunduran Mutu Lele Dumbo Pada Penyimpanan Suhu Chilling Dengan Perlakuan Cara Mati. *Jurnal Pengolahan Hasil Perikanan Indonesia*: 1-16.

Santoso dan Nugraha. 2008. Pengendalian Penyakit Ice-ice Untuk meningkatkan produksi Rumput Laut Indonesia. 37-42. *Jurnal Saintek Perikanan 3 (II)*: 37-42.

Williams PA, Phillips GO, Stephen Am, Churms SC. 2006. Gums and Mucilages. Di Dalam: Stephen AM, Phillips GO, Williams PA, editor. *Food Polysaccharides and their applications*. Ed ke-2. Boca Raton: Taylor & Francis. FL, pp 455-495.

Widodo, S. A. 2008. Karakteristik Sosis Ikan Kurisi Dengan Penambahan Isolat Protein Kedelai dan Karagenan Pada Penyimpanan Suhu Chilling dan Freezing. [Skripsi]. Bogor: Program Studi Teknologi Hasil Perikanan. Institut Pertanian Boogor.

Winamo, F. (2002). *Kimia Pangan dan Gizi*. Jakarta: Gramedia Pustaka.

Yulianti T. 2003. Mempelajari pengaruh karakteristik isolat protein kedelai terhadap mutu sosis [Skripsi]. Fakultas Teknologi Pertanian. Institut Pertanian Bogor. Bogor.



