



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Weny J.a Musa, Boima Situmeang, ...
Assignment title: For writers
Submission title: Anti-cholesterol triterpenoid acids fr...
File name: LJFP_A_1650762.pdf
File size: 1.22M
Page count: 6
Word count: 3,287
Character count: 16,254
Submission date: 20-Aug-2019 09:03PM (UTC-0700)
Submission ID: 1161920574

INTERNATIONAL JOURNAL OF FOOD PROPERTIES
https://doi.org/10.1080/10942312.2019.1650762

Taylor & Francis
Taylor & Francis Group

OPEN ACCESS Check for updates

Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae)

Weny J.A. Musa, Boima Situmeang, and Julinton Sianturi

^aDepartment of Chemistry, Faculty of Mathematic and Natural Sains, Gorontalo State University, Gorontalo, Indonesia; ^bDepartment of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten, Indonesia; ^cDepartment of Chemistry, Universitas Padjadjaran, Bandung, Indonesia

ABSTRACT
Two triterpenoid compounds, 3 β -hydroxy-Olean-12-en-28-oic acid (1) and 3,19-Dihydroxys-12-en-28-oic acid (2) were isolated from the leaves of *Saurauia vulcani* Korth. (Actinidiaceae). The chemical structures of compounds 1–2 were elucidated with spectroscopic data (UV, IR, ¹H, ¹³C, DEPT 135 $^{\circ}$, HMQC, HMBC, ¹H-¹H COSY NMR) and MS as well as compared with previously reported spectra data. All the compounds were evaluated for their anti-cholesterol activity by the Liebermann-Burchard (LB) colorimetric assay. Compound 1–2 showed a remarkable anti-cholesterol activity. Most importantly, the raised concentration of 1–2 exhibited a dose-dependent manner. The discoveries of anti-cholesterol compounds 1–2 were first reported.

ARTICLE HISTORY
Received 1 April 2019
Revised 22 July 2019
Accepted 26 July 2019

KEYWORDS
Anti-cholesterol;
Triterpenoid acids; Pirodot;
Saurauia vulcani Korth;
Actinidiaceae

Introduction

The genus *Saurauia vulcani* Korth. (Actinidiaceae) is known as pirodot plant and mainly distributed in tropical climate.^[1] This plant has not been researched well and less information about its phytochemical studies; however, its biological screening exhibited interesting pharmacological properties, including antihyperglycemic, antihyperlipidemic,^[1] anti-diabetes,^[2,3] anti-cancer,^[4] antihypoglycemic,^[5] and antioxidant.^[6]

Recently, our group screened the presence of triterpenoid compound from the leaves of *Saurauia vulcani* Korth.^[7] As a part of our continuing search for active compounds from this plant, we isolated and described the triterpenoid compounds, 3 β -hydroxy-Olean-12-en-28-oic acid (1) and 3,19-Dihydroxys-12-en-28-oic acid (2). These compounds were first reported from this plant. Compound 1–2 were evaluated for their anti-cholesterol activity by the Liebermann-Burchard (LB) colorimetric assay. As a result, the isolated compounds 1–2 showed a remarkable anti-cholesterol activity.

Materials and methods

UV spectra were measured by using Shimadzu UV-8452A in methanol. The IR spectra were recorded on a Shimadzu FT-IR Spectrometer in KBr. Mass spectra were recorded with Agilent 5977B single quadrupole GC/MS. NMR spectra were obtained with a JEOL JNM A-500 (500 MHz). Chemical shift is denoted in δ (ppm) relative to residual solvent peak as internal standard (CDCl₃: ¹H δ 7.26, ¹³C δ 77.0). Coupling constant (*J*) are reported in Hertz (Hz). Multiplicity abbreviation: s = singlet, d = doublet, dd = double doublet, t = triplet, m = multiplet. Chromatography separations were carried out on silica gel 60 (70–230 mesh, Merck), octa desyl silane (200–400 mesh, Fuji Silysia). TLC plates were pre-coated with silica gel GF254 (Merck, 0.25 mm). The target compounds were detected

CONTACT Boima Situmeang boima@stak.ac.id Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten, Indonesia

^aPresent address: Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, 14424 Potsdam, Germany.
© 2019 Weny J.A. Musa, Boima Situmeang and Julinton Sianturi. Published with license by Taylor & Francis Group, LLC.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae)

by Weny J.a Musa, Boima Situmeang, Julinton Sianturi

Submission date: 20-Aug-2019 09:03PM (UTC-0700)

Submission ID: 1161920574

File name: LJFP_A_1650762.pdf (1.22M)

Word count: 3287

Character count: 16254

Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae)

Weny J.A Musa, Boima Situmeang, and Julinton Sianturi

^aDepartment of Chemistry, ^bFaculty of Mathematic and Natural Sains, Gorontalo State University, Gorontalo, Indonesia; ^cDepartment of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten, Indonesia; ^dDepartment of Chemistry, Universitas Padjadjaran, Bandung, Indonesia

ABSTRACT

Two triterpenoid compounds, 3 β -hydroxy-Olean-12-en-28-oic acid (**1**) and 3,19-Dihydroxyurs-12-en-28-oic acid (**2**) were isolated from the leaves of *Saurauia vulcani* Korth. (Actinidiaceae). The chemical structures of compounds **1–2** were elucidated with spectroscopic data (UV, IR, ¹H, ¹³C, DEPT 135°, HMQC, HMBC, ¹H-¹H COSY NMR) and MS as well as compared with previously reported spectra data. All the compounds were evaluated for their anti-cholesterol activity by the Liebermann-Burchard (LB) colorimetric assay. Compound **1–2** showed a remarkable anti-cholesterol activity. Most importantly, the raised concentration of **1–2** exhibited a dose-dependent manner. The discoveries of anti-cholesterol compounds **1–2** were first reported.

ARTICLE HISTORY

Received 1 April 2019
Revised 22 July 2019
Accepted 26 July 2019

KEYWORDS

Anti-cholesterol;
Triterpenoid acids; Pirdot;
Saurauia vulcani Korth;
Actinidiaceae



Introduction

The genus *Saurauia vulcani* Korth. (Actinidiaceae) is known as pirdot plant and mainly distributed in tropical climate.^[1] This plant has not been researched well and less information about its phytochemical studies; however, its biological screening exhibited interesting pharmacological properties, including antihyperglycemic, antihyperlipidemic,^[1] anti-diabetes,^[2,3] anti-cancer,^[4] antihypoglycemic,^[5] and antioxidant.^[6]

Recently, our group^[1] screened the presence of triterpenoid compound from the leaves of *Saurauia vulcani* Korth.^[7] As a part of our continuing search for active compounds from this plant, we isolated and described the triterpenoid compounds, 3 β -hydroxy-Olean-12-en-28-oic acid (**1**) and 3,19-Dihydroxyurs-12-en-28-oic acid (**2**). These compounds were first reported from this plant. Compound **1–2** were evaluated for their anti-cholesterol activity by the Liebermann-Burchard (LB) colorimetric assay. As a result, the isolated compounds **1–2** showed a remarkable anti-cholesterol activity.

Materials and methods

UV spectra were measured by using Shimadzu UV-8452A in methanol. The IR spectra were recorded on a Shimadzu FT-IR Spectrometer in KBr. Mass spectra were recorded with Agilent 5977B single quadrupole GC-MS. NMR spectra were obtained with a JOEL JNM A-500 (500 MHz). Chemical shift is denoted in δ (ppm) relative to residual solvent peak as internal standard (CDCl₃: ¹H δ 7.26, ¹³C δ 77.0). Coupling constant (*J*) are reported in Hertz (Hz). Multiplicity abbreviations: s = singlet, d = doublet, dd = double doublet, t = triplet, m = multiplet. Chromatography separations were carried out on silica gel 60 (70–230 mesh, Merck), octa desyl silane (200–400 mesh, Fuji Silysia). TLC plates were precoated with silica gel GF254 (Merck, 0.25 mm). The target compounds were detected

CONTACT Boima Situmeang  boima@stak.ac.id  Department of Chemistry, Sekolah Tinggi Analis Kimia Cilegon, Banten, Indonesia

*Present address: Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, 14424 Potsdam, Germany.

© 2019 Weny J.A Musa, Boima Situmeang and Julinton Sianturi. Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

12 by spraying with 10% H₂SO₄ in ethanol, followed by heating. All other commercially available reagents and solvents were used as purchased.

Plant material

The leaves of *Saurauia vulcani* Korth. (Actinidiaceae) were collected in Silangkitang Village, North Tapanuli, North Sumatera Province, Indonesia in July 2017. The plant was identified by the staff of the LIPI, Serpong, Indonesia and a voucher specimen (No. FR-SP 0101) was deposited at the herbarium.

Extraction and isolation

Dried ground leaves (1 kg) of *Saurauia vulcani* Korth. (Actinidiaceae) were extracted exhaustively with 9 hexane, ethyl acetate (EtOAc), and methanol (MeOH) at room temperature and then filtered. The evaporation resulted in the crude extracts of *n*-hexane (15.5 g), EtOAc (12.1 g) and MeOH (10.9 g), respectively. The EtOAc extract (12.1 g) 11 subjected to column chromatography over silica gel (70–320 mesh) using a gradient elution of mixture of 1 hexane/EtOAc (9/1 to 1/9) as eluting solvents yielded 11 fractions (E01-E11). Fraction E07 (192 mg) was subjected to reverse-phase column chromatography over ODS RP-18 using a gradient elution of mixture of MeOH/H₂O (7/3 to 10/0) as eluting solvents yielded 24 fractions (E07-1- E07-24). Further purification of fractions E07-11- E07-15 (23 mg) and E07-7- E07-10 (28 mg) by reverse-phase column chromatography using an isocratic elution of mixture of MeOH/H₂O (3/1) gave compounds 1 (13.7 mg) and 2 (11.3 mg), respectively.

Cholesterol determination

Liebermann–Burchard test was applied to measure cholesterol, which give 17 deep green color.^[8] Reductions of cholesterol levels of isolated compounds 1–2 were determined by comparison with the blank absorbance (cholesterol solution). The stock cholesterol solution was made by mixing cholesterol salt (100 mg) in 100 mL ethanol (1000 ppm) at 45°C in water bath. To 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of cholesterol solution in a cuvette (BRAN® UV cuvette) were added acetic anhydride (2 mL) and H₂SO₄ (0.1 mL) and diluted with ethanol (EtOH) to a final volume of 5 mL, respectively (blank solutions). To 0.5 mL of cholesterol solutions were added 0.025, 0.05, 0.1, 0.2, 0.3, and 0.4 mL of solution 1 and 2 (1000 ppm), acetic anhydride (2 mL) and H₂SO₄ (0.1 mL) and diluted with EtOH to a final volume of 5 mL, respectively. After being incubated in the dark for 15 min at room temperature, the absorbance of the solutions was measured at 423 nm.

Statistical analysis

Experiments were carried out independently three times and averages are presented. Statistical analysis was performed by unpaired two-tailed t-test (Excel). Differences were considered significant at **p* < .05, ***p* < .01 and ****p* < .001.

Result and discussion

The leaves of *Saurauia vulcani* Korth. (Actinidiaceae) (1 kg) 1 were grounded and successfully extracted with *n*-hexane, EtOAc and MeOH 11 room temperature. The EtOAc extract was chromatographed over column chromatography packed with silica gel (70–320 mesh) by gradient elution. The fractions were repeatedly subjected to reverse-phase column chromatography to give two anti-cholesterol compounds 1–2 (Figure 1).

Compound 1 was obtained as a white amorphous powder. The molecular formula of 1 was established to be C₃₀H₄₈O₃ from GC-MS Agilent MS which showed a [M+ Na]⁺ pseudo-molecular peak *m/z* 479.3496, found 479.3466, together with nuclear magnetic resonance (NMR) data, thus requiring seven




Figure 1. Chemical structures of compound 1–2.

degrees of unsaturation. The ultraviolet (UV) spectrum exhibited absorption peak at λ_{\max} nm (log ϵ): 202 nm explain no conjugated bond in compound. The Infrared (IR) spectrum showed absorption of hydroxyl group (3400 cm^{-1}), C-H aliphatic (2987 cm^{-1}), C-H bending (1456 cm^{-1}), gem dimethyl stretch (1374 cm^{-1}), a stretching of the C-OH group (1040 cm^{-1}), carbonyl group (1688 cm^{-1}), olefinic group C = C (656 cm^{-1}). The $^1\text{H-NMR}$ of **1** showed the presence of a triplet signal of olefinic proton resonating at δ_{H} 5.21 ppm ($J = 3.9\text{ Hz}$) as a position, an oxygenated methine resonating at δ_{H} 3.17 ppm (1H; dt , $J = 11.0, 3.9\text{ Hz}$) indicates as an axial position, double doublet signal of methine resonating at δ_{H} 2.90 ppm (1H; dd , $J = 9.7, 3.9\text{ Hz}$) indicated as α orientation, seven overlap signals of methyl groups, resonating at δ_{H} 1.17–0.77 ppm. The resonances suggesting the presence of a triterpene with olefin and alcohol functionalities of **1**. A total 30 carbon resonances were observed in the $^{13}\text{C NMR}$. The distortionless enhancement by polarization transfer (DEPT 135°) and Heteronuclear Multiple-Quantum Correlation (HMQC) experiments showed the peaks corresponding to two olefinic carbons at δ_{C} 122.2 and 144.2, one carbonyl carbon at δ_{C} 178.2, one sp^3 oxygenated methine, ten methylenes, three methines sp^3 , seven methyl, and six quaternary carbons. The Heteronuclear Multiple Bond Correlation (HMBC) spectrum confirmed the position of the C3 (δ_{C} 77.7 ppm) oxygenated methine carbon by showing correlations of H₃-23 and H₃-24. In addition, the signal of gem dimethyl showed mutual correlation between H-23 with C-24, H-24 with C-23, H-29 with C-30 and H-30 with C-29. The HMBC spectrum exhibited the correlation of H-11 with C-9, C-12 and C-13 and H-12 with C-9, C-11 C-13 and C-18 thus confirming the position of the double bond at C12 and C13, respectively (Figure 2). Furthermore, the location of carboxylic acid at C28 was established by the HBC correlation of H22 and H16 with the C28 carbonyl. The Correlation Spectroscopy ($^1\text{H-}^1\text{H COSY}$) experiment showed a correlation between H-11




Figure 2. Selected HMBC correlations of 1–2.

with H-12, H-3 with H-5 and H-18 with H-19. A comparison of the NMR data of **1** with 3 β -hydroxy-12(13)-en, 28-oic acid revealed that the structures of two compounds are very similar.^[9,10] Therefore, compound **1** was identified as a 3 β -hydroxy-Olean-12-en-28-oic acid.

Compound **2** was obtained as a white amorphous powder. The molecular formula of **2** was established to be C₃₀H₄₈O₄ from GC-MS Agilent MS which showed a [M+ Na]⁺ pseudo-molecular peak *m/z* 495.3445, found 495.3447, together with NMR data (Table 1), thus requiring seven degrees of unsaturation. The UV spectrum exhibited absorption peak at λ_{\max} nm (log ϵ): 204 nm explain no conjugated bond in compound. The IR spectrum showed absorption of hydroxyl group, C-H aliphatic, carbonyl group, C-H stretching, gem dimethyl stretch, and carbon *sp*² at ν_{\max} 3437, 2981, 1700, 1460, 1385, 727 cm⁻¹, respectively. The ¹H and ¹³C NMR spectra of **2** were resembled with compound **1** except for the signal of gem dimethyl at C20. The methine carbon C-19 was resonated downfield as quaternary carbon at δ 73.2 ppm since it is attached to a hydroxyl group. Furthermore, the location of hydroxyl at C-19 was also established by the HMBC correlation of H₃-30 and H₃-29 with C-19 and C20, respectively. A comparison of the NMR data of **2** with 3,19-Dihydroxy-12-ursen-28-oic acid revealed that the structures of two compounds are very similar.^[11] Therefore, compound **2** was identified as a 3,19-Dihydroxyurs-12-en-28-oic acid (**2**). The isolated compounds **1-2** were first reported from this plant.

3 β -hydroxy-Olean-12-en-28-oic acid (**1**)

A white amorphous powder; λ_{\max} MeOH nm (log ϵ): 202 (5.25); IR (KBr) ν_{\max} cm⁻¹: 3400, 2987, 1456, 1374, 1040, 1688, 656; ¹H NMR (CDCl₃, 500 MHz) see Table 1; ¹³C NMR (CDCl₃ 125 MHz), see Table 1; GC-MS Agilent MS: *m/z* 479.3466 [M+ Na]⁺ calculated 479.3496.

18

Table 1. NMR data (500 MHz for ¹H and 125 MHz for ¹³C in CDCl₃) for **1** and **2**.

Position	1		2	
	¹³ C NMR δ_C (mult.)	¹ H NMR δ_H (integral, mult., J Hz)	¹³ C NMR δ_C (mult.)	¹ H NMR δ_H (integral, mult., J Hz)
1	38.4	0.97 (1H; m); 1.55 (1H; m)	39.5	1.01 (2H; m)
2	27.2	1.52 (2H; m)	27.1	1.04 (2H; m)
3	77.7	3.17 (1H; dt, 11.0; 3.9)	78.6	3.16 (1H; dd, 11.1; 4.5)
4	39.5	-	39.5	-
12	55.4	0.83 (1H; m)	56.3	0.83 (1H; m)
6	18.3	1.51 (1H; m); 1.42 (1H; m)	15.2	1.51 (2H; m)
7	33.1	1.33 (2H; t, 3.6)	33.9	1.33 (2H; t, 4.2)
8	39.2	2	40.7	-
9	47.7	1.53 (1H; m)	48.2	1.71 (1H; m)
10	36.9	-	37.9	2
11	23.3	1.13 (2H; m)	24.4	1.35 (2H; m)
12	122.2	5.21 (1H; t, 3.9)	128.9	5.27 (1H; t, 3.9)
13	144.2	-	139.7	-
14	41.6	-	42.3	2
15	27.6	1.50 (2H; m)	28.1	1.13 (2H; m)
16	23.2	1.53 (1H; m); 1.92 (1H; m)	26.5	1.53 (1H; m); 1.92 (1H; m)
17	42.0	-	48.0	-
18	41.4	2.90 (1dd; 9.7; 3.9)	54.5	2.64 (1H; td, 8.5, 4.5)
19	46.0	2.07 (2H; m)	73.2	-
20	30.4	-	42.3	-
21	33.6	1.60 (2H; m)	32.7	1.30 (2H; m)
22	33.1	1.30 (17 m)	38.5	1.64 (1H; m); 1.73 (1H; m)
23	27.9	0.9 (3H; s)	29.4	0.98 (2; s)
24	15.5	0.79 (3H; s)	15.9	0.79 (3H; s)
25	15.5	0.77 (3H; s)	16.4	0.94 (3H; s)
26	16.8	0.84 (3H; s)	17.4	0.84 (3H; s)
27	25.4	1.17 (3H; s)	24.7	1.68 (3H; s)
28	178.2	-	179.2	-
29	28.6	0.98 (3H; s)	27.3	1.13 (3H; s)
30	23.1	0.79 (3H; s)	16.7	0.95 (3H; d, 4.6)

3,19-Dihydroxyurs-12-en-28-oic acid (2)

A white amorphous powder; UV MeOH λ_{\max} nm (log ϵ); 204 (5.23); IR (KBr) ν_{\max} cm^{-1} : 3437, 2981, 1700, 1460, 1385, 727; ^1H NMR (CDCl_3 , 500 MHz) see Table 1; ^{13}C NMR (CDCl_3 , 125 MHz), see Table 1; GC-MS Agilent MS: m/z 495.3447 [$\text{M} + \text{Na}$] $^+$ calculated 495.3445.

The anti-cholesterol effects of the two isolated compounds were carried out according to the described methods in the previous paper.^[12-14] With the isolated pure compound 1-2 in hand, we investigated an anti-cholesterol activity by them. Cholesterol level was determined by the LB colorimetric assay using UV Vis spectrophotometer.^[15,16] The cholesterol rate decreased when the isolated compounds 1-2 were treated, indicating that the anti-cholesterol induced by them. At a diluted concentration (5 ppm), the anti-cholesterol activity of 1-2 was limited (86% for 1 and 94% for 2). However, dose dependence of anti-cholesterol activity of 1-2 concentration was observed. The cholesterol level was dramatically reduced over a range of 10-80 ppm. The cholesterol levels were approximately 67%, 56%, 40%, 27% and 24% for 1 and 73%, 65%, 61%, 52% and 37% for 2 at the concentration of 10, 20, 40, 60 and 80 ppm, respectively (Figure 3). This result clearly showed the anti-cholesterol effect of isolated compound 1-2. It is also worth noting that an anti-cholesterol of compounds 1-2 were influenced by the presence of hydroxyl, double bond, and carboxylic acid groups. Orientation of the hydroxyl group of 2 showed less effective at reducing the cholesterol level. These results suggested that the additional of hydroxyl group into structure 2 may be a critical pharmacophore for its anti-cholesterol activity. The investigation of the *in vivo* using isolated compounds 1-2 is ongoing.




Figure 3. The anti-cholesterol effects of the isolated compounds 1-2. The level of cholesterol was reduced in a dose-dependent manner. Cholesterol level of the untreated compound or negative control was calculated as 100%. All experiments were conducted in triplicate ($n = 3$) and compared with negative control. Standard deviation (SD) is indicated by error bars. ns: not significant, * $p < .05$, ** $p < .01$, *** $p < .001$.

Conclusion

Two triterpenoids, 3 β -hydroxy-Olean-12-en-28-oic acid (**1**) and 3,19-Dihydroxyurs-12-en-28-oic acid (**2**) were isolated from the leaves of *Saurauia Vulcani* Korth (Actinidiaceae). The isolated compounds **1–2** showed an anti-cholesterol effect by the LB colorimetric assay. The discoveries of anti-cholesterol **1–2** were first reported. Insignificant effect of **2** than **1** caused by orientation of the hydroxyl group that probably interacts with other functional group through hydrogen bonding formation and therefore the activity was reduced. Unlike compound **2**, intramolecular reaction of **1** does not occur. These results suggested that the additional of hydroxyl group of **2** at the position C-19 is a critical point for important structural feature for anti-cholesterol activity in triterpenoid structures.

Acknowledgments

We thank Dr. Sofa Fajriah and Dr. Ahmad in the research center for chemistry, Indonesian Science Institute for NMR measurements. We are grateful to Mr. Irfan Junaedi in the center laboratory of Sekolah Tinggi Analisis Kimia Cilegon for the biological assay.

References

- [1] Hutahean, S.; Tanjung, M.; Sari, D. P.; Ningsih, V. E. Antihyperglycemic and Antihyperlipidemic Effects of Pirdot (*saurauia Vulcani* Korth.) Leaves Extract in Mice. *IOP Conf. Ser. Earth Environ. Sci.* **2018**, *130*(1), 012042. DOI: 10.1088/1755-1315/130/1/012042.
- [2] Situmorang, R. O. P.; Harianja, A. H.; Silalahi, J. Karo's Local Wisdom: The Use of Woody Plants for Traditional Diabetic Medicines. *Indo J. Forest Res.* **2015**, *2*(10), 121–130. DOI: 10.20886/ijfr.2015.2.2.830.121-130.
- [3] Ginting, G. A.; Rosidah, R.; Sitorus, P.; Satria, D. Wound Healing Activity of *Saurauia Vulcani* Korth. Aqueous Leaves Extract Evaluation on Excision Wound in Hyperglycemia Rats. *JIBS.* **2018**, *5*(3), 52–57.
- [4] Silalahi, M.; Supriatna, J.; Walujo, E. B.; Nisyawati, N. Local Knowledge of Medicinal Plants in Sub-ethnic Batak Simalungun of North Sumatera, Indonesia. *Biodiversitas.* **2015**, *16*, 44–54. DOI: 10.13057/biodiv/d160106.
- [5] Sitorus, P.; Rosidah, R.; Satria, D. Hypoglycemic Activity of Ethanolic Extract of *Saurauia Vulcani* Korth. Leaves. *Asian J. Pharm. Clin. Res.* **2018**, *11*, 35–36. DOI: 10.22159/ajpcr.2018.v11s1.26561.
- [6] Saragih, R. R. *Phytochemical Screening, Antioxidant and Antibacterial Activities of Methanol and Etyl Acetate Extracts of the Leaves of Pirdot (saurauia Vulcani Korth.) From Tigarunggu Region*; Universitas Sumatera Utara: Medan, 2018.
- [7] Suparman, A. C.; Kadarusman, M.; Situmeang, B. Triterpenoid Compound from Pirdot Plant (*sauralia* Sp). *J. Itekmata.* **2018**, *3*(1), 12–19.
- [8] Sheftel, A. G. Determination of Total and Free Cholesterol. *J. Lab. Clin. Med.* **1944**, *29*(8), 875–878.
- [9] Uddin, G.; Wairulah; Siddiqui, B. S.; Alam, S.; Sadat, A.; Ahmad, A.; Uddin, A. Chemical Constituent and Phytotoxicity of Solvent Extracted Fraction of Steam Bark of *Grewia Optiva* Drummond Ex Burret. *Middle-Eas J. Sci. Res.* **2011**, *8*(1), 85–91.
- [10] Martins, D.; Carrion, L. L.; Ramos, D. F.; Salome, K. S.; Da Silva, P. E. A.; Barison, A.; Nunez, C. V. Triterpenes and the Antimycobacterial Activity of *Duroia Macrophylla* Huber (rubiacaceae). *Biomed Res. Int.* **2013**, *2013*, article ID 605831. DOI:10.1155/2013/605831.
- [11] Ayatollahi, A. M.; Ghanadian, M.; Afsharypour, S.; Abdella, O. M.; Mirzai, M.; Askari, G. Pentacyclic Triterpenes in *Euphorbia Microsciadia* with Their T-cell Proliferation Activity. *Iran. J. Pharm. Res.* **2011**, *10*(2), 287–294.
- [12] Kenny, A. P. The Determination of Cholesterol by the Liebermann-burchard Reaction. *Biochem. J.* **1952**, *52*(4), 611–619.
- [13] Xiong, Q.; Wilson, W. K.; Pang, J. The Liebermann-burchard Reaction: Sulfonation, Desaturation, and Rearrangement of Cholesterol in Acid. *Lipids.* **2007**, *42*(1), 87–96. DOI: 10.1007/s11745-006-3013-5.
- [14] Barbour, H. M. Enzymatic Determination of Cholesterol and Triglyceride with the Abbott Bichromatic Analyser. *Ann. Clin. Biochem.* **1997**, *14*(1), 22–28. DOI: 10.1177/000456327701400104.
- [15] Liebermann, C. Ueber das Oxychinoterpen. *Dtsch. Chem. Ges.* **1885**, *18*, 1803–1809.
- [16] Burchard, H. Beitrage zur kenntnis des cholesterins. Inaugural-dissertation, universitat Rostock. **1889**, 61-I, 25–27.

Anti-cholesterol triterpenoid acids from *Saurauia vulcani* Korth. (Actinidiaceae)

ORIGINALITY REPORT

26%

SIMILARITY INDEX

17%

INTERNET SOURCES

26%

PUBLICATIONS

21%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Julinton Sianturi, Mayshah Purnamasari, Darwati, Desi Harneti, Tri Mayanti, Unang Supratman, Khalijah Awang, Hideo Hayashi. "New bisamide compounds from the bark of *Aglaia eximia* (Meliaceae)", *Phytochemistry Letters*, 2015
Publication 4%
 - 2 De Ji, Yun-Fei LIU, Qiao-Han WANG, Chun-Feng ZHANG, Zhong-Lin YANG. "Isolation, structure characterization and quantification of related impurities in asperosaponin VI", *Chinese Journal of Natural Medicines*, 2013
Publication 2%
 - 3 repositori.upf.edu
Internet Source 2%
 - 4 Awang, Khalijah, Xe-Min Loong, Kok Hoong Leong, Unang Supratman, Marc Litaudon, Mat Ropi Mukhtar, and Khalit Mohamad. "Triterpenes and steroids from the leaves of *Aglaia exima* (Meliaceae)", *Fitoterapia*, 2012.
Publication 2%
-

5	synapse.koreamed.org Internet Source	2%
6	www.thieme-connect.com Internet Source	1%
7	www.mdpi.com Internet Source	1%
8	Karuppiah Pillai Manoharan. "Triterpenoids from <i>Eugenia grandis</i> : structure elucidation by NMR spectroscopy", <i>Magnetic Resonance in Chemistry</i> , 03/2007 Publication	1%
9	Sianturi, Julinton, Desi Harneti, Darwati, Tri Mayanti, Unang Supratman, and Khalijah Awang. "A New (–)-5',6-dimethoxyisolariciresinol-(3'',4''-dimethoxy)-3 α -O- β -D-glucopyranoside from the bark of <i>Aglaia eximia</i> (Meliaceae)", <i>Natural Product Research</i> , 2016. Publication	1%
10	uad.portalgaruda.org Internet Source	1%
11	ojs.jmolekul.com Internet Source	1%
12	Zhou, Guo-Ping, Yan Yu, Ming-Ming Yuan, Tengfei Ji, Hui-Zheng Fu, and Rui-Jian Zhong. "Four New Triterpenoids from <i>Callicarpa kwangtungensis</i> ", <i>Molecules</i> , 2015. Publication	1%

13	Nurhayati Bialangi, Adam Mustapa, Yusda Salimi, Ari Widianoro, Boima Situmeang. "Isolation of Steroid Compounds from Suruhan (<i>Peperomia pellucida</i> L. Kunth) and Their Antimalarial Activity", <i>Asian Journal of Chemistry</i> , 2018 Publication	1%
14	umexpert.um.edu.my Internet Source	1%
15	Sianturi, Julinton, Mayshah Purnamasari, Darwati, Desi Harneti, Tri Mayanti, Unang Supratman, Khalijah Awang, and Hideo Hayashi. "New bisamide compounds from the bark of <i>Aglaia eximia</i> (Meliaceae)", <i>Phytochemistry Letters</i> , 2015. Publication	1%
16	Guo Sun, Qin Dai, Hong-Xia Zhang, Zhi-Jian Li, Zhi-Zhi Du. " New Sweet-Tasting C21-Pregnane Glycosides from Pericarps of ", <i>Journal of Agricultural and Food Chemistry</i> , 2016 Publication	1%
17	Submitted to Universiti Teknologi Malaysia Student Paper	1%
18	Submitted to UIN Syarif Hidayatullah Jakarta Student Paper	1%
19	Hiroaki Itoh, Kotaro Tokumoto, Takuya Kaji, Atmika Paudel, Suresh Panthee, Hiroshi	1%

Hamamoto, Kazuhisa Sekimizu, Masayuki
Inoue. "Total Synthesis and Biological Mode of
Action of WAP-8294A2: A Menaquinone-
Targeting Antibiotic", The Journal of Organic
Chemistry, 2017

Publication

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On