

Antibacterial and Phytochemical Activity Test of Brown Macroalgae Extract towards *Vibrio Algynoliticus* Bactery through in- litre Fertilization

by Juliana Jane L

Submission date: 05-Nov-2019 07:58PM (UTC+1000)

Submission ID: 1207456882

File name: Jurnal_cek_Turnitin.pdf (63.98K)

Word count: 3138

Character count: 17957

Antibacterial and Phytochemical Activity Test of Brown Macroalgae Extract towards *Vibrio Alginolyticus* Bacteria through in- vitro Fertilization

'Juliana, 'Jane L. Dangeubun and 'Diana Y. Syahailatua

'State University of Gorontalo Indonesia, Polytechnic State Fisheries of Tual,
Gorontalo, Indonesia

'Ray a Satheam Street, km 5, 0911-21377 Maluku Tenggara, Indonesia

Abstract: In the long term this research aims at finding a method to enhance the control of bacterial diseases in a safer and environmentally friendly fish farming. Meanwhile, the special target to attain is to obtain active antibacterial compounds through in vitro fertilization based on solvent polarity level and brown macroalgae active extract test through in vitro fertilization which can be used to increase fish survival rate in controlling bacterial diseases in fish farming. The extraction with methanol solvent results in the highest crude extract in all extracted seaweeds. This is justifiable because the greatest yield is those resulted by methanol solvent. We can see that seaweed species of *Sargassum polygocystum*, *S. cristaeifolium*, *S. hemphyllium*, turbinaria ornate and padina australis can produce the highest inhibition zone of 12.0 mm compared to other seaweed species extract in 100% concentration which is then followed by *S. cristaeifolium* methanol extract, turbinaria water extract and Padina australis water extract. Based on qualitative test chemical compound content of brown seaweed species, they contain phenolic, flavonoid, steroid and tannin compound. Whilst all qualitative tested seaweed contain no saponin. Thus, it is highly possible to develop those compounds as natural antibacterial and immunostimulant.

Key words: Antibacterial and phytochemical. macroalgae. in-vitro fertilization. antibacterial. qualitative

INTRODUCTION

Seaweed contains primary and secondary metabolite. The primary metabolites are vitamin, mineral, fibre, alginate and karaginan to be made as cosmetics substance for skin treatment. Besides its economical primary content, the secondary metabolite content of seaweed can be potentially produced as various bioactive metabolites with wide range of activities such as antibacterial, antiviral, antifungal and cytotoxic.

Green, red, or brown seaweed are potential resources of bioactive compound highly beneficial for developing pharmaceutical industry as antibacterial, anti tumor, anticancer or as reversal agent and agrochemical industry primarily for antifungal, fungicide and herbicide (Bijanti, 2005). According to Kordi, seaweed is greatly utilized by coastal communities as external medicine such as natural antiseptic. Research finding of Pringgienies shows seaweed potential as pathogens antibacterial inducing infectious diseases. One of such infectious diseases which commonly infect farmed fish is that causing scarlet fever disease.

Fish illness is one of the obstacles of fish farming industry. This is so because epidemic can trigger massive

fish mortality or farmed shrimp. The high rate of farmed fish mortality can decrease fish production which also decreases income as compared to the expenses for fish farming such as fish fry purchasing, fish feed purchasing, fishpond making, wage expenses and et cetera besides, the ill fish will only have lower sale value than that of healthy fish, especially for fresh fish selling.

On the basis of its cause, the fish diseases are divided into two categories namely infecting and non infecting diseases. Infecting diseases are diseases caused by pathogen infection into the wet mother body. The pathogenic diseases of fish are virus, bacteria, parasites and fungus. Meanwhile, non infectious diseases are diseases caused by other than pathogenic infection, such as environmental degrading quality, malnutrition and genetic physical defect. All these things can occur either in freshwater fish farming or in sea fish farming.

The control towards bacterial diseases in either freshwater fish farming or sea fish farming has been done all this time by using antibiotics. Antibiotics usage is beneficial when used appropriately based on appropriate diagnosis and dosage, easy to obtain and has stronger and faster visible effect. However, consistent antibiotics application can trigger resistance, leave residue in fish

body and pollute the environment which then can poison non targeted organism. Therefore, it is highly important to search other safer and more effective and environmentally friendly method to control fish diseases at safe level. *V. alginolyticus* and *aeromonas hydrophila* bacteria are one of pathogenic found in both sea water and freshwater causing fish diseases and death. One of alternatives to solve the problem is by utilizing active compound of brown macroalgae which are safe and environmentally friendly.

One of the successful diseases control attempt is the usage of antimicrobe compound made of marine algae using crude extract of *Padina australis* which is given by effective immersion in mouse seafish medication which are affected by *V. alginolyticus* bacteria by increasing mouse seafish survival rate to 100%. It is greatly assumed that the extract contains phenol compound which are lethal for *V. alginolyticus* bacteria. It is stated by Xiao-Jun, Chkhikvishvili and Ramazanov (2000) that *Sargassum furcatum*, *Dictyota* sp, *Sargassum desfontainii*, *Padina pavonica* and other kind of brown algae contain phenol compound such as florotannin a kind of tannin which is potential for antibacterial.

On the basis of this research, it is highly assumed that brown macroalgae is highly potential to be made as natural medicine substance in controlling vibriosis and aeromonas disease. Thus, it is necessary to conduct antibacterial activity test with the lowest dosage of several kinds of brown algae in South East Maluku water area which has never been analyzed academically to obtain bioactive compound which can increase fish survival rate and antibacterial activity in controlling fish disease.

Special research objective: The general objective of this research is to find safer and environmentally friendly bacterial disease control method in fish farming which can function as medicine in disease control of fish seed by using some brown algae. Whilst, the special research objectives are. Knowing the appropriate solvent based on polarity level to obtain bioactive compound of brown algae *Padina australis*, *Padina tetrastomatika*, *Sargassum polycystum*, *Sargassum cristaeifolium*, *Dictyota dichotoma*, *Turbinaria ornata*, *Turbinaria decurrens*, *Hydroclathrus clathratus* which function as antibacterial.

Searching for active compound of brown algae *Padina australis*, *Padina tetrastomatika*, *Sargassum polycystum*, *Sargassum cristaeifolium*, *Dictyota dichotoma*, *Turbinaria ornata*, *Turbinaria decurrens*, *Hydroclathrus clathratus* which have antibacterial quality and can be used in fish farming safely and environmentally friendly.

MATERIALS AND METHODS

The research can be done in two stages. The first stage aims at extracting active compound of *Medina australis*, *Padina tetrastomatika*, *Sargassum polycystum*, *Sargassum cristaeifolium*, *Sargassum olgocystum*, *Dictyota dichotoma*, *Turbinaria ornata*, *Hydroclathrus clathratus* which are antibacterial to obtain the lowest concentration of the active compound to resist *V. alginolyticus* bacteria and *A. hydrophila* through *in vitro* fertilization. In order to know the chemical compound of brown algae, we conduct phytochemical test which is antibacterial for fish. The whole steps of the first stage is conducted in basic chemical laboratory and pathological laboratory in state fishing polytechnic.

Extraction, phytochemical active compound of brown algae shows antibacterial characteristic and resistability test towards *V. alginolyticus* and *A. hydrophila* bacteria through *in vitro*. Sample collection and extraction of brown algae. At this stage, the researcher conducts several steps including sample preparation and active substance extraction. At the preparation stage, brown algae is taken in dry condition in South East Maluku and crushed by cutting them into tiny parts into powder. The next step is active substance extraction. The method used for extraction is Harbon method in 198° which has been modified using three kinds of solvents based on its polarity level.

Antibacterial test: The three produced thick extract (extract n hexana, dycoloromethan and methanol) undergo antibacterial test to find which extract can actively resist bacteria. This test use disk testing where sterilized disk paper is immersed in each extract. After 15-30 min or some, the disk will be attached to TCBS media which has been inoculated with *V. alginolyticus* bacteria. The measurement is conducted during incubation period for 24 h at 25°C by observing the existence of clear zone formed around the disk paper.

Phytochemical test of brown algae species potential As antibacterial

Alkaloids compound (culvenor-fitzgerul method): The 4 grams of brown powdered algae is crushed by using crusher, then little chloroform is added until it become pasta. Afterwards, 10 mL of ammoniac-chloroform of 0,05 N is added and is crushed again, the layer of 10 mL H₂SO₄ 2N is formed and strongly shaken. Then, it is cooled until it forms two layers. Then, a fleec of cotton is inserted in the tip of pipette to filter it. Sulphate acid layer is taken and poured into small reaction tube. Filtrate is tested by mayer reactor. The formation of white deposit with mayer reactor shows the existence of alkaloid

Flavanoid compound analysis (shinoda method cyanidin test): About 0.5 mg powdered sample is extracted with 5 mL methanol and is heated for 5 min in tube reaction. The extract is added with some drops of thick HCl and a little bit of magnesium powder. If it change color into red or yellow it means the extract contains flavanoid.

Saponin compound analysis (foam test): For the saponin test it is suggested to use dried sample because the test used is foam test. The dried sample is crushed and is poured into reaction tube and is added with 10 ml distilled water and is boiled for 2-3 min. Afterwards it is cooled and shaken powerfully. Constant foam for 5 min means an existence of saponin content.

Polyphenol compound analysis: mL, extract (ethanol, n-hexan-ethanol) is added with FeCl₃ 1%. Terpenoida compound is signed with the emergence of blue color, black, or purple. Terpenoid and steroid analysis (Lieberman-burchard 'method) several drops of chloroform in alkaloid test is placed on drop plates and is added with 3 drops of anhydride acetat then is dried and added with 3 drops of thick H₂SO₄.

The observed parameter: The observed parameter in this research is clear zone diameter of each extract and chemical content of various species of brown algae.

RESULTS AND DISCUSSION

Extraction of active substance in brown algae. The extraction result of brown algae can be seen in Table 1. The resulted extract from extraction process of various species of brown algae differ variously based on solvent kind used such as methanol extract, ethyl acetat extract and extract n-hexan with comparison (1:3). The yield of each extract can be seen in Table 1. Yield is the comparison of extract weight resulted with first weight of used substance and is stated in percentage (%). Extraction of brown seaweed. Extraction with methanol solvent can produce the highest crude extract in all extracted seaweed. This is justiciable because the highest yield produced is yield produced with methanol solvent. Thus in line with the statement stating that methanol can extract organic compound, some part of fat and tannin causing great methanol extract (Heath and Remeccius, 1987). The extraction result is affected by several factors namely natural condition of natural resources, extraction method, size of particle sample and condition and length of sample storage. This is so because during maceration there is a mixing of extracted substances which enlarge the possibility of collision between particles causing cell splitting with the hope that the expected component can come out of network substance and dissolve in the

Table 1: Yield result of crude extract of sargasum cristafolium, ethyl acetat and n-hexan

| Kind of RL | Dried weight (gram) | Rendemen (%) | | |
|--------------------------|---------------------|--------------|--------------|---------|
| | | Methanol | Ethyl acetat | n-hexan |
| <i>S.cristaeofolium</i> | 100 | 6.91 | 1.50 | 1.0 |
| <i>S.polyvestum</i> | 100 | 11.71 | 3.70 | 2.7 |
| <i>S.hemiphyllum</i> | 100 | 9.80 | 5.21 | 2.5 |
| <i>P.tetrastomatica</i> | 100 | 8.61 | 4.53 | 2.4 |
| <i>P.australis</i> | 100 | 9.70 | 6.40 | 3.5 |
| <i>Turbinaria ornata</i> | 100 | 8.50 | 3.40 | 2.5 |

Table 2: Phytochemical test of brown algae species crude extract

| Secondary/metabolit | Test method | A | B | C | D |
|---------------------|---|---|---|---|---|
| Phenolic | FeCl ₃ : 5% reactor t | + | + | + | + |
| Flavonoid | Thick HCl+Mg reactor | - | - | - | - |
| | H ₂ SO ₄ 2N reactor | + | + | + | + |
| | NaOH 10% reactor | + | + | + | + |
| Steroid | Lieberman-burchard reactor | + | + | + | + |
| Triterpenoid | | - | - | - | - |
| Saponin | HCl+H ₂ O reactor | - | - | - | - |
| Tanin | FeCl ₃ 1% reactor | + | + | + | - |

Explanation: A = Ethyl Acetat extract of sargasum cristaeofolium, B = Hexana extract of sargasum cristaeofolium, C = Ethyl Acetat extract of sargasum olygocystum, D = Hexana extract of sargasum olygocystum

Table 3: The result of clear zone towards vibrio alginoliticus (Cm) Bactery

| Kind of substance | Concentration (%) | | | | |
|--------------------------------|-------------------|----|----|------|------|
| | 15 | 25 | 50 | 75 | 100 |
| Sargassum. Methanol 1:4 | - | - | - | - | - |
| Sargassum cristafolium | - | - | - | - | - |
| Sargassum olygocystum methanol | - | - | - | 0.6 | 1.00 |
| Sargassum oligocystum. N-Hexan | - | - | - | - | 1.20 |
| Sargassum. hemiphyllum N-Hexan | - | - | - | - | - |
| Turbinaria ornate aguades | - | - | - | 0.6 | 1.00 |
| Padina australis aguades | - | - | - | 0.65 | 1.00 |
| Padina tetrastomatica | - | - | - | - | - |

solvent and in order to magnify fastering and reaction between active substance component with used solvent (Gaspez, 1991).

Phytochemical test of brown seaweed extract: To develop *A. acuminata* as antibacterial and immunostimulant, it is suggested to know the chemical compound of the extract from several kinds of seaweed extract. The gualitative test of chemical compound content extract of brown algae.

Based on the chemical compound content gualitative testof various brown seaweed as shown in Table 2, it is known that all kinds of brown seaweed as seen in Table 3 we can tell that all kinds of brown seaweed contain phenolic, flavonoid and steroid, steroid and tannin compound. Whulst, all kinds of gualitatively tested brown seaweed contains no sapponin. Therefore, it is highly possible that these compounds develop as antibacterial and natural immunostimulant.

According to Satria (2005), flavonoid compound can function as antioxidants by resisting kinds of coxidation reaction and can reduce hydroxyl,

superoxide and radical peroxy. Antioxidant compound captures free radicals, metal solidify and singlet oxygen formation reducer.

Phenol compound can also change surface tension which damages selective permeability of microbe membrane cell producing essential metabolic and inactivating bacterial system. The damage of this membrane enables nucleotide and amino acid to come out of cell. Besides, the damage can also prevent the intrusion of essential substances membrane cell also controls active transportation into the cell. This can cause bacterial cell death or resist bacterial development (Volk dan Wheeler, 1988).

Tannin compound contained in Widuri leave also can function as antimicrobe. Kim and Fung state that tannin can formulate complex protein with protein and hydrophobic interaction. When hydrogen and tannin and protein are bound, it is possible that the protein will be sedimented. This phenomenon is well known as protein denaturation. When enzyme protein of microbe is denaturated, the enzyme will be inactive so that the microbe metabolism will be disturbed which will cause cell damage.

Steroid working mechanism in impeding microbe is by damaging plasma membrane which leak out cytoplasm cell and leads to cell death (Putra, 2007). Afterwards, according to Cowan (1999), phenolic compound is an antibacterial which disturb cytoplasm membrane function. Phenol is also a compound of OH group bound in aromatic ring. Phenolic is also secondary metabolite spread in plants. Phenolic compound in plants can be in the form of simple phenol, anthraquinone, phenol acid, coumarin, flavonoid, lignin and tannin.

Antibacterial activity test from brown seaweed extract towards *Vibrio alginolyticus* and *Aeromonas hydrophila* bacteria through *in-vitro* fertilization. In Table 3, we can tell that not all extract can impede *A. hydrophila* bacteria. It is known that *Sargassum oligocystum*, *S. cristaefolium*, *S. hemphylium*, *Turbinaria ornata* and *Padina australis* seaweed species can produce resistance zone even though, each extract produce different resistance zone. N-hexane extract of *Sargassum oligocystum* has the greatest resistance zone of 12.0 mm compared to other seaweed extracts in 100% concentration. Then, it is followed by *S. cristaefolium* methanol extract, *turbinaria* water extract, *Padina australis* water extract.

S. oligocystum N-Hexane extract has the highest resistance because, this extract contains phenol, flavonoid, steroid triterpenoid and tannin which collaborate to resist *V. alginolyticus* bacteria through *in vitro* fertilization. However, this resistance is still categorized as mild category. This is according to resistance

criteria for bacteria according to Stout as referred in Rachdiati who states that resistance zone with 5-10 mm average is included in mild criteria. Meanwhile, resistance zone of <5 mm is included in weak criteria. This is caused by different ability of each extract with different quality and quantity content and different media extract absorption system so that the number of various bacteria and pathogenicity level result in different resistance zone.

The resistance or killing power of an antimicrobe is greatly affected by many factors such as concentration factor. Darkuni states that the ability of an antimicrobe substance which can abolish the ability towards certain microorganism depends on the concentration of antimicrobe substance. In other words, antimicrobe substance in a bacterial environment greatly determine the bacteria survival ability. Therefore, there are certain bacteria which can survive and even have active metabolism in an antimicrobe environment. Volk and Wheeler (1988), state that the main factor to determine how antimicrobe substance can work effectively is concentration, duration for the substance to work, temperature and number of species and microorganism. Mallawa and Halid state that the measurement of growth resistance zone becomes the standard of bioactivity which is affected by many factors such as functional group activity, bacterial resistance of bioactive substance, active substance level and tested bacterial density.

CONCLUSION

Based on the result and analysis of the research, we can conclude that:

- All extracts can produce resistance zone. However, each extract produces different resistance zone, except that of water extract *Padina tetrastomatica*, which has no resistance power in all concentration.
- Based on qualitative test, the chemical compound content of many kinds of brown seaweed, we can tell that all kinds of brown seaweed contain phenolic, flavonoid and steroid, tannin and all kinds of qualitatively tested seaweed contain no saporin. Thus, it is possible to develop these compounds as antibacterial and natural immunostimulant.

SUGGESTION

It is necessary to develop further research about the kinds of brown algae with appropriate dosage as antibacterial active compound towards *V. alginolyticus*, *A. hydrophila* in tiger grouper fish and tilapia fish at laboratory scale.

REFERENCES

- Bijanti, R., 2005. Fish Hematology. Airlangga University Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia..
- Chkhikvishvili, I.D. and Z.M. Ramazanov, 2000. Phenolic substances of brown algae and their antioxidant activity. *Applied Biochem. Microbiol.*, 36: 289-291.
- Cowan, M.M., 1999. Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*, 12: 564-582.
- Gaspez, V., 1991. Experimental Design Methods Unutuk Agricultural Science and Biological Engineering, Bandung, Indonesia., Sciences.
- Heath, H.B. and G. Reinesccius, 1987. Flavour Chemistry and Technology. Von Nostrand Reinhold Co., New York, USA...
- Putra, I.N.K., 2007. Study of antimicrobial power plant extracts some material against microbial preservative nira nira vandall and compounds active ingredients. Postgraduate Program of Brawijaya University, University of Brawijaya, Malang, Indonesia.
- Satria, E., 2005. The antioxidant potential of the fruit flesh young and old fruit pulp gods crown (*Phaleria macrocarpa* (Scheff Boer). Master Thesis, Faculty of Mathematics and Natural Sciences, Bogor Agricultural Institute, Bogor, Indonesia.
- Volk, W.A. and M.F. Wheeler, 1993. Basic Microbiology. Sth Edn., Erlangga Press, Jakarta.

Antibacterial and Phytochemical Activity Test of Brown Macroalgae Extract towards Vibrio Algynoliticus Bactery through in- litre Fertilization

ORIGINALITY REPORT

7 %

SIMILARITY INDEX

0 %

INTERNET SOURCES

7 %

PUBLICATIONS

0 %

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

7%

★ J. L, Dangeubun, and Syahailatua D. Y.
"Phytochemistry And Antibacterial Activity Test From Methanol Extract of Alstonia Acuminata Tree Bark Against Vibrio Harveyi Bacterium in In-Vitro Manner", Journal of Biology and Life Science, 2014.

Publication

Exclude quotes Off

Exclude bibliography On

Exclude matches Off

Antibacterial and Phytochemical Activity Test of Brown Macroalgae Extract towards Vibrio Algynoliticus Bactery through in- litre Fertilization

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5