










AACL Bioflux

14

H Index

Country	Romania - IIII SJR Ranking of Romania
Subject Area and Category	Agricultural and Biological Sciences Aquatic Science Ecology, Evolution, Behavior and Systematics Environmental Science Management, Monitoring, Policy and Law Water Science and Technology
Publisher	Bioflux Publishing House
Publication type	Journals
ISSN	18449166, 18448143
Coverage	2006, 2009-2020
Scope	Information not localized
	 Join the conversation about this journal





AAAL Bioflux ← Show this widget in your own website

Q3

Aquatic Science

best quartile

SJR 2019

0.28

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```


powered by scimagojr.com

A **andrew** 3 months ago

I would like to know the paper publish in AAAL Bioflux, 2020, Volume 13, Issue 2. I would like to know if this paper indexed by Scopus, how about the level of this paper Q3 or Q4 and how about the SJR ?

Thank you for your help.

reply



[Aquaculture, Aquarium, Conservation & Legislation](#)

You are here > [Home](#)

AACL Bioflux

[Instructions to authors](#)

[Submission letter](#)

[Model of paper](#)

[Reviewer information pack](#)

[Editorial Board Expanded](#)

[Coverage / databases](#)

[Volume 13\(6\)/2020](#)

[Volume 13\(5\)/2020](#)

[Volume 13\(4\)/2020](#)

[Volume 13\(3\)/2020 \(June, 30\)](#)

[Volume 13\(2\)/2020 \(April, 30\)](#)

[Volume 13\(1\)/2020 \(February, 28\)](#)

[Volume 12\(6\)/2019 \(December, 30\)](#)

[Volume 12\(5\)/2019 \(October, 30\)](#)

[Volume 12\(4\)/2019 \(August, 30\)](#)

[Volume 12\(3\)/2019 \(June, 30\)](#)

[Volume 12\(2\)/2019 \(April, 30\)](#)

[Volume 12\(1\)/2019 \(February, 28\)](#)

[Volume 11\(6\)/2018 \(December, 30\)](#)

[Volume 11\(5\)/2018 \(October, 30\)](#)

[Volume 11\(4\)/2018 \(August, 30\)](#)

[Volume 11\(3\)/2018 \(June, 30\)](#)

[Volume 11\(2\)/2018 \(April, 30\)](#)

[Volume 11\(1\)/2018 \(February, 28\)](#)


[Volume 10\(6\)/2017 \(December, 30\)](#)

[Volume 10\(5\)/2017 \(October, 30\)](#)

[Volume 10\(4\)/2017 \(August, 30\)](#)

[Volume 10\(3\)/2017 \(June, 30\)](#)

About us




BIOFLUX SRL

Identity: Research organization, CAEN 7219, CUI: 23753059 from 18.04.2008

Address: 54 Ceahlau street, Cluj-Napoca 400488, Romania,

European Union, Europe.



AACL Bioflux

[Instructions to authors](#)

[Submission letter](#)

[Model of paper](#)

[Reviewer information pack](#)

[Editorial Board Expanded](#)

[Coverage / databases](#)

[Volume 13\(6\)/2020](#)

[Volume 13\(5\)/2020](#)

[Volume 13\(4\)/2020](#)

[Volume 13\(3\)/2020 \(June, 30\)](#)

[Volume 13\(2\)/2020 \(April, 30\)](#)

[Volume 13\(1\)/2020 \(February, 28\)](#)

[Volume 12\(6\)/2019 \(December, 30\)](#)

[Volume 12\(5\)/2019 \(October, 30\)](#)

[Volume 12\(4\)/2019 \(August, 30\)](#)

[Volume 12\(3\)/2019 \(June, 30\)](#)

[Volume 12\(2\)/2019 \(April, 30\)](#)

[Volume 12\(1\)/2019 \(February, 28\)](#)

[Volume 11\(6\)/2018 \(December, 30\)](#)

[Volume 11\(5\)/2018 \(October, 30\)](#)

[Volume 11\(4\)/2018 \(August, 30\)](#)

[Volume 11\(3\)/2018 \(June, 30\)](#)

[Volume 11\(2\)/2018 \(April, 30\)](#)

[Volume 11\(1\)/2018 \(February, 28\)](#)

[Volume 10\(6\)/2017 \(December, 30\)](#)

[Volume 10\(5\)/2017 \(October, 30\)](#)

[Volume 10\(4\)/2017 \(August, 30\)](#)

[Volume 10\(3\)/2017 \(June, 30\)](#)

Aquaculture, Aquarium, Conservation & Legislation - International Journal of the Bioflux Society

ISSN 1844-9166 (online)

ISSN 1844-8143 (print)

Published by Bioflux - bimonthly -

in cooperation with The Natural Sciences Museum Complex (Constanta, Romania)

Peer-reviewed (each article was independently evaluated before publication by two specialists)

The journal includes original papers, short communications, and reviews on Aquaculture (Biology, Technology, Economics, Marketing), Fish Genetics and Improvement, Aquarium Sciences, Fisheries, Ichthyology, Aquatic Ecology, Conservation of Aquatic Resources and Legislation (in connection with aquatic issues) from wide world.

The manuscripts should be submitted to zoobiomag2004@yahoo.com

Editor-in-Chief:

Petrescu-Mag I. Valentin: USAMV Cluj, Cluj-Napoca, University of Oradea (Romania)

Gavriloaie Ionel-Claudiu (reserve): SC Bioflux SRL, Cluj-Napoca (Romania).

Editors:

Abdel-Rahim Mohamed M.: National Institute of Oceanography and Fisheries, Alexandria (Egypt)

Adascalitei Oana: Maritime University of Constanta, Constanta (Romania)

Amira Aicha Beya: Badji Mokhtar Annaba University, Annaba (Algeria)

Arockiaraj A. Jesu: SRM University, Chennai (India)


Appelbaum Samuel: Ben-Gurion University of the Negev (Israel)

Baharuddin Nursalwa: Universiti Malaysia Terengganu, Terengganu (Malaysia)


Boaru Anca: USAMV Cluj, Cluj-Napoca (Romania)

Volume 10(2)/2017 (April, 30)	Botha Miklos: Bioflux SRL, Cluj-Napoca (Romania)
Volume 10(1)/2017 (February, 28)	Breden Felix: Simon Fraser University (Canada)
Volume 9(6)/2016 (December, 30)	Burny Philippe: Universite de Liege, Gembloux (Belgium)
Volume 9(5)/2016 (October, 30)	Caipang Cristopher M.A.: Temasek Polytechnic (Singapore)
Volume 9(4)/2016 (August, 30)	Chapman Frank: University of Florida, Gainesville (USA)
Volume 9(3)/2016 (June, 30)	Coroian Cristian: USAMV Cluj, Cluj-Napoca (Romania)
Volume 9(2)/2016 (April, 30)	Creanga Steofil: USAMV Iasi, Iasi (Romania)
Volume 9(1)/2016 (February, 28)	Cristea Victor: Dunarea de Jos University of Galati, Galati (Romania)
Volume 8(6)/2015 (December, 30)	Das Simon Kumar: Universiti Kebangsaan Malaysia, Bangi, Selangor (Malaysia)
Volume 8(5)/2015 (October, 30)	Dimaggio Matthew A.: University of Florida (USA)
Volume 8(4)/2015 (August, 30)	Dimaggio Matthew A.: University of Florida (USA)
Volume 8(3)/2015 (June, 30)	Firica Cristian Manuel: Spiru Haret University Bucharest, Craiova (Romania)
Volume 8(2)/2015 (April, 30)	Georgescu Bogdan: USAMV Cluj, Cluj-Napoca (Romania)
Volume 8(1)/2015 (February, 28)	Karayucel Ismihan: University of Sinop, Sinop (Turkey)
Volume 7(6)/2014 (December, 30)	Khamesipour Faham: Shiraz University, Shiraz (Iran)
Volume 7(5)/2014 (October, 30)	Khamesipour Faham: Shiraz University, Shiraz (Iran)
Volume 7(4)/2014 (August, 30)	Kosco Jan: Presov University, Presov (Slovakia)
Volume 7(3)/2014 (June, 30)	Kovacs Eniko: USAMV Cluj, Cluj-Napoca (Romania)
Volume 7(2)/2014 (April, 15)	Mehrad Bahar: Gorgan University of Agricultural Sciences and Nat. Res. (Iran)
Volume 7(1)/2014 (February, 15)	Miclaus Viorel: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(6)/2013 (November, 15)	Mihociu Tamara: R&D National Institute for Food Bioresources (Romania)
Volume 6(5)/2013 (September, 15)	Molnar Kalman: Hungarian Academy of Sciences, Budapest (Hungary)
Volume 6(4)/2013 (July, 25)	Muchlisin Zainal Abidin: Universiti Sains (Malaysia), Syiah Kuala University (Indonesia)
Volume 6(3)/2013 (May, 15)	Muntean George Catalin: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(2)/2013 (March, 15)	Muntean George Catalin: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(1)/2013 (January, 15)	Nowak Michal: University of Agriculture in Krakow (Poland)
Volume 5(5)/2012 (December, 30)	Nyanti Lee: Universiti Malaysia Sarawak, Sarawak (Malaysia)
Volume 5(4)/2012 (September, 30)	Olivotto Ike: Universita Politecnica delle Marche, Ancona (Italy)
Volume 5(3)/2012 (July, 30)	Oroian Firuta Camelia: USAMV Cluj, Cluj-Napoca (Romania)
Volume 5(2)/2012 (June, 30)	Papuc Tudor: USAMV Cluj, Cluj-Napoca (Romania)
Volume 5(1)/2012 (March, 15)	Parvulescu Lucian: West University of Timisoara (Romania)
Volume 4(5)/2011 (December, 30)	Pasarin Benone: USAMV Iasi, Iasi (Romania)
Volume 4(4)/2011 (October, 30)	Pattikawa Jesaja Ajub: Pattimura University, Ambon (Indonesia)
Volume 4(3)/2011 (July, 30)	Petrescu Dacinia Crina: Babes-Bolyai University, Cluj-Napoca (Romania)

Volume 4(2)/2011 (April, 30)	Petrescu-Mag Ruxandra Malina: Babes-Bolyai University, Cluj-Napoca (Romania)
Volume 4(1)/2011 (January, 30)	Petrovici Milca: West University of Timisoara (Romania)
Volume 3(5)/2010 (December, 5)	Pratasik Silvester Benny: Sam Ratulangi University, Manado (Indonesia)
Volume 3(4)/2010 (December, 1)	Proorocu Marian: USAMV Cluj, Cluj-Napoca (Romania)
Volume 3(3)/2010 (November, 15)	Ray Sunuram: Khulna University (Bangladesh)
Volume 3(2)/2010 (July, 30)	Rhyne Andrew: Roger Williams University; New England Aquarium, Boston (USA)
Volume 3(1)/2010 (February, 28)	Ruchin Alexander B.: Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Saransk (Russia)
Volume 2(4)/2009 (October, 30)	Safirescu Calin: USAMV Cluj, Cluj-Napoca (Romania)
Volume 2(3)/2009 (July, 30)	Serrano Jr. Augusto E.: University of the Philippines Visayas (Philippines)
Volume 2(2)/2009 (April, 30)	Sima Nicusor Flaviu: USAMV Cluj, Cluj-Napoca (Romania)
Volume 2(1)/2009 (January, 30)	Tlusty Michael F.: New England Aquarium, Boston (USA)
Volume 1(2)/2008 (December, 30)	Vesa Stefan Cristian: Iuliu Hatieganu UMF, Cluj-Napoca (Romania)
Volume 1(1)/2008 (September, 30)	Vintila Iuliana: Dunarea de Jos University of Galati, Galati (Romania)
Volume Pilot/2007 (December, 30) - available printed only	Wariaghli Fatima: University Mohammed V in Rabat, Rabat (Morocco)
Pontus Euxinus, Volume 1 (1980) - Parent Journal	Yusli Wardiatno: Bogor Agricultural University, Bogor (Indonesia).



design: www.simple-webdesign.com



AACL Bioflux

Volume 11(3)/2018

Instructions to authors

First pages, AACL Bioflux 11(3):i-vi.

Submission letter

Nyanti L., Soo C. L., Danial-Nakhaie M. S., Ling T. Y., Sim S. F., Grinang J., Ganyai T., Lee K. S. P., 2018 Effects of water temperature and pH on total suspended solids tolerance of Malaysian native and exotic fish species. AACL Bioflux 11(3):565-575.

Model of paper

Reviewer information pack

Ompi M., Svane I., 2018 Comparing spawning, larval development, and recruitments of four mussel species (Bivalvia: Mytilidae) from South Australia. AACL Bioflux 11(3):576-588.

Editorial Board Expanded

Coverage / databases

Volume 13(6)/2020

Volume 13(5)/2020

Volume 13(4)/2020

Naorbe M. C., Serrano Jr. A. E., 2018 Effects of heavy metals on cell density, size, specific growth rate and chlorophyll a of *Tetraselmis tetraathele* under controlled laboratory conditions. AACL Bioflux 11(3):589-597.

Volume 13(3)/2020 (June, 30)

Salosso Y., Jasmanindar Y., 2018 Diversity of brown macroalgae in Kupang Bay waters and alginate content potential and its phytochemistry. AACL Bioflux 11(3):598-605.

Volume 13(2)/2020 (April, 30)

Volume 13(1)/2020 (February, 28)

Rameli N. M., Kader M. A., Aznan A. S., Musa N., 2018 Effect of *Cosmos caudatus* extract on antibacterial activity and lethality activity of brine shrimp. AACL Bioflux 11(3):606-612.

Volume 12(6)/2019 (December, 30)

Volume 12(5)/2019 (October, 30)

Volume 12(4)/2019 (August, 30)

Firdaus M., Lelono T. D., Saleh R., Bintoro G., Salim G., 2018 The expression of the body shape in fish species *Harpadon nehereus* (Hamilton, 1822) in the waters of Juata Laut, Tarakan city, North Kalimantan. AACL Bioflux 11(3):613-624.

Volume 12(3)/2019 (June, 30)

Volume 12(2)/2019 (April, 30)

Volume 12(1)/2019 (February, 28)

Adibrata S., Yulianda F., Boer M., Nurjaya I. W., 2018 Growth rate, mortality and exploitation for managing bar-cheeked coral trout grouper *Plectropomus maculatus* (Bloch, 1790) in Lepar Island and Pongok Island waters, South Bangka Regency, Indonesia. AACL Bioflux 11(3):625-634.

Volume 11(6)/2018 (December, 30)

Volume 11(5)/2018 (October, 30)

Volume 11(4)/2018 (August, 30)

Volume 11(3)/2018 (June, 30)

Paruntu C. P., Darwisito S., Rumengan A. P., Sinjal J. H., 2018 The effects of monoculture or polyculture of tiger grouper (*Epinephelus fuscoguttatus*) and rabbitfish (*Siganus canaliculatus*) on the growth performance of tiger grouper in floating net cage. AACL Bioflux 11(3):635-644.

Volume 11(2)/2018 (April, 30)

Volume 11(1)/2018 (February, 28)

Andrei (Guriencu) R. C., Cristea V., Cretu M., Dediu L., Docan A. I., 2018 The effect of feeding rate on growth performance and body composition of Russian sturgeon (*Acipenser gueldenstaedtii*) juveniles. AACL Bioflux 11(3):645-652.

Volume 10(6)/2017 (December, 30)

Volume 10(5)/2017 (October, 30)

Volume 10(4)/2017 (August, 30)

Volume 10(3)/2017 (June, 30)

- Volume 10(2)/2017 (April, 30)
- Volume 10(1)/2017 (February, 28)
- Volume 9(6)/2016 (December, 30)
- Volume 9(5)/2016 (October, 30)
- Volume 9(4)/2016 (August, 30)
- Volume 9(3)/2016 (June, 30)
- Volume 9(2)/2016 (April, 30)
- Volume 9(1)/2016 (February, 28)
- Volume 8(6)/2015 (December, 30)
- Volume 8(5)/2015 (October, 30)
- Volume 8(4)/2015 (August, 30)
- Volume 8(3)/2015 (June, 30)
- Volume 8(2)/2015 (April, 30)
- Volume 8(1)/2015 (February, 28)
- Volume 7(6)/2014 (December, 30)
- Volume 7(5)/2014 (October, 30)
- Volume 7(4)/2014 (August, 30)
- Volume 7(3)/2014 (June, 30)
- Volume 7(2)/2014 (April, 15)
- Volume 7(1)/2014 (February, 15)
- Volume 6(6)/2013 (November, 15)
- Volume 6(5)/2013 (September, 15)
- Volume 6(4)/2013 (July, 25)
- Volume 6(3)/2013 (May, 15)
- Volume 6(2)/2013 (March, 15)
- Volume 6(1)/2013 (January, 15)
- Volume 5(5)/2012 (December, 30)
- Volume 5(4)/2012 (September, 30)
- Volume 5(3)/2012 (July, 30)
- Volume 5(2)/2012 (June, 30)
- Volume 5(1)/2012 (March, 15)
- Volume 4(5)/2011 (December, 30)
- Volume 4(4)/2011 (October, 30)
- Volume 4(3)/2011 (July, 30)
- Amira A. B., Bougdah M., 2018 Influence of Mafragh and Seybouse inputs (sediment and salts) on the productivity of Annaba Bay. AACL Bioflux 11(3):653-665.**
- Wiyono E. S., Raharjo S. S. S., Permana S. M., 2018 Fishermen acceptance on introduction of fishing technology: perception and its development strategies. AACL Bioflux 11(3):666-677.**
- Ahmadi, Muhammad, Emmy Lilimantik, 2018 Phototactic response of climbing perch *Anabas testudineus* to different colors and light pattern of LED light traps. AACL Bioflux 11(3):678-689.**
- Krome C., Jauncey K., Lohaus G., Focken U., 2018 Phytate analysis and phytase application in *Jatropha curcas* kernel meal for use in aquaculture feeds. AACL Bioflux 11(3):690-700.**
- Paul A. K., Alam M. M., Islam M. S., Hussain M. A., Das S. K., 2018 Feeding behaviour of mud crab *Scylla serrata* in north of Sundarbans, Bangladesh. AACL Bioflux 11(3):701-708.**
- Diaf A., Benchikh N., Bouhali F. Z., Ladaimia S., Dahel A., Djebar A. B., 2018 Structure of the ichthyological stands in the Gulf of Annaba, Algerian east coast. AACL Bioflux 11(3):709-723.**
- Collazos-Lasso L. F., Gutierrez-Espinosa M. C., Aya-Baquero E., 2018 Induced reproduction of the sailfin pleco, *Pterygoplichthys gibbiceps* (Kner, 1854) (Pisces: Loricariidae). AACL Bioflux 11(3):724-729.**
- Benchikh N., Diaf A., Ladaimia S., Bouhali F. Z., Dahel A., Djebar A. B., 2018 European anchovy *Engraulis encrasicolus* (Linnaeus, 1758) from the Gulf of Annaba, east Algeria: age, growth, spawning period, condition factor and mortality. AACL Bioflux 11(3):730-743.**
- Sartimbul A., Rohadi E., Ikhsani S. N., Listiyaningsih D., 2018 Morphometric and meristic variations among five populations of *Sardinella lemuru* Bleeker, 1853 from waters of Bali Strait, northern and southern-east Java and their relation to the environment. AACL Bioflux 11(3):744-752.**
- Ardiansyah, Dahlia, Hartinah, Ibrahim, Wahidah, 2018 Improvement of the nutritive quality of *Sargassum* powder through *Aspergillus niger*, *Saccharomyces cerevisiae*, and *Lactobacillus* spp. fermentations. AACL Bioflux 11(3):753-764.**
- Kepel R. C., Mantiri D. M. H., Paransa D. S. J., Paulus J. J. H., Nasprianto, Wagey B. T., 2018 Arsenic content, cell structure, and pigment of *Ulva* sp. from Totok Bay and Blongko waters, North Sulawesi, Indonesia. AACL Bioflux 11(3):765-772.**
- Asadi M. A., Kemalawati R., Guntur G., Luthfi O. M., 2018 The PAH characterization and the acute toxicity of diesel oil and chemically dispersed diesel oil to *Litopenaeus vannamei*. AACL Bioflux 11(3):773-781.**
- El-Shafiey M. H. M., Mabroke R. S., Mola H. R. A., Hassaan M. S., Suloma A., 2018 Assessing the suitability of different carbon sources for Nile tilapia, *Oreochromis niloticus* culture in BFT system. AACL Bioflux 11(3):782-795.**

Volume 4(2)/2011 (April, 30)

Volume 4(1)/2011 (January, 30)

Volume 3(5)/2010 (December, 5)

Volume 3(4)/2010 (December, 1)

Volume 3(3)/2010 (November, 15)

Volume 3(2)/2010 (July, 30)

Volume 3(1)/2010 (February, 28)

Volume 2(4)/2009 (October, 30)

Volume 2(3)/2009 (July, 30)

Volume 2(2)/2009 (April, 30)


Volume 2(1)/2009 (January, 30)

Volume 1(2)/2008 (December, 30)

Volume 1(1)/2008 (September, 30)

Volume Pilot/2007 (December, 30) - available printed only

Pontus Euxinus, Volume 1 (1980) - Parent Journal



- Ompi M., Lomoindong F., Mandagi S. V., 2018 Effects of artificial and natural materials ropes as substrates for settling and metamorphosis of pediveliger pearl oyster, *Pinctada maxima* (Jameson, 1901). AACL Bioflux 11(3):796-803.
- Irnawati R., Surilayani D., Susanto A., Munandar A., Rahmawati A., 2018 Potential yield and fishing season of anchovy (*Stolephorus* sp.) in Banten, Indonesia. AACL Bioflux 11(3):804-809.
- Vincentius A., Nessa M. N., Jompa J., Saru A., Nurdin N., Rani C., 2018 Influential factors analysis towards mangrove cover and production of demersal fish in Maumere Bay, Indonesia. AACL Bioflux 11(3):810-822.
- Pratiwi F. D., Zainuri M., Purnomo P. W., Purwanti F., 2018 Evaluation of weevils (*Neochetina* spp.) existence in Rawa Pening Lake (abundance and impact on target and non-target plant). AACL Bioflux 11(3):823-832.
- Kadim M. K., Pasingi N., Kasim F., 2018 Spatial and temporal distribution of phytoplankton in the Gorontalo Bay, Indonesia. AACL Bioflux 11(3):833-845.
- El-Husseiny O. M., Goda A. M. A. S., Mabroke R. S., Soaudy M., 2018 Complexity of carbon sources and the impact on biofloc integrity and quality in tilapia (*Oreochromis niloticus*) tanks. AACL Bioflux 11(3):846-855.
- Taqwa F. H., Supriyono E., Budiardi T., Utomo N. B. P., Affandi R., 2018 Optimization of physiological status of glass eel (*Anguilla bicolor bicolor*) for transport by salinity and temperature acclimatization. AACL Bioflux 11(3):856-867.
- Effendy I. J., Hutabarat J., Ambariyanto A., Basuki F., 2018 Protein content and free amino acid composition of abalone (*Haliotis asinina*) broodstock fed by different fresh macroalgae and formulated diet. AACL Bioflux 11(3):868-876.
- Supriyantini E., Soenarjo N., Santosa G. W., Ridlo A., Sedjati S., Ambariyanto A., 2018 Effectiveness and efficiency of red seaweed *Gracilaria verrucosa* as biofilter in absorption of Pb in seawater. AACL Bioflux 11(3):877-883.
- Kantun W., Mallawa A., Tuwo A., 2018 Reproductive pattern of yellowfin tuna *Thunnus albacares* in deep and shallow sea FAD in Makassar Strait. AACL Bioflux 11(3):884-893.
- Bensaad-Bendjedid L., Belhaoues S., Kerdoussi A., Djebbari N., Tahri M., Bensouilah M., 2018 Age structure, growth and shell form of *Cerastoderma glaucum* (Bivalvia: Cardiidae) from El Mellah lagoon, Algeria. AACL Bioflux 11(3):894-904.
- Kusmana C., Sukwika T., 2018 Coastal community preference on the utilization of mangrove ecosystem and channelbar in Indramayu, Indonesia. AACL Bioflux 11(3):905-918.
- Limmon G. V., Rijoly F., Ongkers O. T. S., Loupatty S. R., Pattikawa J. A., 2018 Community structure of reef fish in the southern waters of Ambon Island, eastern Indonesia. AACL Bioflux 11(3):919-924.

Nikdehghan N., Kashiri H., Hedayati A. A., 2018 CuO nanoparticles-induced micronuclei and DNA damage in *Cyprinus carpio*. AACL Bioflux 11(3):925-936.


Bensafia N., Khati W., 2018 Seasonal variations in four trace metals (Cu, Zn, Pb, Cd) in sponges *Sarcotragus spinosulus* of the Gulf of Annaba, Northeast Algeria. AACL Bioflux 11(3):937-942.

Hasbullah D., Karim M. Y., Zainuddin, Trijuno D. D., 2018 The effect of salinity on the body chemical composition and RNA/DNA ratio of the hybrid brackishwater Nile tilapia *Oreochromis niloticus*. AACL Bioflux 11(3):943-947.

Chadijah A., Sulistiono, Haryani G. S., Affandi R., Mashar A., 2018 Species composition of *Telmatherina* caught in the vegetated and rocky habitats in Matano Lake, South Celebes, Indonesia. AACL Bioflux 11(3):948-955.

design: www.simple-webdesign.com

[Home](#) | [Archive](#) | [Volume 1 \(1\) / 2008](#) | [CEEX 140](#) | [Volume 1 \(2\) / 2008](#) | [Volume 2 \(1\) / 2009](#) | [Volume 2 \(2\) / 2009](#) | [Volume 2 \(3\) / 2009](#) | [Volume 2 \(4\) / 2009](#) | [Pilot](#) | [Volume 3 \(1\) / 2010](#) | [Volume 3\(2\)/2010 \(July, 30\)](#) | [Volume 3\(3\)/2010](#) | [Volume 3\(4\)/2010](#) | [Volume 3\(5\)/2010 - ACVAPEDIA 2010](#) | [Volume 4\(1\)/2011](#) | [Volume 4\(2\)/2011 - ACVAPEDIA 2010](#) | [Volume 4\(3\)/2011](#) | [Volume 4\(4\)/2011](#) | [Volume 4\(5\)/2011](#) | [Volume 5\(1\)/2012 \(March, 15\)](#) | [Volume 5\(2\)/2012](#) | [Volume 5\(3\)/2012](#) | [Volume 5\(4\)/2012](#) | [Volume 5\(5\)/2012 \(December, 30\)](#) | [Volume 6\(1\)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas \(HAKI\), 27-29th of November, 2012](#) | [Volume 6\(2\)/2013 - ACVAPEDIA 5th edn., Hungary, Szarvas \(HAKI\), 27-29th of November, 2012](#) | [Volume 6\(3\)/2013](#) | [Volume 6\(4\)/2013](#) | [Volume 6\(5\)/2013](#) | [Volume 6\(6\)/2013](#) | [Volume 7\(1\)/2014](#) | [Volume 7\(2\)/2014](#) | [Volume 7\(3\)/2014](#) | [Volume 7\(4\)/2014](#) | [Volume 7\(5\)/2014](#) | [Volume 7\(6\)/2014](#) | [Volume 8\(1\)/2015](#) | [Volume 8\(2\)/2015](#) | [Volume 8\(3\)/2015](#) | [Volume 8\(4\)/2015](#) | [Volume 8\(5\)/2015](#) | [Volume 8\(6\)/2015](#) | [Volume 9\(1\)/2016](#) | [Volume 9\(2\)/2016](#) | [Volume 9\(3\)/2016](#) | [Volume 9\(4\)/2016](#) | [Volume 9\(5\)/2016](#) | [Volume 9\(6\)/2016](#) | [Volume 10\(1\)/2017](#) | [Volume 10\(2\)/2017](#) | [Volume 10\(3\)/2017](#) | [Volume 10\(4\)/2017](#) | [Volume 10\(5\)/2017](#) | [Volume 10\(6\)/2017](#) | [Volume 11\(1\)/2018](#) | [Volume 11\(2\)/2018](#) | [Volume 11\(3\)/2018](#) | [Volume 11\(4\)/2018](#) | [Pontus Euxinus, Volume 1, 1980](#) | [Volume 11\(5\)/2018](#) | [Volume 11\(6\)/2018](#) | [Volume 12\(1\)/2019](#) | [Volume 12\(2\)/2019](#) | [Volume 12\(3\)/2019](#) | [Volume 12\(4\)/2019](#) | [Volume 12\(5\)/2019](#) | [Volume 12\(6\)/2019](#) | [Volume 13\(1\)/2020](#) | [Volume 13\(2\)/2020](#) | [Volume 13\(3\)/2020](#) | [Contact](#) | [Site Map](#)



Spatial and temporal distribution of phytoplankton in the Gorontalo Bay, Indonesia

Miftahul K. Kadim, Nuralim Pasingi, Faizal Kasim

Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Gorontalo City, Gorontalo Province, Indonesia.

Corresponding author: M. K. Kadim, miftahulhairkadim@ung.ac.id

Abstract. Phytoplankton has an important role in coastal ecosystems due to its ability to process photosynthesis as the base of producer for the aquatic food chain. The purpose of this study is to determine the spatial and seasonal distribution pattern of phytoplankton in Gorontalo Bay, Indonesia. The samples were taken from 5 sampling sites in the Gorontalo Bay, Indonesia in three different seasons: west monsoon season (in May), transition season (in June) and east monsoon season (in July). The results showed that ecological dynamics in the Gorontalo Bay was indicated by the various physics and chemical characteristics of waters spatially and seasonally. Based on the abundance of phytoplankton in the waters during the study, the east monsoon season was the most productive time in the Gorontalo Bay with an average of number of phytoplankton of 673026 cells L⁻¹ followed by transition and west monsoon season with the average counts of phytoplankton of 352704 and 3839 cells L⁻¹ respectively. Spatially, the highest abundance of phytoplankton was found in the station 3 with an average of 9891 cell L⁻¹, followed by station 4 with an average of 8770 cell L⁻¹, station 1 with an average of 7493 cell L⁻¹, station 2 with an average of 6343 cell L⁻¹, and, finally, station 5 with the average of 2651 cell L⁻¹.

Key Words: monsoon season, phytoplankton distribution, Gorontalo Bay.

Introduction. Tomini Bay is relatively fertile and rich in natural marine potential (Yusron & Edward 2000). It is also known as a marine tourism region and has a variety of potential pelagic fish resources (Wiadnyana 1998). The Bay of Gorontalo is part of Tomini Bay area which goes into the administrative area of Gorontalo City, Indonesia. The geographic conditions of the bay give consequences for the mass circulation of water between the waters within the bay and the surrounding as it is directly related to the Maluku Sea, Tolo Bay and the Celebes Sea (Setyadji & Priatna 2011). The existence of phytoplankton in the Gorontalo Bay is one of the ecological studies that need to be conducted regularly to monitor and ensure aquatic ecosystem balance.

Phytoplankton has an important role in coastal ecosystems due to its ability to process photosynthesis. Most of marine organisms depend on phytoplankton as a base producer for the aquatic food chain (Townsend et al 2000). Phytoplankton forms the essential base of the aquatic food web and plays a major role in aquatic productivity and ecosystem health (Nursuhayati et al 2013). The abundance and composition of phytoplankton in waters can be used as an indication of the fertility rate and water conditions changes (Karydis & Tsirtsis 1996; Thoha & Amri 2011; Radiarta 2013). Phytoplankton availability influences the distribution of consumer organisms especially the herbivore zooplankton. The high nutritional contents make phytoplankton become the valuable feed items in the aquatic food chain (Natrah et al 2007; Khatoon et al 2009; Banerjee et al 2010).

The phytoplankton distribution in the Gorontalo Bay is influenced by the dynamics occurring within and outside the water. Internal factors such as ecological interactions that take place in the bay waters and external factors such as seasons and human activities around the bay have the potential to affect the stability of the aquatic ecosystem. The purpose of this study is to determine the spatial and seasonal distribution pattern of phytoplankton in the Gorontalo Bay, Indonesia.

Material and Method

Description of the study sites. The study area was localized throughout the Gorontalo City coastal region called as Gorontalo Bay. This research was carried out in west monsoon (in May), transition (in June) and east monsoon season (in July) in the year 2017. Samplings were performed in 5 stations, all displaying diverse types of land use. In the main land near from site 1 there are rare resident settlements and a steep cliff close to site 2 with no human activities. Moreover, land use near from the site 3 is Bone River estuary. In the site 4 there is an oil port and in the site 5 there are densely populated settlements. Sampling in each station was performed out in 3 sub-stations by dragging the line-transect toward the offshore as presented on the map on the Figure1.




Figure 1. Map of the study area showing the locations of stations.

Physical and chemical parameters establishment. Physical parameters such as temperature, salinity, pH, and dissolved oxygen were conducted *in situ* in each station, while chemical parameters performed *ex situ* and determined using spectrophotometry according to the APHA (APHA 2012) methods consist of chlorophyll-*a*, nitrate, ammonium and orthophosphate.

The water sample for phytoplankton analysis purpose was collected using plankton-net (25 µm in mesh size) by horizontally towing at 15 points then compiled into 5 sample bottles preserved with a 10% lugol solution. In the meantime, water samples for chemical parameters measuring purpose were taken in depth ranging from 0 to 2 m from the surface in every site.

Phytoplankton performance. Phytoplankton was identified taxonomically using light microscope referred to Davis (1955) taxonomical guide. Phytoplankton abundances were measured using Lackey drop-transect method. Biological diversity index (H') and dominance index (D) were calculated according to the following equation (Shannon & Wiener and Sampson index 1949 in Odum 1998).

$$H' = - \sum_{i=1}^s \left(\frac{N_i}{N} \right) \ln \left(\frac{N_i}{N} \right)$$

$$D = \sum_{i=1}^s \left(\frac{N_i}{N} \right)^2$$

Where N_i symbol is individual amount of the phytoplankton species, and N is total individual amount, then s is a total phytoplankton species.

Data analysis. The variety of water parameters and counts of phytoplankton species in every sampling site and different season were performed descriptively. Furthermore, the correlation between physico-chemical water parameters and the abundance of phytoplankton in the Gorontalo Bay were analyzed using Microsoft Excel.

Results

Physico-chemical characteristics. Physico-chemical characteristics and chlorophyll-*a* in the Gorontalo Bay were studied parallel to the phytoplankton performance at the same time in the sampled stations. The results are showed in Table 1.

Tomini Bay water surface temperature varied with a minimum of 26.5°C at station 3 in the east monsoon season and a maximum of 29.8°C at station 1 in the west monsoon season. Although water temperature effect on communities and ecosystem processes is debated, it affects phytoplankton growth rates strongly (Edwards et al 2016). Renaud et al (2002) mentioned that phytoplankton tolerate a range of temperature 20-30°C and optimum temperature to phytoplankton development is 27-30°C.

The pH value showed maximum of 6.67 during transition season and a minimum of 5.67 at stations 2, 4 and 5 in in the west monsoon season. Marine phytoplankton in general is resistant to climate change in terms of ocean acidification and does not increase or decrease in its growth rate according to ecological relevant ranges of pH and CO₂ (Berge et al 2010).

The salinity is the main physical parameter that can be attributed to the plankton diversity and acting as a limiting factor, which influences the distribution of plankton communities (Sridhar et al 2006). The salinity ranged between 30.93 and 34.80 ppt. The minimum salinity was observed at station 5 in transition monsoon, while the maximum level was detected at station 1 in west monsoon period.

The dissolved oxygen (DO) concentration in water depends on its temperature and salinity. The quantity of organic matter presenting in the aquatic environment also depends on DO in water. If the decomposition of organic matter is in great proportion then it will absorb of DO in water too much (Shakweer 2003). The DO varied between a maximum of 6.15 mg L⁻¹ in west monsoon period at station 4 and a minimum of 3.59 mg L⁻¹ in east monsoon season at station 3. Phytoplankton are responsible for dissolved oxygen level in surface water, which upturns due to photosynthesis (Ramaraj et al 2010). Concentration of DO naturally associates with microalgae densities in waters (Kunlasak et al 2013).

Table 1

Waters characteristics of Gorontalo Bay in three different seasons in year 2017

<i>Parameters</i>	<i>West monsoon season</i>					<i>Transition season</i>					<i>East monsoon season</i>				
	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>
Temperature	29.80	29.70	28.43	29.07	29.50	29.40	28.17	26.73	28.27	26.83	29.60	28.40	26.50	27.27	28.40
pH	6.00	5.67	6.00	5.67	5.67	6.00	6.00	6.00	6.00	6.67	6.00	6.00	6.00	6.00	6.00
Salinity	34.8	34.13	34.63	35.03	34.03	32.73	32.37	31.53	31.9	30.93	32.10	32.9	33.27	30.87	31.8
DO	6.00	5.40	5.36	6.15	6.14	5.32	4.85	5.38	5.63	5.76	4.45	4.35	3.59	4.41	4.46
NO ₃	0.058	0.060	0.146	0.042	0.084	0.089	0.132	0.079	0.098	0.126	0.061	0.051	0.084	0.117	0.144
NH ₃	0.057	0.069	0.142	0.117	0.122	0.130	0.063	0.206	0.035	0.105	0.051	0.073	0.193	0.054	0.336
PO ₄	0.026	0.057	0.038	0.024	0.044	0.003	0.046	0.013	0.024	0.006	0.059	0.052	0.094	0.137	0.088
Chlorophyll- <i>a</i>	1.789	1.536	1.175	0.651	0.821	1.806	1.610	1.377	1.182	0.984	1.806	1.969	2.332	3.744	1.775

1-5: stations.

The maximum concentration of phosphate (0.137 mg L^{-1}) was shown at station 4 east monsoon season, while the minimum value (0.003 mg L^{-1}) was recorded at station 1 transition period. The role of dissolved inorganic phosphorus could be considered as an important nutrient for marine phytoplankton in the oligotrophic settings and the need for evaluating nutrient limitation at the taxa and/or single cell level, rather than inferring it with nutrient concentrations and ratios or bulk enzyme activities (Mackey et al 2007).

Nitrate is the most stable form of inorganic nitrogen in the oxygenated waters (Nassar et al 2014). Dissolved nitrate attained a maximum of 0.146 in west monsoon season at station 3 and a minimum of 0.042 at station 4. Nitrate and phosphate are primary nutrients for algae growth (Haag 2007). Reduction of nitrogen and phosphorus concentrations also can be a limiting factor to phytoplankton growth (Freeman 2002).

Ammonium is the nitrogenous end product of the bacterial decomposition of natural organic matter containing nitrogen. In the presence of high ammonium concentrations, the phytoplankton productivity could be high or even higher if the cells are using NH_4^+ rather than NO_3^- (Dugdale et al 2007). The maximum concentration of ammonium (0.336 mg L^{-1}) occurred in east monsoon season at station 5 while the minimum (0.035 mg L^{-1}) was found in transition monsoon season at station 4. Hutchinson (1967) stated that Chlorophyta and diatom preferred nitrogen form of nitrate and ammonia.

Ocean phytoplankton production plays a considerable role in the ecosystem. Primary production can be indexed by chlorophyll concentration (Gui & Subrahmanyam 2018). Chlorophyll-*a* is the most commonly used indicator of phytoplankton biomass in the marine environment. It is relatively simple and cost effective to measure when compared to phytoplankton abundance and is thus routinely included in many surveys (Davies et al 2018). The highest chlorophyll-*a* occurred in east monsoon season with the number of average 3.744 mg L^{-1} . Season is one of some factors that can affect the variability of Chlorophyll-*a* concentration in the water. The upwelling in Gorontalo Bay generally occurs in the east season. The mass of water in the upper layers of the waters is driven by the eastern winds then the vacuum of this coated water is filled by the water mass from the nutrient-rich bottom. In addition, Al-Azri et al (2010) stated that the highest concentrations of chlorophyll-*a* (3 mg m^{-3}) were recorded during the southwest monsoon when upwelling is active along the coast of Oman.

Spatial distribution and composition of phytoplankton. The phytoplankton of Gorontalo Bay was represented by 39 taxa from 7 classes and 3 divisions: Chrysophyta (30 taxa), Chlorophyta (7 taxa) and Cyanophyta (2 taxa). Chrysophyta was the most dominant group, since it constitutes about 63.3% of the total average phytoplankton abundance of $182.326 \text{ cell L}^{-1}$, mainly due to the flourishing of *Chaetoceros affinis* (average of $25.441 \text{ cell L}^{-1}$), *Bacteriastrum hyalinum* (average of $19.624 \text{ cell L}^{-1}$), *Chaetoceros brightwelli* (average of $17.539 \text{ cell L}^{-1}$), *Chaetoceros messanensis* (average of $17.007 \text{ cell L}^{-1}$) and *Bacteriastrum delicatulum* (average of $15.660 \text{ cell L}^{-1}$). Cyanophyta was the second dominant group forming about 22.7% of the total average phytoplankton with $65.334 \text{ cell L}^{-1}$ and a relative high occurrence of *Oscillatoria limosa* (average of $42.140 \text{ cell L}^{-1}$) and *Anabaena* sp. (average of $23.194 \text{ cell L}^{-1}$). Whereas, the other algae division, Chlorophyta as the least dominant group forming collectivity about 14.1% (total average $40.558 \text{ cell L}^{-1}$) with relative high occurrence of *Pediastrum simplex* (average of $17.381 \text{ cell L}^{-1}$) and *Pediastrum duplex* (average of $11.708 \text{ cell L}^{-1}$).

Recording of 39 phytoplankton species in this study is considered to be high since compared to the previous study that was reported by Kadim & Arsad (2016) that in the west of Tomini Bay founded 29 species of phytoplankton which dominated by Chrysophyta. In addition, Setyadji & Priatna (2011) stated that in the eastern part of Tomini Bay there were 33 species phytoplankton was recorded and dominated by Bacillariophyceae.

This research shows that the highest abundance of phytoplankton found in the station 3 with an average of 9891 cell L^{-1} followed by station 4 with an average of 8770 cell L^{-1} , station 1 with an average of 7493 cell L^{-1} and station 2 with an average of 6343 cell L^{-1} . Furthermore, the lowest population density of phytoplankton was recorded at

station 5 with the average of 2651 cell L⁻¹. This may be due to rising up of organic materials which entered the river. The sewage discharge can cause phytoplankton productivity growth where sewage was the main source of nitrogen and phosphorus (Burford et al 2012). The distribution and abundance of phytoplankton in tropical waters varied substantially because of the environmental fluctuations during the seasons (Rajkumar et al 2009).

Species diversity index (H') of phytoplankton population during this research varied from 0.433 to 2.712 and the dominance index (D) ranged between 0.071 and 0.302 (Table 2). In the west monsoon period, H' values with an average of 1.287-2.262 were recorded. Meanwhile, in the transition period, H' increased with the average from 2.268 to 2.712. In general, based on Shannon-Wiener diversity index, these two seasons performed an average diversity condition. On the other hand, the lower diversity value recorded during the east monsoon season with an average of 0.433 to 0.904. Diversity depends on key ecological processes such as competition, predation, succession and therefore changes in these processes can alter the species diversity index through changes in evenness (Stirling & Wilsey 2001). The species diversity index of plankton communities can serve as an indicator that the ecosystem is under the influence of pollution or eutrophication (Telesh 2004 in Nassar & Gharib 2014). An increasing in diversity values means the water quality is recovered. Lower species diversity indicated influence of pollution (Nassar & Gharib 2014). According to the dominance index value (D) in the west monsoon to east monsoon seasons, there is no species domination in Gorontalo Bay.

Table 2

Diversity index (H') and dominance index (D) of phytoplankton in the Gorontalo Bay

Seasons	Index	Stations				
		1	2	3	4	5
West monsoon	H'	1.890	2.034	1.287	2.262	1.813
	D	0.177	0.171	0.302*	0.121	0.180
Transition	H'	2.304	2.469	2.268	2.712*	2.284
	D	0.190	0.131	0.129	0.078	0.142
East monsoon	H'	0.723	0.904	0.433	0.874	0.760
	D	0.103	0.071	0.096	0.110	0.118

Note: * maximum values.

Seasonal fluctuation of phytoplankton. Result of phytoplankton analysis according to their abundance, spatial-seasonal fluctuation is successively presented on Table 3 and Figures 2 and 3.

The total phytoplankton in the whole investigated areas showed the highest counts during east monsoon season with an average of 673026 cell L⁻¹ followed by transition season with an average of 352704 cell L⁻¹ and west monsoon season with an average of 3839 cell L⁻¹. It can be seen from Table 3 that in the Gorontalo Bay the east monsoon season was the most productive season. Awwaludin et al (2005) stated that in the east monsoon season, phytoplankton distribution was concentrated in the north part of Tomini Bay with the abundance ranged between 80010 and 1082520 cell m⁻³ which consisted of 45 species and dominated by Bacillariophyceae. In the transitional season, the number of phytoplankton cell tends to be homogeneity recorded as 26 taxa with the abundance ranged from 25000 to 1500000 cell m⁻³ dominated by *Chaetoceros* and *Rhizosolenia* (Wiadnyana 1998).

In this study, the high abundance of phytoplankton during east monsoon season was associated with high flourishing of *Anabaena* sp. (61863 cell L⁻¹ in average), *Oscillatoria limosa* (59902 cell L⁻¹ in average), *Chaetoceros affinis* (53724 cell L⁻¹ in average), *Chaetoceros messanensis* (44783 cell L⁻¹ in average) and *Bacteriastrium hyalinum* (42086 cell L⁻¹ in average). On the other hand, some phytoplankton species showed their maximum abundance during transition season. These taxa are namely

Oscillatoria limosa (average of 66366 cell L⁻¹), *Chlorella* sp. (average of 7817 cell L⁻¹) and *Rhizosolenia imbricata* (average of 7732 cell L⁻¹).

Table 3

Seasonal fluctuation of phytoplankton abundance (cell L⁻¹) in the Gorontalo Bay

Class	Species	West monsoon	Transition	East monsoon
Division: Chlorophyta				
Trebouxiophyceae	<i>Chlorella</i> sp.	141	7817	5761
Zygnematophyceae	<i>Cosmarium</i> sp.	118	2042	2533
	<i>Hyalotheca mucosa</i>	76	1880	5533
	<i>Spirogyra</i> sp.	223	1196	4258
Chlorophyceae	<i>Pediastrum duplex</i>	282	8202	26641
	<i>Pediastrum simplex</i>	0	12858	39283
	<i>Golenkinia</i> sp.	94	2734	0
Division: Chrysophyta				
Bacillariophyceae	<i>Achnanthes brevipes</i>	94	4955	3146
	<i>Dyitylum criophilum</i>	203	1126	4331
	<i>Synedra gailonii</i>	218	4836	11956
	<i>Synedra montana</i>	249	6519	6832
	<i>Synedra ulna</i>	102	6092	7257
	<i>Pleurosigma formosum</i>	24	0	5598
	<i>Amphora obtusa</i>	235	3913	8581
	<i>Cymbella cistula</i>	24	1196	2942
	<i>Gyrosigma</i> sp.	12	352	1136
	<i>Nitzschia</i> sp.	326	5801	6178
	<i>Rhizosolenia castracanei</i>	317	5784	8711
	<i>Rhizosolenia imbricata</i>	642	9665	7020
	<i>Rhizosolenia pungens</i>	509	3238	9766
	<i>Rhizosolenia shrubsolei</i>	353	2136	5720
	<i>Surirella</i> sp.	142	2691	9725
	<i>Corethron hystrix</i>	0	162	597
	<i>Thalassionema nitzschioides</i>	0	8031	11147
	<i>Thalassiothrix frauenfeldii</i>	0	5895	14056
	<i>Trachyneis aspera</i>	352	0	2243
Mediophyceae	<i>Biddulphia aurita</i>	141	85	8989
	<i>Hemiaulus membranaceus</i>	355	3845	11858
	<i>Lauderia anulata</i>	106	5468	5042
	<i>Bacteriastrum delicatulum</i>	176	13456	36039
	<i>Bacteriastrum hyalinum</i>	212	16575	42086
	<i>Chaetoceros affinis</i>	1752	20846	53724
	<i>Chaetoceros brightwelli</i>	690	12986	38940
	<i>Chaetoceros messanensis</i>	0	6237	44783
Coscinodiscophyceae	<i>Coscinodiscus eccentricus</i>	259	3904	4944
	<i>Coscinodiscus radiatus</i>	74	1837	3906
	<i>Coscinodiscus</i> sp.	86	940	2860
Division: Cyanophyta				
Cyanophyceae	<i>Anabaena</i> sp.	142	7577	61863
	<i>Oscillatoria limosa</i>	153	66366	59902

Chaetoceros as a group from Bacillariophyceae appeared frequently in the waters. It has high tolerance in waters salinity fluctuation (Kennish 1990), has segmented cell form where each segment is self-contained, and more adaptable to its environment and is a phytoplankton group favored by fish and shrimp larvae. In addition, *Anabaena* and *Oscillatoria* from Cyanophyceae class also appeared in the waters with a high frequency. They can survive in unfavorable environmental conditions since they have ability to fixing nitrogen in the air directly (Meeks & Elhai 2002; Conradie et al 2008).

The spatial distribution of abundance of phytoplankton species varied substantially among all seasons (Figure 2). The fluctuation of total phytoplankton abundance among sub stations located in front of estuary has alteration with a cyclic-like pattern. In west monsoon season, the higher abundance of total phytoplankton species was separate between in front of estuary and western waters. It was move and being fully concentrated at western waters during transition season. The fully concentrated in front of estuary is back later in east monsoon. The continually lower abundances were located in eastern waters of Gulf Gorontalo (predominantly at sub stations XIII, XIV, and XV).




Figure 2. Tempo-spatial variations of phytoplankton distribution and species-class fluctuation during different season: (a) west monsoon, (b) transition and (c) east monsoon. The interpolation map was constructed by GIS software using the kriging method.

In general, the average of species and class occurrence was rise from time to time of season (Figure 3). Further analysis to comparison in term of ratio between occurrence of average amount of classes and their member of species has also a rising pattern. While in east monsoon the higher ratio was significant, the lag distance (i.e. differences value between ratios) from west monsoon to transition season was higher than transition to east monsoon season.




Figure 3. Average of total abundance of phytoplankton species and classes and their ratio of comparison during three phase of season.

Bacillariophyceae and Mediophyceae from Chrysophyta division have a high species presence variety in Gorontalo Bay. Nybakken (1993) stated that phytoplankton species dominated by Chrysophyta mainly from Bacillariophyceae. The marine phytoplankton community is characterized by high density dominated by diatoms and high species biodiversity (Nursuhayati et al 2013). Diatom dominance among phytoplankton groups has been reported by Shah et al (2008), Ananthan et al (2008), Muylaert et al (2009), Setyadji & Priatna (2011), and Nassar et al (2014).

Phytoplankton growth rate and distribution are tremendously influenced by environmental factors. The affecting parameters differ for each species (Adenan et al 2013). The variety of planktonic diatom species existing in temperate waters is essential. Horizontal distribution influence factors are temperature, salinity, and current. *Rhizosolenia* as the great of number species found dispersed along tropical seawater (circumtropical) and has being adapted with warm temperature (Morrissey & Sumich 2012; Cokrowati et al 2014). Water temperature is very vital for metabolic processes, growth and reproductive ability among aquatic organisms. Power plants can affect temperature fluctuations in the water and can influence the composition of phytoplankton communities. The effect is primarily for phytoplankton existence such as declining of community composition, abundance, and their productivity in water column (Chuang et al 2009; Choi et al 2012).

In addition, species from the Chlorophyta and Cyanophyta divisions in the waters have relatively low species diversity, although *Oscillatoria limosa* exhibits a fairly high abundance in the three distinct seasons. *Oscillatoria* dominance will be in line with the high value of BOD. The dominance of the *Oscillatoria* can be an indication of the decline of water quality or pollution by organic waste disposal (Kumari et al 2008; Rangpan 2008; Panich-pat et al 2009).

Phytoplankton abundance and phisyo-chemical parameters correlation. The relationship between the abundance of phytoplankton and environmental parameters is shown in Table 4. The parameters which have positive correlation with phytoplankton abundance are pH ($r = 0.15$; $p > 0.05$), NO_3 ($r = 0.06$; $p > 0.05$), NH_3 ($r = 0.04$; $p > 0.05$), PO_4 ($r = 0.65$; $p < 0.05$), and chlorophyll-*a* ($r = 0.83$; $p < 0.05$), while it was inversely correlated with temperature ($r = 0.51$; $p > 0.05$), salinity ($r = 0.47$; $p > 0.05$), and dissolved oxygen ($r = 0.83$; $p < 0.05$).

Table 4

Correlation matrix between the physico-chemical parameters and the total counts of phytoplankton in the Gorontalo Bay

	<i>Phyto.</i>	<i>Temp.</i>	<i>pH</i>	<i>Salinity</i>	<i>DO</i>	<i>NO₃</i>	<i>NH₃</i>	<i>PO₄</i>	<i>Chl-a</i>
Phyto.	1.00								
Temp.	-0.78	1.00							
pH	0.54	-0.95	1.00						
Salinity	-0.77	1.00	-0.96	1.00					
DO	-0.98	0.64	-0.37	0.63	1.00				
NO ₃	0.50	-0.93	1.00	-0.94	-0.33	1.00			
NH ₃	0.96	-0.56	0.28	-0.55	-0.99	0.23	1.00		
PO ₄	0.74	-0.15	-0.17	-0.13	-0.85	-0.21	0.90	1.00	
Chl- <i>a</i>	0.96	-0.57	0.28	-0.55	-1.00	0.24	1.00	0.90	1.00

Conclusions. The most productive season in Gorontalo Bay, Indonesia was shown in east monsoon season with the abundance of phytoplankton in average of 673026 cell L⁻¹, followed by the transition and west monsoon season with the mean counts of phytoplankton of 352704 cell L⁻¹ and 3839 cell L⁻¹ consecutively. The highest variety of phytoplankton species found in waters is from the Chrysophyta group. Spatially, the highest abundance of phytoplankton founded in the Station 3 with an average of 9891 cell L⁻¹ followed by Station 4 with an average of 8770 cell L⁻¹, Station 1 with an average of 7493 cell L⁻¹, Station 2 with an average of 6343 cell L⁻¹, and Station 5 with the average of 2651 cell L⁻¹.

Acknowledgements. We would like to thank Directorate of Research and Development, Ministry of Research, Technology, and Higher Education of Republic Indonesia for funding support of this research. We also thank the staff of Fisheries and Marine Science Faculty Laboratory of Gorontalo State University for technical help.

References

- Adenan N. S., Yusoff F. M., Shariff M., 2013 Effect of salinity and temperature on the growth of diatoms and green algae. *Journal of Fisheries and Aquatic Science* 8(2): 397-404.
- Al-Azri A. R., Piontkovski S. A., Al-Hashmi K. A., Goes J. I., do Gomes H. R., 2010 Chlorophyll *a* as a measure of seasonal coupling between phytoplankton and the monsoon periods in the Gulf of Oman. *Aquatic Ecology* 44(2):449-461.
- Ananthan G., Sampathkumar P., Soundarapandian P., Kannan L., 2008 Phytoplankton composition and community structure of Ariyankuppam estuary and Verampattinam coast of Pondicherry. *Journal of Fisheries and Aquatic Science* 3(1):12-21.
- APHA, 2012 Standard methods for the examination of water and wastewater. 22nd edition, Rice E. W., Baird R. B., Eaton A. D., Clesceri L. S. (eds), American Public Health Association (APHA), American Water Works Association (AWWA) and Water Environment Federation (WEF), Washington, D.C., USA, 277 pp.
- Awwaludin A., Suwarso S., Setiawan R., 2005 [The distribution of abundance and structure of plankton communities in the eastern seasons in the Tomini Bay]. *Jurnal Penelitian Perikanan Indonesia* 11(6):33-56. [in Indonesian]
- Banerjee S., Khatoon H., Shariff M., Yusoff F. M., 2010 Enhancement of *Penaeus monodon* shrimp postlarvae growth and survival without water exchange using marine *Bacillus pumilus* and periphytic microalgae. *Fisheries Science* 76(3):481-487.
- Berge T., Daugbjerg N., Andersen B. B., Hansen P. J., 2010 Effect of lowered pH on marine phytoplankton growth rates. *Marine Ecology Progress Series* 416:79-91.
- Burford M. A., Revill A. T., Smith J., Clementson L., 2012 Effect of sewage nutrients on algal production, biomass and pigments in tropical tidal creeks. *Marine Pollution Bulletin* 64(12):2671-2680.

- Choi K. H., Kim Y. O., Lee J. B., Wang S. Y., Lee M. W., Lee P. G., Ahn D. S., Hong J. S., Soh H. Y., 2012 Thermal impacts of a coal power plant on the plankton in open coastal water environment. *Journal of Marine Science and Technology* 20(2):187-194.
- Chuang Y., Yang H., Lin H., 2009 Effects of a thermal discharge from a nuclear power plant on phytoplankton and periphyton in subtropical waters. *Journal of Sea Research* 61:197-205.
- Cokrowati N., Amir S., Abidin Z., Setyono B. D. H., Damayanti A. A., 2014 [Abundance and composition of phytoplankton in Kodek Bay Pemenang Lombok Utara]. *Depik* 3(1): 21-26. [in Indonesian]
- Conradie K. R., Du Plessis S., Venter A., 2008 Re-identification of "*Oscillatoria simplicissima*" isolated from the Vaal River, South Africa, as *Planktothrix pseudagardhii*. *South African Journal of Botany* 74:101-110.
- Davies C. H., Ajani P., Armbrecht L., Atkins N., Baird M. E., et al., 2018 A database of chlorophyll a in Australian waters. *Scientific Data* 5:180018. doi: 10.1038/sdata.2018.18.
- Davis C. C., 1955 *The marine and freshwater plankton*. Michigan State University Press, USA, 562 pp.
- Dugdale R. C., Wilkerson F. P., Hogue V. E., Marchi A., 2007 The role of ammonium and nitrate in spring bloom development in San Francisco Bay. *Estuarine, Coastal and Shelf Science* 73:17-29.
- Edwards K. F., Thomas M. K., Klausmeier C. A., Litchman E., 2016 Phytoplankton growth and the interaction of light and temperature: a synthesis at the species and community level. *Limnology and Oceanography* 6:1232-1244.
- Freeman S., 2002 *Biological science*. Upper Saddle River, Prentice Hall, 1017 pp.
- Gui F., Subrahmanyam M. V., 2018 Chlorophyll variations over coastal area of China due to typhoon Rananim. *Indian Journal of Geo-Marine Sciences* 47:804-811.
- Haag A. L., 2007 Algae bloom again. *Nature* 447:520-521.
- Hutchinson G. E., 1967 *A treatise on limnology*. Volume II. Introduction to lake biology and the limnoplankton. John Wiley and Sons, New York, 1115 pp.
- Kadim M. K., Arsad S., 2016 Distribution and abundance of microalgae based on coastal characteristic and ecology in Bone Bolango coastal region, Indonesia. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* 18(2):395-401.
- Karydis M., Tsiertsis G., 1996 Ecological indices: a biometric approach for assessing eutrophication levels in the marine environment. *Science of the Total Environment* 186(3):209-219.
- Kennish M. J., 1990 *Ecology of estuaries*. Volume II: biological aspects. CRC Press, Inc., Boca Raton, Florida, 391 pp.
- Khatoon H., Banerjee S., Yusoff F. M., Shariff M., 2009 Evaluation of indigenous marine periphytic *Amphora*, *Navicula* and *Cymbella* grown on substrate as feed supplement in *Penaeus monodon* postlarval hatchery system. *Aquaculture Nutrition* 15(2):186-193.
- Kumari P., Dhadse S., Chaudhari P. R., Wate S. R., 2008 A biomonitoring of plankton to assess quality of water in the lakes of Nagpur city. In: *Proceedings of Taal: the 12th World Lake Conference*, pp. 160-164.
- Kunlasak K., Chitmanat C., Whangchai N., Promya J., Lebel L., 2013 Relationship of dissolved oxygen with chlorophyll-a and phytoplankton composition in tilapia ponds. *International Journal of Geosciences* 4:46-53.
- Mackey K. R. M., Labiosa R. G., Calhoun M., Street J. H., Post A. F., Paytan A., 2007 Phosphorus availability, phytoplankton community dynamics, and taxon-specific phosphorus status in the Gulf of Aqaba, Red Sea. *Limnology and Oceanography* 52(2):873-885.
- Meeks J. C., Elhai J., 2002 Regulation of cellular differentiation in filamentous cyanobacteria in free-living and plant-associated symbiotic growth states. *Microbiology and Molecular Biology Reviews* 66(1):94-121.
- Morrisey J. F., Sumich J. L., 2012 *Introduction to the biology of marine life*. 10th edition, Courier Kendallville, USA, 463 pp.

- Muyllaert K., Sabbe K., Vyverman W., 2009 Changes in phytoplankton diversity and community composition along the salinity gradient of the Schelde estuary (Belgium/The Netherlands). *Estuarine, Coastal and Shelf Science* 82(2):335-340.
- Nassar M. Z. A., Gharib S. M., 2014 Spatial and temporal patterns of phytoplankton composition in Burullus Lagoon, Southern Mediterranean Coast, Egypt. *The Egyptian Journal of Aquatic Research* 40(2):133-142.
- Nassar M. Z., Mohamed H. R., Khiray H. M., Rashedy S. H., 2014 Seasonal fluctuations of phytoplankton community and physico-chemical parameters of the north western part of the Red Sea, Egypt. *The Egyptian Journal of Aquatic Research* 40(4):395-403.
- Natrah F. M. I., Yusoff F. M., Shariff M., Abas, F., Mariana N. S., 2007 Screening of Malaysian indigenous microalgae for antioxidant properties and nutritional value. *Journal of Applied Phycology* 19(6):711-718.
- Nursuhayati A. S., Yusoff F. M., Shariff M., 2013 Spatial and temporal distribution of phytoplankton in Perak estuary, Malaysia, during monsoon season. *Journal of Fisheries and Aquatic Science* 8(4):480-493.
- Nybakken J. W., 1993 *Marine biology: an ecological approach*. Harper Collins Publishers, ISBN: 0065008227, 462 pp.
- Odum E. P., 1998 [Fundamentals of ecology]. 3rd edition, Samingan T. (ed), Universitas Gadjah Mada Press, Yogyakarta, 697 pp. [in Indonesian]
- Panich-pat T., Yenwaree W., Ongmali R., 2009 Monitoring of water quality using phytoplankton, protozoa, and benthos as bioindicator in Chadeebucha Canal, Nakhon Pathom Province. *Applied Environmental Research* 31(2):1-14.
- Radiarta I. N., 2013 [The relationship between distribution of phytoplankton with water quality in Alas Strait, Sumbawa Regency, West Nusa Tenggara]. *Jurnal Bumi Lestari* 13(2):234-243. [in Indonesian]
- Rajkumar M., Perumal P., Prabu V. A., Perumal N. V., Rajasekar K. T., 2009 Phytoplankton diversity in Pichavaram mangrove waters from south-east coast of India. *Journal of Environmental Biology* 30(4):489-498.
- Ramaraj R., Tsai D. D. W., Chen P. H., 2010 Algae growth in natural resources. *Journal of Soil and Water Conservation* 42(4):439-450.
- Rangpan V., 2008 Effects of water quality on periphyton in the Pattani River, Yala Municipality, Thailand. Doctoral dissertation, Universiti Sains Malaysia, 290 pp.
- Renaud S. M., Thinh L. V., Lambrinidis G., Parry D., 2002 Effect of temperature on growth, chemical composition and fatty acid composition of tropical Australian microalgae grown in batch cultures. *Aquaculture* 211(1-4):195-214.
- Setyadji B., Priatna A., 2011 [Spatial and temporal distribution of plankton in Tomini Bay, Sulawesi]. *BAWAL* 3(6):387-395. [in Indonesian]
- Shah M. M. R., Hossain M. Y., Begum M., Ahmed Z. F., Ohtomi J., Rahman M. M., Alam M. J., Islam M. A., Fulanda B., 2008 Seasonal variations of phytoplanktonic community structure and production in relation to environmental factors of the southwest coastal waters of Bangladesh. *Journal of Fisheries and Aquatic Science* 3: 102-113.
- Shakweer L., 2003 Ecological and fishery investigations of Nozha Hydrome near Alexandria 2000-2001. 1. Chemistry of Nozha Hydrome water under the conditions of fertilizers applications. *Bull Natl Inst Oceanogr Fish* 29:387-425.
- Sridhar R., Thangaradjou T., Kumar S. S., Kannan L., 2006 Water quality and phytoplankton characteristics in the Palk Bay, southeast coast of India. *Journal of Environmental Biology* 27(3):561-566.
- Stirling G., Wilsey B., 2001 Empirical relationships between species richness, evenness, and proportional diversity. *The American Naturalist* 158(3):286-299.
- Thoha H., Amri K., 2011 [Composition and abundance of phytoplankton in south Kalimantan waters]. *Oceanologi dan Limnologi di Indonesia* 37(2):371-382. [in Indonesian]
- Townsend C. R., Harper J. L., Begon M., 2000 *Essentials of ecology*. Blackwell Science, Oxford, UK, 553 pp.

- Wiadnyana N. N., 1998 [Distribution and variation of phytoplankton pigments in Tomini Bay, North Sulawesi]. Seminar Kelautan Ambon: LIPI-UNHAS, pp. 248-259. [in Indonesian]
- Yusron E., Edward., 2000 [Waters condition and biodiversity in the Tomini Bay waters, North Sulawesi]. Seminar Nasional Pendayagunaan Sumberdaya Hayati Dalam Pengelolaan Lingkungan, Fakultas Biologi, Universitas Kristen Satya Wacana, Salatiga. [in Indonesian]

Received: 27 April 2018. Accepted: 31 May 2018. Published online: 21 June 2018.

Authors:

Miftahul K. Kadim, Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Jl. Jendral Sudirman. No. 6, Gorontalo City, Gorontalo Province, Indonesia, e-mail: miftahulhairkadim@ung.ac.id

Nuralim Pasingi, Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Jl. Jendral Sudirman. No. 6, Gorontalo City, Gorontalo Province, Indonesia, e-mail: pasisinginuralim@gmail.com

Faizal Kasim, Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Gorontalo State University, Jl. Jendral Sudirman. No. 6, Gorontalo City, Gorontalo Province, Indonesia, e-mail: faizalkasim@ung.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Kadim M. K., Pasingi N., Kasim F., 2018 Spatial and temporal distribution of phytoplankton in the Gorontalo Bay, Indonesia. *AAFL Bioflux* 11(3): 833-845.