


ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 20 - Number 1 - January 2019





Biodiversitas

11

H Index

CountryIndonesia -  [SJR Ranking of Indonesia](#)**Subject Area and Category**[Agricultural and Biological Sciences](#)
[Animal Science and Zoology](#)
[Plant Science](#)[Biochemistry, Genetics and Molecular Biology](#)
[Molecular Biology](#)**Publisher**[Biology department, Sebelas Maret University Surakarta](#)**Publication type**[Journals](#)**ISSN**[1412033X, 20854722](#)**Coverage**[2014-2020](#)**Scope**

"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biological aspects of plants, animals and microbes at the level of gene, species, and ecosystem.

[Homepage](#)[How to publish in this journal](#)[Contact](#)[Join the conversation about this journal](#)**Quartiles**

Animal Science and Zoology

Molecular Biology

Plant Science

2015

2016

2017

2018








2019

SJR



Citations per document





Biodiversitas

Q3

Animal Science and Zoology

best quartile

SJR 2019

0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="265 728 408 740">
```

K **KETUT SUADA** 2 months ago

Dear Editors

May I know the reason/s of Why my article title "The potential of various indigenous *Trichoderma* spp. to suppress *Plasmodiophora brassicae* the pathogen of clubroot disease on cabbage" DOI: 10.13057/biodiv/d180418, in BIODIVERSITAS VOL 18/4 OCT 2017, PAGES:1424-1429, was justified as "SHORT COMMUNICATION", WHILE THE DATA IN THE ARTICLE WAS COMPLETE INCLUDING TO DIVERSITY AND EVEN ITS EFFECT TO THE TRICHODERMA IN PLANT (CABBAGE), CAN YOU

BIODIVERSITAS

Journal of Biological Diversity
Volume 20 – Number 1 – January 2019

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD (COMMUNICATING EDITORS):

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Alan J. Lymbery** (Australia), **Alireza Ghanadi** (Iran), **Ankur Patwardhan** (India), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Faiza Abbasi** (India), **Hassan Pourbabaie** (Iran), **I Made Sudiana** (Indonesia), **Ivan Zambrana-Flores** (United Kingdom), **Joko R. Witono** (Indonesia), **Katsuhiko Kondo** (Japan), **Krishna Raj** (India), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Maresh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Muhammad Akram** (Pakistan), **Mochamad A. Soendjoto** (Indonesia), **Mohib Shah** (Pakistan), **Mohamed M.M. Najim** (Srilanka), **Morteza Eighani** (Iran), **Pawan K. Bharti** (India), **Paul K. Mbugua** (Kenya), **Rasool B. Tareen** (Pakistan), **Seweta Srivastava** (India), **Sayed Aliakbar Hedayati** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Stavros Lalas** (Greece), **Subhash Santra** (India), **Sugiyarto** (Indonesia), **T.N. Prakash Kammardi** (India)

EDITOR-IN-CHIEF:

Sutarno

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, Email: unsjournals@yahoo.com

ONLINE:

biodiversitas.mipa.uns.ac.id

.....

EXPERTISE AND CORRESPONDING EMAIL OF THE COMMUNICATING EDITORS:

GENETIC DIVERSITY: **Agnieszka B. Najda** (agnieszka.najda@up.lublin.pl), **Alan J. Lymbery** (a.lymbery@murdoch.edu.au), **Darlina Md. Naim** (darlinamdn@usm.my), **Mahendra K. Rai** (pmkrai@hotmail.com).

SPECIES DIVERSITY: **Joko R. Witono** (jrwitono@yahoo.com), **Katsuhiko Kondo** (k3kondo@nodai.ac.jp), **Livia Wanntorp** (livia.wanntorp@nrm.se), **Maresh K. Adhikari** (mkg_adh@wlink.com.np), **Maria Panitsa** (mpanitsa@upatras.gr), **Mohib Shah** (mohibshah@awkm.edu.pk), **Paul K. Mbugua** (paulkmbugua@gmail.com), **Rasool B. Tareen** (rbtareen@yahoo.com).

ECOSYSTEM DIVERSITY: **Abdel Fattah N.A. Rabou** (arabou@iugaza.edu), **Alireza Ghanadi** (aghannadi@yahoo.com), **Ankur Patwardhan** (ankurpatwardhan@gmail.com), **Bambang H. Saharjo** (bhsaharjo@gmail.com), **Daiane H. Nunes** (nunesdaiane@gmail.com), **Faiza Abbasi** (faeza.abbasi@gmail.com), **Ghulam Hassan Dar** (profdar99@gmail.com), **Hassan Pourbabaie** (hassan_pourbabaie@yahoo.com), **I Made Sudiana** (sudianai@yahoo.com), **Ivan Zambrana-Flores** (izambrana@gmail.com), **Krishna Raj** (krishnarajise@yahoo.co.uk), **Mahdi Reyahi-Khoram** (phdmrk@gmail.com), **Mochamad A. Soendjoto** (masoendjoto@gmail.com), **Mohamed M.M. Najim** (mnajim@kln.ac.lk), **Morteza Eighani** (morteza_eighani@yahoo.com), **Pawan K. Bharti** (gurupawanbharti@rediffmail.com), **Seweta Srivastava** (seweta.21896@lpu.co.in), **Sayed Aliakbar Hedayati** (Hedayati@gau.ac.ir), **Shahabuddin** (shahabsaleh@gmail.com), **Shahir Shamsir** (shahirshamsir@gmail.com), **Shri Kant Tripathi** (sk_tripathi@rediffmail.com), **Stavros Lalas** (slalas@teilar.gr), **Subhash Santra** (sesantra@yahoo.com), **Sugiyarto** (sugiyarto_ys@yahoo.com), **T.N. Prakash Kammardi** (prakashtnk@yahoo.com).

ETHNOBIOLOGY: **M. Jayakara Bhandary** (mbjaikar@gmail.com), **Muhammad Akram** (makram_0451@hotmail.com).



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers(double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

A charge Starting on January 1, 2017, publishing costs waiver is granted to foreign (non-Indonesian) authors who first publish the manuscript in this journal, especially for graduate students from developing countries. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m⁻², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cir" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaad AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

THIS PAGE INTENTIONALLY LEFT BLANK

After the publication of Biodiversitas vol. 15, no. 2, October 2014, furthermore, all manuscripts will be published online as soon as the revision is approved (no time constraints), but they will be printed in the next issue, namely Biodiversitas vol. 16, no. 1, April 2015, and so on.

Examples of how to cite:

- *Prior to the date of printing (before 1 April 2015):*
Tavankar F, Bonyad AE. 2014. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas DOI: 10.13057/biodiv/d160101. (Year is referring to the download time)
- *After the print date (after 1 April 2015):*
Tavankar F, Bonyad AE. 2015. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas 16 (1): 1-9.

NOTIFICATION

Starting on January 1, 2019, Biodiversitas will be issued monthly. New issues will also be found at [Smujo](#), as an online first publication.



Home | Archives | Contact Us | Affiliate Program
Copyright © Biodiversitas, 2012. All Rights Reserved



All content on this website, unless otherwise noted, is licensed under a [Creative Commons Attribution 4.0 License](#)

[Home](#)[Content and Archives](#)[Aims and Scopes](#)[Editorial Board](#)[Guidance for Authors](#)[Ethical Guidelines](#)[Charges](#)[Membership](#)[Previously Reviewers](#)[Conference Events](#)**Biodiversitas Vol. 20, No. 1, January 2019**

only search Biodiversitas

Table of Content

[Cover \[PDF\]](#)[Editorial Board & Guidance for Authors \[PDF\]](#)

1. The sufficiency of existed protected areas in conserving the habitat of proboscis monkey (*Nasalis larvatus*)
WARDATUTTHOYIBAH, SATYAWAN PUDYATMOKO, SENA ADI SUBRATA, MUHAMMAD ALI IMRON [\[PDF\]](#)
2. The prevalence and species of root-knot nematode which infect on potato seed in Central Java, Indonesia
MUTALA'LIAH, SIWI INDARTI, ARIF WIBOWO [\[PDF\]](#)
3. Genetic diversity and phylogenetic analysis of two Indonesian local cattle breeds based on cytochrome b gene sequences
TETY HARTATIK, DWI NUR HAPPY HARIYONO, YUDI ADINATA [\[PDF\]](#)
4. Impact of the green revolution on the gender's role in wet rice farming: A case study in Karangwangi Village, Cianjur District, West Java, Indonesia
RUHYAT PARTASASMITA, BUDIAWATI S. ISKANDAR, SITI NURAENI, JOHAN ISKANDAR [\[PDF\]](#)
5. The study of suweg (*Amorphophallus paeoniifolius*) and other undergrowth species in teak plantation forest of Temengeng, Blora, Indonesia
SYAMSUL HIDAYAT [\[PDF\]](#)
6. Typology of native species as the shade tree for merbau (*Intsia bijuga*) plantations in Papua, Indonesia based on ecological species group
ELIESER VIKTOR SIRAMI, DJOKO MARSONO, RONGGO SADONO, MUHAMMAD ALI IMRON [\[PDF\]](#)
7. Taxonomy confirmation of *Puntius* cf. *binotatus* from Gunung Tujuh Lake, Jambi, Indonesia based on Cytochrome Oxidase-I (COI) gene
DEWI IMELDA ROESMA, DJONG HON TJONG, WILA KARLINA, DYTA RABBANI AIDIL [\[PDF\]](#)
8. Spatio-temporal distribution of microalgae producing chlorophyll and carotenoid pigments in Bali Strait, Indonesia
ANNA FAUZIAH, DIETRICH G BENGEN, MUJIZAT KAWAROE, HEFNI EFFENDI, MAJARIANA KRISANTI [\[PDF\]](#)
9. Expression of oil palm (*Elaeis guineensis*) polyisoprenoids in response to *Ganoderma boninense* infection
DADANG AFANDI, MOHAMMAD BASYUNI, LOLLIE AGUSTINA P. PUTRI, DIANA CHALIL, INDRA SYAHPUTRA [\[PDF\]](#)
10. Conflict between human and wild Sumatran Elephant (*Elephas maximus sumatranus* Temminck, 1847) in Aceh Province, Indonesia
INDIRA NURUL QOMARIAH, TUTIA RAHMI, ZURAIDAH SAID, ARIEF WIJAYA [\[PDF\]](#)
11. Short Communication: Age and growth of elongated mudskipper, *Pseudapocryptes elongatus* (Cuvier, 1816) from Sundarbans, India
GOPALAN MAHADEVAN, PALANIVEL BHARATHIRAJAN, VELAYUTHAM RAVI, MOJTABA POULADI, MARYAM MIRZAEI VAND KHANGHAH [\[PDF\]](#)
12. Bamboo diversity of Sulawesi, Indonesia
DITA ERVIANTI, ELIZABETH A. WIDJAJA, AGUNG SEDAYU [\[PDF\]](#)
13. Characteristics and causes of forest and land fires in Kapuas District, Central Kalimantan Province, Indonesia
ACHMAD SIDDIK THOHA, BAMBANG HERO SAHARJO, RIZALDI BOER, MUHAMMAD ARDIANSYAH [\[PDF\]](#)
14. Local monitoring of flowering and fruiting of Jernang, *Daemonorops* species in Sumatra, Indonesia
LARS SCHMIDT, HIMMAH RUSTIAMI, IDA THEILADE [\[PDF\]](#)
15. Qualitative morphological diversity of female Pelung Chickens in West Java, Indonesia
INDRAWATI Y. ASMARA, DANI GARNIDA, WIWIN TANWIRIAH, RUHYAT PARTASASMITA [\[PDF\]](#)
16. Growth evaluation of native tree species planted on post coal mining reclamation site in East Kalimantan, Indonesia
DEWI AYU LESTARI, ABBAN PUTRI FIQA, FAUZIAH, SUGENG BUDIHARTA [\[PDF\]](#)
17. Folic acid content and fruit characteristics of five Indonesian dessert banana cultivars
RITA NINGSIH, RITA MEGIA [\[PDF\]](#)
18. Pollen morphology of eight tribes of Asteraceae from Universitas Indonesia Campus, Depok, Indonesia
ANDI SALAMAH, RACHMI LUTHFIKASARI, ASTARI DWIRANTI [\[PDF\]](#)
19. The practice of hydrological restoration to rehabilitate abandoned shrimp ponds in Bunaken National Park, North Sulawesi, Indonesia
RIGNOLDA DJAMALUDDIN, BENJAMIN BROWN, ROY R. LEWIS III [\[PDF\]](#)
20. Relationship between environmental parameters and the plankton community of the Batuhideung Fishing

Grounds, Pandeglang, Banten, Indonesia

NOVERITA D. TAKARINA, WAWAN NURLIANSYAH, WISNU WARDHANA [PDF]

21. The natural forest gaps maintenance diversity of understory birds in Mae Sa-Kog Ma Biosphere Reserve, northern Thailand
SUPALAK SIRI, YUWADEE PONPITUK, MONGKOL SAFOOWONG, DOKRAK MAROD, PRATEEP DUENGKAE [PDF]
22. Local wisdom of Talang Mamak Tribe, Riau, Indonesia in supporting sustainable bioresource utilization
PRIMA WAHYU TITISARI, ELFIS, IRINA SAFITRI ZEN, KHAIRANI, NADIATUL JANNA, NUNUT SUHARNI, TIKA PERMATA SARI [PDF]
23. The importance of in-situ conservation area in mining concession in preserving diversity, threatened and potential floras in East Kalimantan, Indonesia
ABBAN PUTRI FIQA, FAUZIAH, DEWI AYU LESTARI, SUGENG BUDIHARTA [PDF]
24. Short Communication: Further morphological evidence for separating *Mukia* Arn. from *Cucumis* L.
MENTARI PUTRI PRATAMI, TATIK CHIKMAWATI, RUGAYAH [PDF]
25. Short Communication: Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia
MAULANA KHALID RIEFANI, MOCHAMAD ARIEF SOENDJOTO, ANDY M. MUNIR [PDF]
26. Distribution and habitat preferences of Sumatran elephant (*Elephas maximus sumatranus*) in Riau, Indonesia
YOHANES WISHNU SUKMANTORO, HADI S. ALIKODRA, AGUS P. KARTONO, EFRANSJAH [PDF]
27. Distribution Body size characteristics and polymorphism in GH and GHRH genes of Simeulue Buffalo of Aceh, Indonesia
KARTINI ERIANI, RAHMI RAHMI, IKHWAN JAMIL, ROSNIZAR ROSNIZAR, AL AZHAR [PDF]
28. Terrestrial gastropod diversity, distribution and abundance in areas with and without anthropogenic disturbances, Northeast Algeria
HADJER BELHIOUANI, MOHAMED EL-HADEF EL-OKKI, FATIMA-ZOHRA AFRI-MEHENNAOUI, LEILA SAHLI [PDF]
29. Short Communication: Dye-yielding plant resources of Maharashtra, India: A checklist
PATIL SONALI HINDURAO, KURLAPKAR DILIP DAMODAR, GAIKWAD DATTATRAYA KRISHNA [PDF]
30. Initiating the establishment of commercial stand at Tasik Besar Serkap, Riau, Indonesia: An early growth assessment of two peat swamp tree species
HENTI HENDALASTUTI RACHMAT, ATOK SUBIAKTO [PDF]
31. The response of East Kalimantan, Indonesia local rice cultivars against iron stress
NURHASANAH, HELMA SUCI LESTARI, WIDI SUNARYO [PDF]
32. Local biodiversity conservation in Sigi, Central Sulawesi, Indonesia: Analysis of the effect of elevation, land accessibility, and farmers income and perception on vegetation diversity in agroforestry systems
NINA DWI LESTARI, DIDI SUPRAYOGO, ARIEF RACHMANSYAH [PDF]
34. Using micro-morphological fruit characters in resolving some of ambiguities in Iranian *Acer* L. (Sapindaceae) species
FATEMEH ASADI, FARIBA SHARIFNIA, FAHIMEH SALIMPOUR, AHMAD MAJD [PDF]
35. Comparative study of floristic diversity along altitude in the northern slope of the central Alborz Mountains, Iran
HALIME MORADI, FARIDEH ATTAR [PDF]
36. Presence of *Anisakis* nematode larvae in Indian mackerel (*Rastrelliger* spp.) along the Indian Ocean southern coast of East Java, Indonesia
EKO SETYOBUDI, IZANATUR ROHMAH, RIZKA FAUZIANA SYARIFAH, LISA RAMATIA, MURWANTOKO, DINI WAHYU KARTIKA SARI [PDF]
37. Response of polyisoprenoid concentration and profile in three groups of mangrove seedlings of coping with long-term salinity
MOHAMMAD BASYUNI, HIROSHI SAGAMI, SHIGEYUKI BABA, HIROSUKE OKU [PDF]
38. Morphology, palynology, and stipe anatomy of four common ferns from Pekanbaru, Riau Province, Indonesia
NERY SOFIYANTI, DYAH IRIANI, FITMAWATI, AFNI ATIKA MARPAUNG [PDF]
39. Short Communication: Early stability test of mutant candidates of Bangka local cassava, Indonesia
TRI LESTARI, ERIES DYAH MUSTIKARINI, RION APRIYADI, SAIPUL ANWAR [PDF]
40. Species diversity of Orchids in Bukit Barisan Selatan National Park, Lampung, Indonesia
Esti Munawaroh, Yuzammi [PDF]
41. Analysis of land use change and its relation to land potential and elephant habitat at Besitang Watershed, North Sumatra, Indonesia
WANDA KUSWANDA, AHMAD DANY SUNANDAR [PDF]
42. Comparison of true mangrove stands in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia
FAIZAL KASIM, MIFTAHUL KHAIR KADIM, SITTI NURSINAR, ZULKIFLI KARIM, ALDIN LAMALANGO [PDF]
43. Physiological status of high and low metabolism *Hevea* clones in the difference stage of tapping panel dryness
RADITE TRISTAMA, PUTRI ARISMA SIREGAR MAWADDAH, LUBIS ADE-FIPRIANI, JUNAIDI [PDF]
44. Short Communication: Exploration of bidara laut (*Strychnos lucida*) parent trees in Gunung Tunak Ecotourism Park, West Nusa Tenggara, Indonesia
ALI SETYAYUDI, KRISNAWATI, RYKE NANDINI [PDF]
45. The diversity of *Smilax* (Smilacaceae) in Besiq-Bermai and Bontang Forests, East Kalimantan, Indonesia
SITI SOFIAH, LULUT DWI SULISTYANINGSIH [PDF]
46. Salinity and subsequent freshwater influences on the growth, biomass, and polyisoprenoids distribution of *Rhizophora apiculata* seedlings
MOHAMMAD BASYUNI, MASRIDA WASILAH, POPPY ANJELISA ZAITUN HASIBUAN, NURDIN SULISTYONO, SUMARDI, YUNTHA BIMANTARA, RAHMAH HAYATI, HIROSHI SAGAMI, HIROSUKE OKU [PDF]
47. Population status of endemic tree kokoleceran (*Vatica bantamensis*) in Ujung Kulon National Park, Indonesia
IYAN ROBIANSYAH, DODO, ARIEF HAMIDI [PDF]

Comparison of true mangrove stands in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia

FAIZAL KASIM^{1,*}, MIFTAHUL KHAIR KADIM¹, SITTI NURSINAR¹, ZULKIFLI KARIM^{1,2},
ALDIN LAMALANGO^{1,2}

¹Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Universitas Negeri Gorontalo, Gorontalo, Indonesia. Jl. Jenderal Sudirman No. 6, Gorontalo City 96128, Gorontalo Province, Indonesia. Tel./fax. +62-435-821125, *email: faizalkasim@ung.ac.id, kasim.faizal@gmail.com

²Gorontalo Coastal Ecological and Spatial Mapping Studies Club (GCEMS). Dulalowo, Kota Tengah, Gorontalo 96138, Indonesia

Manuscript received: 3 November 2018. Revision accepted: 30 December 2018.

Abstract. Kasim F, Kadim MK, Nursinar S, Karim Z, Lamalango A. 2019. Comparison of true mangrove stands in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia. *Biodiversitas* 20: 259-266. This study aimed to investigate and compare the current status of mangrove areas, as well as the composition and species diversity of mangrove stands in both regions of Dudepo and Ponelo Islands. The results showed that the mangrove areas calculated using the segmentation method in classifying image of Landsat-8 OLI (acquisition on September 2017) were 279.46 ha (Dudepo Island) and 113.35 ha (Ponelo Island) respectively. A total of 13 species of true mangrove were recorded from both islands, using survey method from 9 transect lines (TL), with a distance ranging from 40 to 210 meters (1-6 quadrats) per transect. The mean densities of trees were $2133 \pm 329.78 \text{ ha}^{-1}$ (Dudepo Island) and $2111 \pm 234.28 \text{ ha}^{-1}$ (Ponelo Island), while those of saplings and seedlings were $58 \pm 13.48 \text{ ha}^{-1}$ and $1425 \pm 113.96 \text{ ha}^{-1}$ (Dudepo Island), and $79 \pm 14.51 \text{ ha}^{-1}$ and $2963 \pm 443.22 \text{ ha}^{-1}$ (Ponelo Island). The mean diameter and basal area were $19.73 \pm 10.65 \text{ cm}$ and $84.22 \pm 67.67 \text{ m}^2 \text{ ha}^{-1}$ (Dudepo Island), $17.04 \pm 1.46 \text{ cm}$ and $60.07 \pm 15.12 \text{ m}^2 \text{ ha}^{-1}$ (Ponelo Island), respectively. The Importance Value Index (IVI) ranged between 3.97-114.87 (Dudepo Island) and 6.04-82.18 (Ponelo Island). The dominant and codominant species based on IVI in both islands were *Rhizophora apiculata* Blume and *R. stylosa* Griff. The indexes of diversity, richness, and evenness of mangrove species in both islands were 0.34-1.70, 0.48-1.18, 0.47-0.94 (trees), 0.00-1.10, 0.00-1.82, 0.00-1.00 (saplings), and 0.00-1.48, 0.00-1.44, 0.72-1.00 (seedlings), respectively. The Bray-Curtis similarity index between Dudepo and Ponelo Islands, based on the overall values of community attributes, was 0.75.

Keywords: Basal area, Bray-Curtis, Diversity, Gorontalo, mangroves

INTRODUCTION

Until the end of 2000, Indonesia's mangrove forest area was estimated to be 3,112,989 ha or 22.6% of the total mangrove area in the world (Giri et al. 2011). Therefore, mangrove resources are important biological wealth of Indonesia's coastal areas. Mangrove forest becomes a transitional ecosystem in almost all coastal areas in the Indonesian Archipelago, ranging from Sumatra, Java, Kalimantan, Bali, Sulawesi, Maluku, to Papua island (Kusmana et al. 2003; Rugayah, 2007; Marbawa et al. 2014; Prasetyo et al. 2014; Muhtadi et al. 2016; Setiawan et al. 2017; Wouthuyzen and Ahmad, 2018). Besides occupying large area, the mangroves in Indonesia are also known to have high diversity. Snedaker (1984) noted that there were at least 202 species of mangroves in Indonesia, 43 of which (consists of 33 species of trees and several species of shrubs) were true mangrove species, and the rest were other species that lived around mangroves, known as associated mangroves. Mangroves provide a variety of benefits, both goods and services for most communities in coastal areas. The essential ecological and economic functions of mangrove forests are that they provide materials needed by people for commercial, recreational, and also fishery purposes through environmental services

as a spawning ground, nursery habitat for marine fauna (Zhang et al. 2007; Giri et al. 2011; Nfotabong-Atheull et al. 2011; Joshi and Ghose 2014; Hutchison et al. 2014).

The diversity of true mangrove species in northern Gorontalo coastal area was reported to be quite high. Kasim et al. (2017) recorded 19 species in eastern coast area, dominated by members of the Rhizophoraceae family both at genus and species levels. Meanwhile, two other species in this region which are important for global conservation are *Aegiceras floridum* Roem. & Schult (local name *Tongge*) and *Ceriops decandra* (Griff.) Ding Hou (local name *Posi-posi*). Further studies of mangrove forests condition are necessary for resource conservation and coastal land management. For Gorontalo-which is geographically laid in the heart of Wallacea (Gorlinski 2012)-the comprehensive studies of biodiversity in broader areas of mangrove distribution could provide the best choice of management strategies. In term of mangrove complexity study, the significance of the results will be obtained by excluding the species of associated mangrove into the analysis (Blanco et al. 1999). Hence, in this study, we only studied the true mangroves and their distribution. The study aimed to compare the community attributes (species composition, species importance value, and diversity) of true mangroves in Dudepo and Ponelo Islands,

North Gorontalo District, Indonesia.

MATERIALS AND METHODS

Site of research

Dudepo and Ponelo Islands are two large islands in North Gorontalo District, Indonesia. They are separated

administratively: Anggrek Sub-district and Ponelo Kepulauan Sub-district. Both areas are facing the Sulawesi Sea (Figure 1). The sites of research were located in 3 villages in Ponelo Kepulauan Sub-district, and 4 hamlets in Dudepo Island (Table 1).




Figure 1. The research locations in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia. The red dots showed the station positions and the line transects in each location

Table 1. Detailed information on the observed mangroves per station in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia

Island	Station	Coordinate	Code of Transect	No. of quadrats	pH	Salinity (‰)	Temperature (°C)	Substrate
Dudepo	Hamlet of Makassar	122° 46' 33.52" E, 0° 53' 35.75" N	ST1	3	6.5	28	30	Md
	Hamlet of Upo	122° 46' 51.42" E, 0° 53' 34.8" N	ST2	3	7.4	27-28	27.5-27.8	Md
	Hamlet of Tapia	122° 48' 18.56" E, 0° 53' 1.23" N	ST3	3	7.4	30	29.1-29.	Md, Sn
	Hamlet of Bolongo	122° 47' 30.1" E, 0° 53' 38.1" N	ST4	3	7.1	30-32	29.8-30.1	Md, Ms
Ponelo	Village of Atiola	122° 52' 52.46" E, 0° 51' 43.98" N	ST5	6	7.8	31-32	29-29.7	Md, Ms, Dc
		122° 53' 8.8" E, 0° 51' 33.58" N	ST6	3	8	31-33	29.5	Md, Ms, Dc
	Village of Malambe	122° 51' 44.76" E, 0° 53' 32.63" N	ST7	3	7.6	30-31	29.5-30.2	Ms
		122° 52' 1.01" E, 0° 53' 33.41" N	ST8	3	6.81	30-34	30.1-30.3	Sn, Dc
	Village of Ponelo	122° 53' 33.92" E, 0° 52' 44.85" N	ST9	4	6.82	33-35	29.4-30	Ms, Rc

Note: Md = Mud, Ms = Muddy sand, Dc = Dead coral, Rc = rocky, Sn = Sand

Research procedure

Field survey and data collection

The field survey was conducted from May to August 2017. Details of all stations, transects and locations are presented in Table 1. Data collection of mangrove was done using line-transects and 10 x 10 m quadrats made purposively. The interval of quadrats in each transect was 30 to 50 m, depending on the width of mangrove belt on the station. In each quadrat, data of trees, saplings, and seedlings, and also environmental condition were collected (Table 1). The width of mangrove belts was 140 to 657 m in Dudepo Island, and 40-300 m in Ponelo Island. The total areas of sampling in Dudepo and Ponelo Islands were 0.12 ha and 0.19 ha, respectively, resulted from 3 to 6 quadrats employed in perpendicular or parallel transects to shoreline.

Types and sources of data collected

Only true mangrove species were identified and counted in each quadrat. Mangrove species were identified according to the true mangrove description by Kusmana et al. (2003), Noor et al. (2012) and Wetland International (2017). The species name and diameter at breast height (DBH, at 1.3 m) were recorded. The mangrove plants were classified into three categories based on the growth stages as follows: 1) Seedling was any woody plant from sprouts to young mangrove up to 1.5 m high; 2) Sapling was any woody plant higher than 1.5 m with a DBH of less than 10 cm; 3) Tree was any woody plant with a DBH of more than 10 cm (Cintron and Novelli 1984; Cañizares and Seronay 2016; Joshi and Ghose 2014; Winata et al. 2017).

The data of mangrove area and distribution delineated from other land cover were calculated using Landsat-8 (OLI) Path / Row: 113/059, acquisition on September 21, 2017, obtained from U.S. Geological Survey Visualization Viewer or earth explorer which is available free to download through the <http://earthexplorer.com>.

Data analyses

Landsat image analysis for mangrove distribution area

To determine the mangrove distribution area, the scene of image was cropped to the area being observed. Then, it was delineated using the mixed method of supervised classification and segmentation of band composites of RGB 654. SAGA GIS 6.3.0 software was used for all processes of mangrove delineation.

Analysis of mangrove structure

Vegetation data were analyzed using importance value index (IVI) of species (Cintron and Novelli 1984).

The importance value index for each species was calculated using this formula:

$$IVI = \text{Relative Density} + \text{Relative Frequency} + \text{Relative Dominance}$$

$$\text{Density of species } i = \frac{\text{number of trees of species } i}{\text{area of quadrats}}$$

$$\text{Relative density of species } i = \frac{\text{Density of species } i}{\text{Density of all species}} \times 100 \%$$

$$\text{Frequency of species } i = \frac{\text{number of quadrats where species } i \text{ was found}}{\text{total quadrats}}$$

$$\text{Relative frequency of species } i = \frac{\text{Frequency of species } i}{\text{frequency of all species}} \times 100 \%$$

$$\text{Dominance of species } i = \frac{\text{Basal area of species } i}{\text{area of quadrats}}$$

$$\text{Relative dominance of species } i = \frac{\text{Dominance of species } i}{\text{Dominance of all species}} \times 100 \%$$

The diversity, richness and evenness of species in stations and locations were analyzed using the Shannon-Wiener diversity index, H' , Margalef wealth index, and Buzas and Gibson's evenness index. The similarity of the species composition and diversity indexes among all observed quadrats, in all stations and locations was then analyzed using multivariate clustering analysis (Bray-Curtis's index) (Hammer 2017). The procedure followed that described by Kasim et al. (2017) using PAleontological STatistics software (PAST Version 3.15).

RESULTS AND DISCUSSION

Mangrove extent and distribution

The map of the mangrove area and distribution based on the analysis of Landsat-8 OLI images between Dudepo and Ponelo Islands is presented in Figure 2, which shows the differences in distribution and mangrove area between the two islands. The mangrove area in Dudepo Island was larger (279.46 ha) than that in Ponelo Island (113.35 ha). The front area of both islands has relatively same situation of mangrove distribution in the north, directly facing the open area of the Sulawesi Sea. Hence, generally, the mangrove is located in the back area in both islands, facing the mainland (Kwandang and Anggrek), protected from the influence of waves. The mangrove distribution in Dudepo Island in the north is limited to the areas of small bay characterized by the presence of river flow influence and also the protection from the influence of waves.

Although there was the same intensive land cover change in upland area of both islands, we observed that in term of habitat loss and natural life of mangrove-the mangrove ecosystem in Dudepo was more protected from the impacts of human activities than that in Ponelo. Not surprisingly, then, that the mangrove area on Dudepo was larger (2.47 times) than that in Ponelo.

Species composition and mangrove community structure

Table 2 shows the 13 species recorded (members of 6 genera of 4 families) which made up the true mangrove community in Dudepo and Ponelo Islands.

The maximum richest taxa in genus and species levels were found in Rhizophoraceae. Since this family always has the richest taxa in most mangrove ecosystems in the world, as proved in the present study, this family is referred to as the "true mangrove family" (Duke et al. 1998).




Figure 2. Map of mangrove extent and distribution areas in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia from the image of Landsat-8 OLI analyzed with mixed method of supervised classification and segmentation technique of composite band RGB 654.

Table 2. Composition of true mangrove species recorded in Dudepo and Ponelo Islands, North Gorontalo District, Indonesia

Family	Genus	Species	Dudepo			Ponelo			Status in IUCN Redlist
			Tr	Sap	Seed	Tr	Sap	Seed	
Myrsinaceae	<i>Aegiceras</i>	<i>A. floridum</i> Roem. & Schult ^{*)}	+						Nt
Avicenniaceae	<i>Avicennia</i>	<i>A. alba</i> Blume ^{*)}				+		+	Lc
		<i>A. lanata</i> Ridley ^{**)}	+		+				Vu
Rhizophoraceae	<i>Bruguiera</i>	<i>B. gymnorhiza</i> (L.) Lam. ^{*)}	+		+				Lc
		<i>B. parviflora</i> Wight & Arn. ex Griffith ^{*)}				+			Lc
		<i>B. sexangula</i> (Lour.) Poir. ^{**)}				+	+	+	Lc
		<i>C. decandra</i> (Griff.) Ding Hou ^{*)}	+						Nt
	<i>Rhizophora</i>	<i>C. tagal</i> (Perr) C.B. Rob. ^{*)}	+		+				Lc
		<i>R. apiculata</i> Blume ^{*)}	+		+	+	+	+	Lc
		<i>R. mucronata</i> Lam. ^{*)}	+	+	+	+	+	+	Lc
		<i>R. stylosa</i> Griff. ^{*)}	+	+	+	+	+	+	Lc
Lythraceae	<i>Sonneratia</i>	<i>S. alba</i> Sm. ^{*)}	+			+	+	+	Lc
		<i>S. caseolaris</i> (L.) Engl. ^{*)}				+		+	Lc
		Sum	9	2	6	8	5	7	

Note: ^{*)} = species also recorded in eastern coast, ^{**)} = species is not recorded in eastern coast, Tr=Tree, Sap=Sampling, Seed=Seedling. status of redlist IUCN was referred to <http://www.iucnredlist.org> Vu=vulnerable, Nt= near threatened, Lc= least concern

Comparison the species composition based on mangrove structure in Dudepo and Ponelo Islands, generally, showed that more tree species were recorded in Dudepo Island (9 species) than in Ponelo Island (8 species). The proportions of plants at different growth stages (trees-saplings-seedlings) in the two islands respectively were 52.94%-11.76%-35.29% in Dudepo Island and 40.00%-

25.00%-35.00% in Ponelo Island. In the context of conservation, we recorded 3 important species in Dudepo and Ponelo Islands, namely *Aegiceras floridum* Roem. & Schult and *Avicennia lanata* Ridley in Dudepo Island and *Bruguiera sexangula* (Lour.) Poir in Ponelo Island which are enlisted as near-threatened and vulnerable species in IUCN's red list.

Furthermore, we noted that the current numbers of species in Dudepo and Ponelo Islands were less than those recorded previously in the eastern coast (18 species of 9 genera of 8 families). However, we were getting a total of 21 species so far as the species richness of true mangrove in North Gorontalo District, by adding the current result recorded in both islands with those recorded in the eastern coast. Relatively, this number is higher than that recently reported by Widyastuti et al. (2018) for mangrove in Segara Anakan, Central Java, Indonesia, and also reported by Cañizares and Seronay (2016) for the mangrove richness in Barangay Imelda, Dinagat Island, Philippines. The number of 21 species as biological wealth of true mangrove species in North Gorontalo is almost comparable with that reported by Barik and Chowdhury (2014) for diversity in the Sundarbans Delta, West Bengal, Eastern India, and also that reported by Wouthuyzen and Ahmad (2018) in the Lease Islands (Saparua and Nusalaut Islands), Maluku Province, Indonesia.

Geographically, Maluku and Gorontalo in Sulawesi Island are parts of the same island group located in Wallacea in the center part and eastern area of the Indonesian archipelago. We argue, then, if this region is a high-diversity habitat for mangrove in Indonesia, though, more detailed information of this is needed to confirm it.

Mangrove structure

Comparison between the density of all mangrove plants, and diameter and basal of trees in Dudepo and Ponelo Island are presented in Table 3. While the mean tree density in Dudepo Island was higher (2133 ± 329.78 individuals.ha⁻¹), the mean density of seedlings was higher in Ponelo Island (2963 ± 443.22 individuals.ha⁻¹). The lower sapling density than that of seedlings in both islands was an interesting record. It seems there were factors controlling mangrove seedling survival to the sapling stage in both islands which we have not understood yet. Therefore, we suggest that the critical step in mangrove maturity occurs between the seedling and sapling stages.

Additionally, Koch (1997) proposed soil salinity, aeration, and resource availability may all be important determinants of mangrove seedling development to the sapling stage. By investigating abiotic factors influencing seedling development into sapling stage of *Rhizophora*

mangle L. in South Florida, then, he concluded that low salinity stress, high light availability, and soil fertility were presumably the dominant factors controlling *R. mangle* seedling development to the sapling stage (height > 85 cm), and it was also hypothesized that soil anoxia was an important stressor in lagoonal-bay estuaries and marsh-mangrove ecotones with minimal tidal exchange. In the current research, the abiotic factors we measured included those related to mangrove growth, nevertheless it is misleading to conclude that those to be the factors influencing the phenomenon in both islands. More detailed and focused research related to this phenomenon is needed.

In spite of that confusing phenomenon, by comparing diameter and basal area of trees, we recorded the bigger mean diameter and basal area of trees in Dudepo Island than in Ponelo Island. The biggest diameter and basal area (29.26 ± 15.46 cm and 200.66 ± 3.57 m²ha⁻¹) and also the highest tree density (2333 ± 91.72 individuals.ha⁻¹) were recorded in transects in Hamlet of Upo in Dudepo Island (ST2), which had relatively lower water salinity and temperature than other stations (Table 1).

Species ranks based on their importance value (IV) in Dudepo and Ponelo Islands are presented in Table 4. Based on IV, we found that *Rhizophora stylosa* Griff and *R. mucronata* Lam, respectively, were the dominant and codominant species in both islands. Importance Value Index (IVI) states the role of a plant species in the community. Theoretically, the greater the IVI of a species, the greater the role of the species in the community. If the IVIs are evenly distributed across all species measured, the community has high evenness index which lead to high diversity index. By applying certain limits for IVI, we can assume the diversity index from the number of species filtered at the limit. For communities having similar species richness the lower the number of species filtered at the limit, the lower the diversity index. Applying the limit value of 10% of species IVI, we got lower number of species in Dudepo Island (6 species with IVI range of 11.14-114.87%) than in Ponelo Island (7 species with IVI range of 13.29-82.18%), which indicated the higher mangrove species diversity index in Ponelo Island than in Dudepo Island. To confirm this, then diversity and dominance indexes were calculated.

Table 3. Comparison of mangrove structure among transects on Dudepo and Ponelo Islands.


Island	Code of Transects	Trees (ha ⁻¹)	No. of Species	Saplings. (ha ⁻¹)	No. of Species	Seedlings (ha ⁻¹)	No. of Species	Diameter (cm)*	Basal Area (m ² ha ⁻¹)*
Dudepo	ST1	2167 ± 104.1	3	0 ± 0	-	1267 ± -	1	17.52 ± 4.95	56.38 ± 0.51
	ST2	2333 ± 91.72	6	0 ± 0	-	1533 ± -	4	29.26 ± 15.46	200.66 ± 3.57
	ST3	2000 ± 134.35	3	133 ± 11.6	1	1533 ± 11.6	2	14.09 ± 3.39	34.65 ± 0.27
	ST4	2033 ± 100.94	3	100 ± 9.18	2	1367 ± 9.18	3	15.9 ± 4.59	45.19 ± 0.45
	Mean	2133 ± 329.78		58 ± 13.48		1425 ± 113.96		19.73 ± 10.65	84.22 ± 67.67
Ponelo	ST5	2217 ± 53.21	6	67 ± 2.8	2	3017 ± 87.22	5	19.45 ± 10.7	89.06 ± 0.98
	ST6	2200 ± 106.94	4	33 ± 2.01	1	3000 ± 82.17	4	16.91 ± 5.2	55.72 ± 0.5
	ST7	2000 ± 94.44	4	100 ± 4.59	1	2933 ± 73.92	4	16.73 ± 7.04	55.19 ± 0.68
	ST8	2033 ± 107.06	5	67 ± 2.92	2	4267 ± 126.51	5	17.22 ± 6.49	55.84 ± 0.64
	ST9	2025 ± 68.77	6	125 ± 3.9	4	1900 ± 55.07	6	14.87 ± 6.53	44.53 ± 0.44
	Mean	2111 ± 234.28		79 ± 14.51		2963 ± 443.22		17.04 ± 1.46	60.07 ± 15.12

Note: * diameter and basal area analyzed on trees category only.

Table 4. Comparison of importance value index (IVI) of species and their rank in Dudepo and Ponelo Islands.

Species	Diameter (cm)	Basal area (m ² ha ⁻¹)	Density (ha ⁻¹)	Frequency	IVI	Rank
Dudepo Island						
<i>R. stylosa</i> Griff.	15.81 ± 4.69	22.59 ± 0.11	1025 ± 3.51	1	114.87	1
<i>R. mucronata</i> Lam.	16.72 ± 5.02	14.64 ± 0.13	600 ± 3.34	0.5	65.5	2
<i>A. lanata</i> Ridley **)	32.55 ± 16.67	24.51 ± 1.04	233 ± 2.79	0.33	53.38	3
<i>S. alba</i> Sm.	37.04 ± 14.85	13.55 ± 0.92	108 ± 2.94	0.17	27.83	4
<i>R. apiculata</i> Blume	17.83 ± 3.91	1.75 ± 0.1	67 ± 1.73	0.17	11.86	5
<i>B. gymnorhiza</i> (L.) Lam.	39.73 ± 20.75	5.26 ± 1.2	33 ± 1.73	0.08	11.14	6
<i>C. tagal</i> (Perr) C.B. Rob. *)	20.19 ± 3.93	1.38 ± 0.11	42 ± 2	0.08	6.93	7
<i>C. decandra</i> (Griff.) Ding Hou	15.92 ± 0.64	0.33 ± 0.01	17 ± 1	0.08	4.51	8
<i>A. floridum</i> Roem. & Schult *)	17.83 ± 0	0.21 ± -	8 ± -	0.08	3.97	9
Ponelo Island						
<i>R. stylosa</i> Griff.	15.05 ± 6.18	14.43 ± 0.08	647 ± 16.08	0.68	82.18	1
<i>R. mucronata</i> Lam.	16.16 ± 6.19	15.97 ± 0.1	647 ± 18.18	0.53	77.92	2
<i>S. alba</i> Sm.	24.84 ± 13.74	18.65 ± 0.44	289 ± 13.29	0.42	60.71	3
<i>A. alba</i> Blume	20.02 ± 6.02	7.23 ± 0.11	211 ± 16.64	0.21	30.19	4
<i>R. apiculata</i> Blume	14.7 ± 4.43	2.53 ± 0.05	132 ± 15.89	0.16	16.87	5
<i>B. sexangula</i> (Lour.) Poir.	15.53 ± 6.1	1.84 ± 0.08	79 ± 10.77	0.16	13.29	6
<i>B. parviflora</i> Wight & Arn. ex Griffith	17.69 ± 5.5	1.84 ± 0.09	68 ± 10.11	0.16	12.8	7
<i>S. caseolaris</i> (L.) Engl.	20.79 ± 5.06	1.32 ± 0.1	37 ± 12.89	0.05	6.04	8

Note: *) enlisted as near-threatened (NT) species, while **) was more critical with vulnerable status (Vu) globally in IUCN's red list

**Figure 3.** Comparison of Shannon-Wiener index, Margalef species richness index, and evenness of tree category among transects in Dudepo and Ponelo Island

The indexes of species diversity, species richness and evenness of mangrove trees in all stations

The Shannon Index (H), the Margalef's species richness index, and the Buzas and Gibson evenness index for the tree category in all transects are shown in Figure 3.

Shannon-Wiener diversity index of mangrove trees in Dudepo Island ranged from 0.34 (ST3) to 1.46 (ST2) while in Ponelo Island was from 1.22 (ST6)-1.70 (ST9). This result confirmed the view above related to comparison of mangrove diversity index based on IVI of species in both islands. Further analysis then showed that the diversity index in Ponelo Island which was higher than the highest index in Dudepo Island (1.46) was found in transects in

ST8 (1.50), ST5 (1.52) and ST9 (1.72). Although Ponelo Island had higher diversity index than Ponelo Island, since the Shannon diversity on both islands was less than 2, collectively, it was categorized into the low mangrove diversity.

When comparing the relationship between quadrat number with the indexes of diversity of Shannon, species richness, and species evenness in all transects employed, we didn't find in both Island the pattern which showed the increasing values of those indexes with the increasing sampling area as argued by Ismaini et al. (2015) for plant diversity they observed in upland regions (Dempo Mountain, South Sumatra).




Figure 4. Dendrogram of similarity among transects in Dudepo Island (ST1, ST2, ST3, and ST4) and Ponelo Island (ST5, ST6, ST7, ST8, and ST9).

Multivariate clustering of mangrove

The Bray-Curtis similarity index was used to cluster the transects of mangroves in Dudepo and Ponelo Island. The mangrove variables compared were the indexes of Shannon diversity, the species richness, the Margalef evenness, number of species and quadrats in each transect. The similarity dendrogram among transects in both islands is shown in Figure 4. In this figure, it was known that both islands had similarity of 0.75. Both islands were similar on the lower value, about 0.72. Furthermore, the high dissimilarity of mangrove in Dudepo Island from Ponelo Island were found in ST1, ST3 and ST4. Transects in station of hamlet Upo (ST2) in Dudepo Island, in another side, was more similar to those in Dudepo Island with a similarity value of about 0.78.

With these results, then we concluded that while the mangrove in Dudepo Island was important by the occurrence of important species of IUCN's red list for conservation and also by the bigger diameter of trees, the true mangrove in Ponelo Island was more important by the higher diversity index and better regeneration.

ACKNOWLEDGEMENTS

The research was carried out and granted within frame of Eminent Grand of Basic Research (PDUPT), the funded year 2017-2018 by the Indonesian State Ministry for Research and Technology (RISTEK), Jakarta, Indonesia, Contract Number: 249 / UN47.D / PL / 2018. We thank the following person: Jalipati Tuheteru, Olpien Umar, and Sandrianto (members of the Gorontalo DEHETO club) for help during field survey and data collection activities.

REFERENCES

- Barik J, Chowdhury S. 2014. True mangrove species of Sundarbans Delta, West Bengal, Eastern India. *Check List* 10 (2): 329-334.
- Blanco JF, Bejarano AC, Lasso J, Cantera JR. 1999. A new look at computation of the complexity index in mangroves: do disturbed forests have clues to analyze canopy height patchiness?. *Wetlands Ecol Manag* 9: 91-101.
- BPS Kabupaten Gorontalo Utara. 2017. Kabupaten Gorontalo Utara Dalam Angka 2017. Badan Pusat Statistik Kabupaten Gorontalo Utara. Kwandang. [Indonesian]
- Cañizares LP, Seronay RA. 2016 Diversity and species composition of mangroves in Barangay Imelda, Dinagat Island, Philippines. *AACL Bioflux*, 9 (3): 518-526
- Cintron G, Novelli YS. 1984. Methods for studying mangrove structure. In: Snedaker SC and Snedaker JG (eds) *The mangrove ecosystem: research methods*, pp. 91-113, UNESCO, Paris.
- Duke NC, Ball MC, Ellison JC. 1998. Factors influencing biodiversity and distributional gradients in mangroves. *Global Ecol Biogeogr Lett* 7: 27-47.
- Giri C, Ochieng E, Tieszen LL, Zhu Z, Singh A, Loveland T, Masek J, Duke N. 2011. Status and distribution of mangrove forests of the world using earth. *Global Ecol Biogeogr* 20: 154-159.
- Gorlinski V. 2012 *Encyclopædia Britannica: Gorontalo Province, Indonesia*. <https://www.britannica.com/place/Gorontalo>. [28 October 2017].
- Hammer Ø. 2017. Reference manual Paleontological Statistics Version 3.15. Natural History Museum University of Oslo. Oslo.
- Hutchison J, Manica A, Swetnam R, Balmford A, Spalding M. 2014. Predicting global patterns in mangrove forest biomass. *Conserv Lett* 7 (3): 233-240.
- Ismaini L, Lailati M, Rustandi, Sunandar D. 2015. Analisis komposisi dan keanekaragaman tumbuhan di Gunung Dempo, Sumatera Selatan. *Pros Sem Nas Masy Biodiv Indon* 1 (6): 1397-1402.
- Joshi HG, Ghose M. 2014. Community structure, species diversity, and aboveground biomass of the Sundarbans mangrove swamps. *Trop Ecol* 55 (3): 283-303.
- Kasim F, Nursinar S, Panigoro C, Karim Z, Lamalango A. 2017. True mangrove of North Gorontalo District, Indonesia, their list, status and habitat-structural complexity in easternmost coast area. *AACL Bioflux* 10 (6): 1445-1455.
- Koch MS. 1997. *Rhizophora mangle* L. Seedling development into the sapling stage across resource and stress gradients in subtropical Florida. *Biotropica* 29 (4): 427-439.
- Kusmana C, Onrizal, dan Sudarmadji. 2003. Jenis-Jenis Pohon Mangrove di Teluk Bintuni, Papua. Fakultas Kehutanan, Institut Pertanian Bogor dan PT Bintuni Utama Murni Wood Industries, Bogor. [Indonesian]
- Marbawa IKC, Astarini IA, Mahardika IG. 2014. Analisis Vegetasi Mangrove untuk Strategi Pengelolaan Ekosistem Berkelanjutan di Taman Nasional Bali Barat. *Ecotrophic*, 8 (1): 24-38. [Indonesian]
- Muhtadi A, Rudi HS, Rusdi L, Zulham AH. 2016. Ecological status of mangrove of Sembilan Island, Langkat District, North Sumatra Province. *Depik* 5 (3): 151-163.
- Nfotabong-Atheull A, Din N, Koum LGE, Satyanarayana B, Koedam N, Dahdouh-Guebas F. 2011. Assessing forest products usage and local residents' perception of environmental changes in peri-urban and rural mangroves of Cameroon, Central Africa. *J Ethnobiol Ethnomed* 7: 41. DOI: 10.1186/1746-4269-7-41
- Noor YR, Khazali M dan Suryadiputra NN. 2012. Panduan Pengenalan mangrove di Indonesia (Cetakan ulang ketiga). PHKA/WI-IP, Bogor. [Indonesian]
- Prasetyo DE, Ferbrian KA, Firman Z, Hani SP, Achmad S, Anisa B, Edy S. 2014. Kajian sosio-ekologis kawasan mangrove di Pesisir Pantai Kecamatan Biduk-Biduk, Kalimantan Timur. *Omni-Akuatika*, 13 (18): 1-9. [Indonesian]
- Rugayah S. 2007. Mangrove plant diversity in Sepanjang Island, East Java. *Biodiversitas* 8 (2): 130-134.
- Setiawan H, Rini P, dan R. Garsetiasih. 2017. Perception and attitude of community towards mangrove ecosystem conservation at Tanakeke Island-South Sulawesi. *Jurnal Penelitian Sosial dan Ekonomi Kehutanan* 14 (1): 57-70.
- Snedaker SC. 1984. The Mangroves of Asia and Oceania: Status and Research Planning. In: Soepadmo E, Rao AN, MacIntosh DJ (eds). 1984. *Proceedings of the Asian Symposium on Mangrove*

- Environment Research and Management, Kuala Lumpur, August 25-29, 1980.
- Wetland International. 2017. Mangrove. <http://www.wetlands.or.id/mangrove/mangrove.php>. [6 September 2017].
- Widyastuti A, Yani D, Nasution EK, Rochmatino. 2018. Diversity of mangrove vegetation and carbon sink estimation of Segara Anakan Mangrove Forest, Cilacap, Central Java, Indonesia. Biodiversitas 19 (1): 246-252.
- Winata A, Yuliana E, Rusdiyanto E. 2017 Diversity and natural regeneration of mangrove vegetation in the tracking area on Kemujan Island, Karimunjawa National Park, Indonesia. AES Bioflux 9 (2): 109-119.
- Wouthuyzen S. dan Ahmad F. 2018. Mangrove Mapping Of The Lease Islands, Maluku Province Using Multi-Temporal And Multi-Sensor Of Landsat Satellite Images. Oseanologi dan Limnologi di Indonesia 3 (1): 19-37.
- Zhang CG, Leung KK, Wong YS., Tam NFY. 2007. Germination, growth and physiological responses of mangrove (*Bruguiera gymnorhiza*) to lubricating oil pollution. Environ Exp Bot 60: 127-136.

***Front cover: Nasalis larvatus* male**
(PHOTO: CHARLES J SHARP)

Published bimonthly

PRINTED IN INDONESIA

ISSN: 1412-033X



E-ISSN: 2085-4722

