



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name



Home

Journal Rankings

Country Rankings


Viz Tools

Help

About Us

AACL Bioflux

Country

Romania - 

14

H Index

Subject Area and Category

Agricultural and Biological Sciences
 Aquatic Science
 Ecology, Evolution, Behavior and Systematics

Environmental Science
 Management, Monitoring, Policy and Law
 Water Science and Technology

Publisher

Bioflux Publishing House

Publication type

Journals

ISSN

18449166, 18448143

Coverage

2006, 2009-2020


Scope

Information not localized








[Join the conversation about this journal](#)

Quartiles +



 SJR +

Citations per document +



AAFL Bioflux

Q3

Aquatic Science

best quartile

SJR 2019

0.28


powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

Metrics based on Scopus® data as of April 2020



AACL Bioflux

Instructions to authors

Submission letter

Model of paper

Reviewer information pack

Editorial Board Expanded

Coverage / databases

Volume 13(6)/2020

Volume 13(5)/2020 (October, 30)

Volume 13(4)/2020 (August, 30)

Volume 13(3)/2020 (June, 30)

Volume 13(2)/2020 (April, 30)

Volume 13(1)/2020 (February, 28)

Volume 12(6)/2019 (December, 30)

Volume 12(5)/2019 (October, 30)

Volume 12(4)/2019 (August, 30)

Volume 12(3)/2019 (June, 30)

Volume 12(2)/2019 (April, 30)

Volume 12(1)/2019 (February, 28)

Volume 11(6)/2018 (December, 30)

Volume 11(5)/2018 (October, 30)

Volume 11(4)/2018 (August, 30)

Volume 11(3)/2018 (June, 30)

Volume 11(2)/2018 (April, 30)

Volume 11(1)/2018 (February, 28)

Volume 10(6)/2017 (December, 30)

Volume 10(5)/2017 (October, 30)

Volume 10(4)/2017 (August, 30)

Aquaculture, Aquarium, Conservation & Legislation - International Journal of the Bioflux Society

ISSN 1844-9166 (online)

ISSN 1844-8143 (print)

Published by Bioflux - bimonthly -

in cooperation with The Natural Sciences Museum Complex (Constanta, Romania)

Peer-reviewed (each article was independently evaluated before publication by two specialists)

The journal includes original papers, short communications, and reviews on Aquaculture (Biology, Technology, Economics, Marketing), Fish Genetics and Improvement, Aquarium Sciences, Fisheries, Ichthyology, Aquatic Ecology, Conservation of Aquatic Resources and Legislation (in connection with aquatic issues) from wide world.

The manuscripts should be submitted to zoobiomag2004@yahoo.com

Editor-in-Chief:

Petrescu-Mag I. Valentin: USAMV Cluj, Cluj-Napoca, University of Oradea (Romania)

Gavriloaie Ionel-Claudiu (reserve): SC Bioflux SRL, Cluj-Napoca (Romania).

Editors:

Abdel-Rahim Mohamed M.: National Institute of Oceanography and Fisheries, Alexandria (Egypt)

Adascalitei Oana: Maritime University of Constanta, Constanta (Romania)

Amira Aicha Beya: Badji Mokhtar Annaba University, Annaba (Algeria)

Arockiaraj A. Jesu: SRM University, Chennai (India)

Appelbaum Samuel: Ben-Gurion University of the Negev (Israel)

Baharuddin Nursalwa: Universiti Malaysia Terengganu, Terengganu (Malaysia)

Boaru Anca: USAMV Cluj, Cluj-Napoca (Romania)

Botha Miklos: Bioflux SRL, Cluj-Napoca (Romania)

Volume 10(3)/2017 (June, 30)	Breden Felix: Simon Fraser University (Canada)
Volume 10(2)/2017 (April, 30)	Burny Philippe: Universite de Liege, Gembloux (Belgium)
Volume 10(1)/2017 (February, 28)	Caipang Cristopher M.A.: Temasek Polytechnic (Singapore)
Volume 9(6)/2016 (December, 30)	Chapman Frank: University of Florida, Gainesville (USA)
Volume 9(5)/2016 (October, 30)	Coroian Cristian: USAMV Cluj, Cluj-Napoca (Romania)
Volume 9(4)/2016 (August, 30)	Creanga Steofil: USAMV Iasi, Iasi (Romania)
Volume 9(3)/2016 (June, 30)	Cristea Victor: Dunarea de Jos University of Galati, Galati (Romania)
Volume 9(2)/2016 (April, 30)	Das Simon Kumar: Universiti Kebangsaan Malaysia, Bangi, Selangor (Malaysia)
Volume 9(1)/2016 (February, 28)	Dimaggio Matthew A.: University of Florida (USA)
Volume 8(6)/2015 (December, 30)	Firica Cristian Manuel: Spiru Haret University Bucharest, Craiova (Romania)
Volume 8(5)/2015 (October, 30)	Georgescu Bogdan: USAMV Cluj, Cluj-Napoca (Romania)
Volume 8(4)/2015 (August, 30)	Karayucel Ismihan: University of Sinop, Sinop (Turkey)
Volume 8(3)/2015 (June, 30)	Khamesipour Faham: Shiraz University, Shiraz (Iran)
Volume 8(2)/2015 (April, 30)	Kosco Jan: Presov University, Presov (Slovakia)
Volume 8(1)/2015 (February, 28)	Kovacs Eniko: USAMV Cluj, Cluj-Napoca (Romania)
Volume 7(6)/2014 (December, 30)	Mehrad Bahar: Gorgan University of Agricultural Sciences and Nat. Res. (Iran)
Volume 7(5)/2014 (October, 30)	Miclaus Viorel: USAMV Cluj, Cluj-Napoca (Romania)
Volume 7(4)/2014 (August, 30)	Mihociu Tamara: R&D National Institute for Food Bioresources (Romania)
Volume 7(3)/2014 (June, 30)	Molnar Kalman: Hungarian Academy of Sciences, Budapest (Hungary)
Volume 7(2)/2014 (April, 15)	Muchlisin Zainal Abidin: Universiti Sains (Malaysia), Syiah Kuala University (Indonesia)
Volume 7(1)/2014 (February, 15)	Muntean George Catalin: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(6)/2013 (November, 15)	Nowak Michal: University of Agriculture in Krakow (Poland)
Volume 6(5)/2013 (September, 15)	Nyanti Lee: Universiti Malaysia Sarawak, Sarawak (Malaysia)
Volume 6(4)/2013 (July, 25)	Olivotto Ike: Universita Politecnica delle Marche, Ancona (Italy)
Volume 6(3)/2013 (May, 15)	Oroian Firuta Camelia: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(2)/2013 (March, 15)	Papuc Tudor: USAMV Cluj, Cluj-Napoca (Romania)
Volume 6(1)/2013 (January, 15)	Parvulescu Lucian: West University of Timisoara (Romania)
Volume 5(5)/2012 (December, 30)	Pasarin Benone: USAMV Iasi, Iasi (Romania)
Volume 5(4)/2012 (September, 30)	Pattikawa Jesaja Ajub: Pattimura University, Ambon (Indonesia)
Volume 5(3)/2012 (July, 30)	Petrescu Dacinia Crina: Babes-Bolyai University, Cluj-Napoca (Romania)
Volume 5(2)/2012 (June, 30)	Petrescu-Mag Ruxandra Malina: Babes-Bolyai University, Cluj-Napoca (Romania)
Volume 5(1)/2012 (March, 15)	
Volume 4(5)/2011 (December, 30)	
Volume 4(4)/2011 (October, 30)	

- Volume 4(3)/2011 (July, 30)
- Volume 4(2)/2011 (April, 30)
- Volume 4(1)/2011 (January, 30)
- Volume 3(5)/2010 (December, 5)
- Volume 3(4)/2010 (December, 1)
- Volume 3(3)/2010 (November, 15)
- Volume 3(2)/2010 (July, 30)
- Volume 3(1)/2010 (February, 28)
- Volume 2(4)/2009 (October, 30)
- Volume 2(3)/2009 (July, 30)
- Volume 2(2)/2009 (April, 30)
- Volume 2(1)/2009 (January, 30)
- Volume 1(2)/2008 (December, 30)
- Volume 1(1)/2008 (September, 30)
- Volume Pilot/2007 (December, 30) - available printed only
- Pontus Euxinus, Volume 1 (1980) - Parent Journal

- Petrovici Milca: West University of Timisoara (Romania)
- Pratasik Silvester Benny: Sam Ratulangi University, Manado (Indonesia)
- Proorocu Marian: USAMV Cluj, Cluj-Napoca (Romania)
- Ray Sunuram: Khulna University (Bangladesh)
- Rhyne Andrew: Roger Williams University; New England Aquarium, Boston (USA)
- Ruchin Alexander B.: Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Saransk (Russia)
- Safirescu Calin: USAMV Cluj, Cluj-Napoca (Romania)
- Serrano Jr. Augusto E.: University of the Philippines Visayas (Philippines)
- Sima Nicusor Flaviu: USAMV Cluj, Cluj-Napoca (Romania)
- Tlusty Michael F.: New England Aquarium, Boston (USA)
- Vesa Stefan Cristian: Iuliu Hatieganu UMF, Cluj-Napoca (Romania)
- Vintila Iuliana: Dunarea de Jos University of Galati, Galati (Romania)
- Wariaghli Fatima: University Mohammed V in Rabat, Rabat (Morocco)
- Yusli Wardiatno: Bogor Agricultural University, Bogor (Indonesia).


AACL Bioflux

Q3 Aquatic Science
best quartile

SJR 2019
0.28

powered by scimagojr.com

design: www.simple-webdesign.com



AACL Bioflux

Instructions to authors

Submission letter

Model of paper

Reviewer information pack

Editorial Board Expanded

Coverage / databases

Volume 13(6)/2020

Volume 13(5)/2020 (October, 30)

Volume 13(4)/2020 (August, 30)

Volume 13(3)/2020 (June, 30)

Volume 13(2)/2020 (April, 30)

Volume 13(1)/2020 (February, 28)

Volume 12(6)/2019 (December, 30)

Volume 12(5)/2019 (October, 30)

Volume 12(4)/2019 (August, 30)

Volume 12(3)/2019 (June, 30)

Volume 12(2)/2019 (April, 30)

Volume 12(1)/2019 (February, 28)

Volume 11(6)/2018 (December, 30)

Volume 11(5)/2018 (October, 30)

Volume 11(4)/2018 (August, 30)

Volume 11(3)/2018 (June, 30)

Volume 11(2)/2018 (April, 30)

Volume 11(1)/2018 (February, 28)

Volume 10(6)/2017 (December, 30)

Volume 10(5)/2017 (October, 30)

Volume 10(4)/2017 (August, 30)

Volume 13(5)/2020

Koniyo Y., Juliana, Pasisingi N., Kalalu D., 2020 The level of parasitic infection and growth of red tilapia (*Oreochromis sp.*) fed with vegetable fern (*Diplazium esculentum*) flour. AACL Bioflux 13(5):2421-2430.

Efendi D. S., Adrianto L., Yonvitner, Wardiatno Y., Agustina S., 2020 The performance of stock indicators of grouper (Serranidae) and snapper (Lutjanidae) fisheries in Saleh Bay, Indonesia. AACL Bioflux 13(5):2431-2444.

Sari L. D., Fadjar M., Widodo M. S., Lutfiatunnisa, Valen F. S., 2020 Growth analysis of Asian seabass (*Lates calcarifer* Bloch 1790) based on Morphometrics in BPBAP Situbondo, East Java. AACL Bioflux 13(5):2445-2451.

Ndobe S., Yasir I., Salanggon A. I. M., Wahyudi D., Ederyan, Muslihudin, Renol, Adel Y. S., Moore A. M., 2020 Eucheumatoid seaweed farming under global change - Tomini Bay seaweed trial indicates *Eucheuma denticulatum (spinosum)* could contribute to climate adaptation. AACL Bioflux 13(5):2452-2467.

Subandiyono S., Hastuti S., 2020 Dietary protein levels affected on the growth and body composition of tilapia (*Oreochromis niloticus*). AACL Bioflux 13(5):2468-2476.

Longdong F. V., Mantjoro E., Kepel R. C., Budiman J., 2020 Adaptation strategy of Bitung fishermen to the impact of fisheries Moratorium policy in Indonesia. AACL Bioflux 13(5):2477-2496.

Zainuddin, Aslamyah S., Nur K., Hadijah, 2020 Substitution of sweet potato flour and corn starch to the growth, survival rate, feed conversion ratio and body chemical composition of juvenile *Litopenaeus vannamei*. AACL Bioflux 13(5):2497-2508.

Leilani A., Restuwati I., 2020 Analysis of the benefits of information and communication technology in extension activities in the District and City of Cirebon, West Java Province, Indonesia. AACL Bioflux 13(5):2509-2521.

Suherman A., Santosa M. A., Ihsan Y. N., Wijayanto D., Juwana S., 2020 The eradication of IUU fishing in Indonesia for fisheries resources sustainability by the Task Force 115. AACL Bioflux 13(5):2522-2537.

Mustikasari D., Nuryanto A., Suryaningsih S., 2020 The existence of blue panchax (*Aplocheilichthys panchax*) in the abandoned tin mining pits water with different age contaminated by heavy metals. AACL Bioflux 13(5):2538-2550.

Volume 10(3)/2017 (June, 30)
Volume 10(2)/2017 (April, 30)
Volume 10(1)/2017 (February, 28)
Volume 9(6)/2016 (December, 30)
Volume 9(5)/2016 (October, 30)
Volume 9(4)/2016 (August, 30)
Volume 9(3)/2016 (June, 30)
Volume 9(2)/2016 (April, 30)
Volume 9(1)/2016 (February, 28)
Volume 8(6)/2015 (December, 30)
Volume 8(5)/2015 (October, 30)
Volume 8(4)/2015 (August, 30)
Volume 8(3)/2015 (June, 30)
Volume 8(2)/2015 (April, 30)
Volume 8(1)/2015 (February, 28)
Volume 7(6)/2014 (December, 30)
Volume 7(5)/2014 (October, 30)
Volume 7(4)/2014 (August, 30)
Volume 7(3)/2014 (June, 30)
Volume 7(2)/2014 (April, 15)
Volume 7(1)/2014 (February, 15)
Volume 6(6)/2013 (November, 15)
Volume 6(5)/2013 (September, 15)
Volume 6(4)/2013 (July, 25)
Volume 6(3)/2013 (May, 15)
Volume 6(2)/2013 (March, 15)
Volume 6(1)/2013 (January, 15)
Volume 5(5)/2012 (December, 30)
Volume 5(4)/2012 (September, 30)
Volume 5(3)/2012 (July, 30)
Volume 5(2)/2012 (June, 30)
Volume 5(1)/2012 (March, 15)
Volume 4(5)/2011 (December, 30)
Volume 4(4)/2011 (October, 30)

Wijayanto D., Bambang A. N., Nugroho R. A., Kurohman F., 2020 The growth model of *Eucheuma cottonii* cultivated in Karimunjawa Islands, Indonesia. *AACL Bioflux* 13(5):2551-2557.

Hadijah, Zainuddin, Aqmal A., Banin D. K., 2020 The effect of marine algae (*Gracilaria verrucosa*) formulated feed on the growth rate, survival rate and chemical composition of abalone (*Haliotis squamata*) reared in marine submersible cages. *AACL Bioflux* 13(5):2558-2566.

Huda M. A., Sunarno M. T. D., Nurhudah M., 2020 Potential addition of black soldier fly carcass meal in sangkuriang catfish (*Clarias gariepinus*) feed formulation. *AACL Bioflux* 13(5):2567-2576.

Haeruddin, Supriharyono, Rahman A., Ghofar A., Iryanthony S. B., 2020 Spatial distribution and heavy metal pollution analysis in the sediments of Garang watershed, Semarang, Central Java, Indonesia. *AACL Bioflux* 13(5):2577-2587.

El Fajri N., Sukendi, Syafriadiman, Windarti, Yusuf M., 2020 Sustainability assessment of swamp ecosystem management in Sawah village, Kampar Regency, Riau. *AACL Bioflux* 13(5):2588-2598.

Mojumder N., Saha D., Utsa S. S., Maruf M. K. K., Paul S. K., 2020 Biology of the endangered queen loach (*Botia dario*) collected from wild sources in Bangladesh. *Bioflux* 13(5):2599-2609.

Setiyono J., Azhar M., Solechan, Trihastuti N., Hakim A. R., 2020 Justification of the ship-sinking policy in the territorial jurisdiction of Indonesia. *AACL Bioflux* 13(5):2610-2618.

Tyas D. E., Haeruddin, Purnomo P. W., 2020 Isolation and identification of biofilm bacteria from bamboo in traditional vannamei shrimp ponds as an ammonia degradation factor. *AACL Bioflux* 13(5):2619-2629.

Riyadi P. H., Romadhon R., Anggo A. D., Herawati V. E., Setyastuti A. I., 2020 PASS and ADMET analyses for eight compounds from Nile tilapia (*Oreochromis niloticus*) viscera waste hydrolysate as anti-inflammatory nutraceutical. *AACL Bioflux* 13(5):2630-2638.


Pangemanan N. P. L., Kepel R. C., Bataragoa N. E., Tumbol R. A., Sahami F. M., Lumenta C., Salaki M. S., Boneka F. B., Mantiri D. M. H., 2020 Morphometric and molecular analysis for identification of the early stages of gobioid fishes in the Poigar River estuary, North Sulawesi, Indonesia. *AACL Bioflux* 13(5):2639-2646.

Muhtadi A., Leidonald R., Rahmadya A., Lukman, 2020 Bathymetry and morphometry of Siais Lake, South Tapanuli, North Sumatra Province, Indonesia. *AACL Bioflux* 13(5):2647-2656.

Ratunil V. B. Jr., Gamboa G. Z. Jr., Ebarsabal G. A., 2020 Does the peak spawning activity of *Glossogobius giuris* in Lake Mainit occur in the early months of a year? *AACL Bioflux* 13(5):2657-2663.

Fahmy A. F., El-Greisy Z. A., Moharram S. G., 2020 Impact of hormonal manipulation on egg quality of *Diplodus sargus*: comparative ultrastructural

Volume 4(3)/2011 (July, 30)
Volume 4(2)/2011 (April, 30)
Volume 4(1)/2011 (January, 30)
Volume 3(5)/2010 (December, 5)
Volume 3(4)/2010 (December, 1)
Volume 3(3)/2010 (November, 15)
Volume 3(2)/2010 (July, 30)
Volume 3(1)/2010 (February, 28)
Volume 2(4)/2009 (October, 30)
Volume 2(3)/2009 (July, 30)
Volume 2(2)/2009 (April, 30)
Volume 2(1)/2009 (January, 30)
Volume 1(2)/2008 (December, 30)
Volume 1(1)/2008 (September, 30)
Volume Pilot/2007 (December, 30) - available printed only
Pontus Euxinus, Volume 1 (1980) - Parent Journal



changes. *AACL Bioflux* 13(5):2664-2675.

Rundupadang S. S., Mallawa A., Kurnia M., 2020 Effectiveness of milkfish (*Chanos chanos*) live bait from aquaculture versus live bait from nature in pole and line fisheries. *AACL Bioflux* 13(5):2676-2686.

Pandelaki L., Rompas R. M., Rembet U. N. W. J., Wantasen A. S., Gerung G. S., Ngangi E. L. A., 2020 Economic value of seagrass ecosystem in Nain Island, South Minahasa Regency, North Sulawesi, Indonesia. *AACL Bioflux* 13(5):2687-2693.

Harbach H., Palm H. W., 2020 Fully controlled experimental recirculating aquaculture system (RAS) for experimental studies with mussels (*Mytilus edulis*-like), focusing on temperature and salinity regimes. *AACL Bioflux* 13(5):2694-2704.

Priyambada A., Fitri A. D. P., Ghofar A., 2020 Potential fishing grounds for *Portunus pelagicus* based on oceanographic factors of the Tukak Sadai waters, Bangka Belitung, Indonesia. *AACL Bioflux* 13(5):2705-2716.

Mulia D. S., Isnansetyo A., Pratiwi R., Asmara W., 2020 Molecular characterization of *Aeromonas caviae* isolated from catfish (*Clarias* sp.). *AACL Bioflux* 13(5):2717-2732.

Jalil J., Makkatenni M., Juhardi J., 2020 Diversity index, similarity index and dominance index of macrozoobenthos in Pangkajene River estuary, Pangkep Regency, Indonesia. *AACL Bioflux* 13(5):2733-2737.

Rachmawati D., Hutabarat J., Samidjan I., Windarto S., 2020 Utilization of papain as feed additive in the fish feed on activity of digestive enzymes, contents of nutrient and minerals of Sangkuriang catfish (*Clarias gariepinus* var. *Sangkuriang*). *AACL Bioflux* 13(5):2738-2744.

Muskananfol M. R., Purnomo P. W., Sulardiono B., 2020 Impacts of environmental factors on macrobenthos distribution and abundance in mangrove ecosystems on the Northern Coast of Java. *AACL Bioflux* 13(5):2745-2756.

Rifqi M., Widigdo B., Mashar A., Nazar F., Wardiatno Y., 2020 Strategy to gain the target of shrimp production on Karawang District coastal area. *AACL Bioflux* 13(5):2757-2769.

Ramadhani D. A., Nugraha M. F. I., Novita H., Rajamuddin M. A. L., Elya B., 2020 Chemical compounds screening of leaves extract from *Eleocharis dulcis* (Burm.f.) Trin. ex Hensch and *in vitro* antibacterial pathogenic test for fish. *AACL Bioflux* 13(5):2770-2778.

Achmad M. J., Subur R., Supyan, Akbar N., 2020 DNA barcode and phylogenetics of green humphead parrotfish (*Bolbometopon muricatum*) in North Maluku waters. *AACL Bioflux* 13(5):2779-2787.

Paransa D. J., Mantiri D. M. H., Kepel R. C., Rumengan A., 2020 Pigment concentration of red algae, *Kappaphycus alvarezii* (Doty) Doty ex Silva during

the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia. *AACL Bioflux* 13(5):2788-2797.

Purnomo A. H., Subaryono, Utomo B. S. B., Paul N., 2020 Institutional arrangement for quality improvement of the Indonesian *Gracilaria* seaweed. *AACL Bioflux* 13(5):2798-2806.

Dangeubun J. L., Letsoin P. P., Syahailatua D. Y., 2020 Growth of *Nannochloropsis* sp. in culture media enriched with shrub-like annual *Clerodendrum minahassae* leaf extract. *AACL Bioflux* 13(5):2807-2815.

Kurohman F., Wijayanto D., Jayanto B. B., 2020 The strategy of capture fisheries development in Pamekasan Regency, Madura Island. *AACL Bioflux* 13(5):2816-2827.

Lutfiatunnisa, Widodo M. S., Faqih A. R., Sari L. D., Valen F. S., 2020 Molecular approach for identification of Asian seabass (*Lates calcarifer* Bloch 1790) based on COI gene sequence from Java, Indonesia. *AACL Bioflux* 13(5):2828-2834.

Picaulima S. M., Wiyono E. S., Baskoro M. S., Riyanto M., 2020 Fleets dynamics of small-scale fisheries in eastern Kei Kecil Island, Maluku Province, Indonesia. *AACL Bioflux* 13(5):2835-2851.

Parvez I., Mahajebin T., Barman T. R., Khan N., Chhanda M. S., Alam M. A., 2020 Evaluation of body shape variation of indigenous and exotic *Anabas testudineus* (Bloch, 1792) in Bangladesh by geometric morphometric analyses. *AACL Bioflux* 13(5):2852-2858.

Sahami F. M., Habibie S. A., 2020 Exploration of adult phase of Nike fish to maintain its sustainability in Gorontalo Bay waters, Indonesia. *AACL Bioflux* 13(5):2859-2867.

Nur M., Rifa'i M. A., Yunita R., Sofia L. A., 2020 Feasibility of floating cage culture based on business scale in Riam Kanan Reservoir, South Kalimantan Province. *AACL Bioflux* 13(5):2868-2877.

Arsad S., Sari L. A., Suherman S. P., Cahyani D., Nadhira T., Yulinda E. N., Musa M., Lusiana E. D., Prasetya F. S., 2020 Utilization of tofu wastewater as *Chlorella pyrenoidosa* growth medium. *AACL Bioflux* 13(5):2878-2885.

Ardiansyah, Amrullah, Dahlia, Jaya A. A., Indrayani, Wahidah S., Hamal R., Jabbar F. B. A., 2020 The selected facultative mixotrophic sulphur-oxidizing bacteria from intensive shrimp ponds. *AACL Bioflux* 13(5):2886-2896.

Paulus C. A., Soewarlan L. C., Ayubi A. A., 2020 Distribution of marine debris in mangrove ecotourism area in Kupang, East Nusa Tenggara, Indonesia. *AACL Bioflux* 13(5):2897-2909.

Turisno B. E., Mahmudah S., Yunanto, 2020 The regulaiton and policy models of Pekalongan Local Government toward mangrove conservation. *AACL Bioflux* 13(5):2910-2920.

Trisyani N., Yusan L. Y., 2020 Proximate analysis and amino acid profile in fresh

meat, meat meal, and shell meal of bamboo clam *Solen* sp. from Kwanyar Coast, Bangkalan-Madura, Indonesia. *AACL Bioflux* 13(5):2921-2927.

Vahira A. D., Riadin A. H., Sarida M., Utomo D. S. C., Setiawan W. A., 2020 Growth performance and sex ratio of *Channa striata* through immersion and bioencapsulation of *Artemia* with recombinant growth hormone. *AACL Bioflux* 13(5):2928-2936.

Zuraida I., Pamungkas B. F., 2020 Effects of acid pretreatment and extraction temperature on the properties of gelatin from striped snakehead (*Channa striata*) scales. *AACL Bioflux* 13(5):2937-2945.

Taridala S. A. A., Nursavista R., Kurnia A., Purbaningsih Y., Alzarliani W. O., Rayuddin, Hartati, 2020 Abalone marketing and its performance during Covid-19 pandemic. *AACL Bioflux* 13(5):2946-2955.

Thovyan A. I., Tapilatu R. F., Sabariah V., Venables S. K., 2020 Plankton abundance and community structure in reef manta ray (*Mobula alfredi*) feeding habitat in the Dampier Strait, Raja Ampat, West Papua, Indonesia. *AACL Bioflux* 13(5):2956-2969.

Sandor Z. J., Biro J. N., Revesz N., Ronyai A., 2020 Comparison of carp filet quality produced in semi-intensive pond system using different type of feeds. *AACL Bioflux* 13(5):2970-2981.

Kurniawan A., 2020 The metal oxides of abandoned tin mining pit waters as an indicator for bacterial diversity. *AACL Bioflux* 13(5):2982-2992.


Latiu C., Cocan D., Uiuu P., Matei M. C., Constantinescu R., Heghedus C., Ihut A., Raducu C., Nicula A. S., Miresan V., 2020 Prespawning and postspawning hematological parameters and oxidative stress biomarkers of brown trout (*Salmo trutta*) (Pisces: Salmonidae), from Somesul Cald River. *AACL Bioflux* 13(5):2993-3003.

Pronina G. I., Mannapov A. G., 2020 Some aspects of fish breeding for immune resistance as shown through a study of common carp. *AACL Bioflux* 13(5):3004-3014.

Wakiah A., Mallawa A., Amir F., 2020 Population dynamics of snakehead fish (*Channa striata*) in Tempe Lake, South Sulawesi, Indonesia. *AACL Bioflux* 13(5):3015-3027.

...

design: www.simple-webdesign.com



Exploration of adult phase of Nike fish to maintain its sustainability in Gorontalo Bay waters, Indonesia

Femy M. Sahami, Sitty A. Habibie

Faculty of Fisheries and Marine Sciences, Gorontalo State University, Gorontalo, 96128, Indonesia. Corresponding author: F. M. Sahami, femysahami@ung.ac.id

Abstract. Nike is an amphidromous goby species that migrates from the Gorontalo Bay waters to the Bone Bolango River and metamorphoses into adult goby. This fish is very popular, and it becomes people's favorite as delicious food and turns into an important economic commodity in Gorontalo. Therefore, the exploration of Nike fish's adult phase as an essential phase in the availability of Nike in the Gorontalo Bay waters needs to be done. This study aims to determine the types of goby as the adult phase of Nike fish in the Bone Bolango River waters based on genetic analysis of the mitochondrial DNA COI gene. Sampling was carried out in April and June 2020. The captured samples are then grouped according to their morphological similarity (shape, color, and body pattern). Each of the two samples from each morphological group was preserved in a 95% ethanol solution for genetic identification needs. This study succeeded in finding the adult phase of six species from a total of nine species of a composer for Nike fish schools in Gorontalo; *Sicyopterus longifilis*, *S. lagocephalus*, *Belobranchus belobranchus*, *B. segura*, *Bunaka gyrinoides*, and *Stiphodon semoni*. *S. longifilis* and *S. lagocephalus* species are fish with the local name Busalo; *B. segura*, *B. belobranchus*, and *B. gyrinoides* are locally named Hundala; and *S. semoni* is locally named Timudu'o. The sustainability of Nike's resources can be done with proper management of this adult phase of Nike fish in the river, such as not overfishing goby or releasing goby that are carrying eggs into the river.

Key Words: amphidromous, Busalo, COI gene, gobi, Hundala, Nike, Timudu'o.

Introduction. Nike fish is a typical commodity found in Gorontalo which has important economic value and is commonly favored by the local community as foodstuff due to its good taste and distinction. It belongs to an amphidromous goby group in the postlarva to juvenile stage. Amphidromy is a diadromous pattern of life, the adult spawn in freshwater and their larvae drift downstream into the sea; larvae develop in the marine environment then migrate back into rivers to grow and spawn (Keith et al 2008; Yamasaki et al 2011; Taillebois et al 2012; Iida et al 2017). In tropical rivers, amphidromous fish species are major contributors to the diversity of ecosystems (Ellien et al 2014; Lejeune et al 2016).

Olii et al (2017) and Pasingi & Abdullah (2018) report that Nike fish in the Gorontalo Bay waters will firstly appear in the sea and then move closer to the river mouth over time until they eventually disappear. Nike's high fishing effort should be balanced with resources and habitat ecology management to ensure its sustainability in the future. Exploring the adult phase of fish is one effort that can be conducted to understand the fish biology and ecology. Thus, the management and conservation plans in preserving stocks and biodiversity can be carried out properly. Many studies recently focused on Nike fish (Usman 2016; Olii et al 2017; Pasingi & Abdullah 2018; Olii et al 2019; Nurjirana et al 2019; Sahami et al 2019a, b; Pasingi et al 2020b; Sahami et al 2020). Whereas adult goby as a potential Nike fish parent is also a commodity that plays an important role in the availability of Nike in the Gorontalo Bay waters.

The DNA barcode method in species identification is used to overcome the limitations of species identification morphologically, including phenotypic plasticity, specific life stages of identification cues, small size, often cryptic ecology, and the possible emergence of new species (Viswambharan et al 2013; Linh et al 2018). The cytochrome oxidase subunit I (COI) gene is the fastest and most reliable gene used as a

barcoding marker to identify and discriminate against species that are very closely related to various animal phyla (Hebert et al 2004; Hubert et al 2008; Viswambharan et al 2013; Linh et al 2018). This study aims to determine the types of gobies as the adult phase of Nike fish in the Bone Bolango River waters based on genetic analysis of the mitochondrial DNA COI gene.

Material and Method

Sampling. Goby fish sampling was carried out in the Bone Bolango River (Figure 1) which was suspected to be an adult phase area of the Nike fish. Sampling was carried out in April and June 2020 by using a hand net and modified electric fishing (12V, 10A) with a radius and depth of less than 1 m. The samples obtained were filled in a labeled zip-lock plastic, then, it was stored and given some ice in a cool box. Sample sorting was carried out at the Hydrology and Biometrics Laboratory of the Faculty of Fisheries and Marine Sciences, Gorontalo State University by grouping caught gobies based on their morphological similarity (shape, color, and body pattern). In total, there were 14 groups of caught fish. Two samples from each morphological group were preserved in a 95% ethanol solution for genetic identification needs.




Figure 1. Map of the research location.

DNA extraction, PCR amplification, and sequencing. Molecular analysis was carried out in the Bionesia Laboratory, Denpasar, through several stages, including a collection of fish tissue, DNA isolation, DNA Polymerase Chain Reaction (PCR), electrophoresis, and DNA sequencing. Isolation of DNA samples was carried out using Qiagen Tissue and Blood Extraction kits following the protocol kit. The mitochondrial DNA Cytochrome Oxidase subunit I (COI) gene was chosen because the resolution of the COI gene at the intraspecific level is better than other core genes. Therefore, it is best to be used to identify species to the intraspecific level (Strüder-Kypke & Lynn 2010).

The mitochondrial DNA COI gene was further amplified using a forward primer FishF1 5'-TCA ACC AAC CAC AAA GAC ATT GGC AC-3' and reverse primer FishR1 5'-TAG ACT TCT GGG TGG CCA AAG AAT CA-3' (Ward et al 2005). Three samples, namely B5, B6, and H7 were amplified using a pair of forwarding primer FF2d 5'-TTC TCC ACC AAC CAC AAR GAY ATY GG-3' and reverse primer FR1d 5'-CAC CTC AGG GTG TCC GAA RAA YCA RAA-3' (Ivanova et al 2007). The PCR profiles used were predenaturation at 94°C for

five minutes, denaturation at 94°C for 30 seconds, primary annealing at 50°C for 30 seconds, elongation at 72°C for 30 seconds, and final elongation at 72°C for seven minutes. This PCR process lasted for 38 cycles. DNA samples that have been amplified and electrophoresed would be sequenced afterward.

Data analysis. The samples that had been amplified by the PCR method were then sequenced by using the Dideoxy Sanger Termination Method through PT Genetika Science Indonesia. Nucleotide sequences from DNA sequencing that have been processed and carried out by CONTIG were then matched BLAST with data available in the GenBank database (www.ncbi.nlm.nih.gov) through the BLAST (Basic Local Alignment Search Tool). The phylogenetic tree was arranged by aligning the identified DNA sequence samples with DNA samples of composer's species for Nike fish schools that were available in the GenBank database (Acc. Num. MN069305.1; MN069306.1; MN069307.1; MN069308.1; MT227827.1; MT227828.1; MT227829.1; MT706639.1; MT706640.1; MT706641.1; MT706720.1; MT706721.1; MT706722.1; MT706723.1; MT706724.1; MT706725.1; MT706726.1; and MT706791.1). The phylogenetic tree was created using the Maximum Likelihood 1000 bootstrap method in MEGA X software (Kumar et al 2018).

Results and Discussion. The local community and general public believe that Nike is a fish with maximum body size; thus juvenile size has become a parent size for Nike fish. Whereas Nike fish as an amphidromous species migrates from the Gorontalo Bay waters to the Bone Bolango River and metamorphoses into an adult goby. The results of the molecular analysis indicate that Nike fish schools are composed by nine species so far; *Sicyopterus pugnans*, *S. longifilis*, *S. cynocephalus*, *S. lagocephalus*, *S. parvei*, *Belobranchus segura*, *B. belobranchus*, *Bunaka gyrinoides*, and *Stiphodon semoni* (Sahami et al 2019b; Sahami et al 2020). The high diversity of species of a composer for Nike fish schools indicates the potential high diversity of goby species in the Bone Bolango River.

Based on its morphological characters, gobies in the Bone Bolango River generally have a torpedo (fusiform) body shape. They have two separate dorsal fins, pectoral fins, pelvic fins, anal fin, and rounded caudal fin. In general, the goby species can be grouped into three large groups known locally as Hundala, Busalo, and Timudu'o. The special characteristics that distinguish between Hundala and Busalo fish are the pelvic fins of Hundala fish are separated and not connected by a membrane and the pelvic fins of Busalo fish are combined to form something like suction discs. In contrast, Timudu'o fish are identified as small gobies found in river waters and have pelvic fins that joined like Busalo fish but with a different morphological appearance.

After editing and cutting sequences, 651 bp of mitochondrial DNA COI were obtained for all the goby samples. A total of 14 groups of goby species were confirmed as adult phases of Nike fish based on the results of COI mitochondrial DNA BLAST. Five of which were Busalo (B1-B5), seven groups were Hundala fish species (H1-H7), and two groups were Timudu'o fish species (T1-T2) as shown in Table 1.

Table 1 shows goby in the Bone Bolango River consisting of four genera and six species; *Sicyopterus longifilis*, *S. lagocephalus*, *Belobranchus belobranchus*, *B. segura*, *Bunaka gyrinoides*, and *Stiphodon semoni*. This species' identification results indicate that the Hundala fish are part of the Eleotridae family, and Busalo and Timudu'o fish are the fish group of the Gobiidae family. The same genetic identity in two or more different morphological groups can be influenced by the environment and can also occur due to sexual dichromatism. Sexual dichromatism usually occurs in freshwater Gobioid fish, in which males have a brighter color than females (Keith 2003). *S. semoni* is one species that shows the phenomenon of sexual dichromatism in which males have a blackish gray body color with greenish staining on the head to the chest area no longer after being captured (T1). On the other hand, females tend to have a yellowish-brown color, and they have transverse lines from the tip of snout to the base of the caudal fin (T2) as described by Maeda & Tan (2013).

Table 1

BLAST results of the mitochondrial COI DNA of adult phases of species of composers for Nike schools

No	Sample codes	Species	NCBI accession number	Query cover (%)	Per. identity (%)
1	B1	<i>Sicyopterus longifilis</i>	MK496959.1	100	99.69
2	B2	<i>Sicyopterus lagocephalus</i>	KF668859.1	100	99.39
3	B3	<i>Sicyopterus lagocephalus</i>	KF668859.1	100	99.54
4	B4	<i>Sicyopterus longifilis</i>	NC_044142.1	99	99.56
5	B5	<i>Sicyopterus longifilis</i>	NC_044142.1	99	99.71
6	H1	<i>Belobranchus belobranchus</i>	KU692346.1	99	99.39
7	H2	<i>Belobranchus segura</i>	MN069308.1	99	99.69
8	H3	<i>Belobranchus segura</i>	MN069308.1	99	99.69
9	H4	<i>Belobranchus belobranchus</i>	MT706791.1	99	99.08
10	H5	<i>Belobranchus belobranchus</i>	MT706791.1	99	99.23
11	H6	<i>Belobranchus segura</i>	MN069308.1	99	99.69
12	H7	<i>Bunaka gyrinoides</i>	MN069307.1	100	99.85
13	T1	<i>Stiphodon semoni</i>	MT706724.1	95	99.70
14	T2	<i>Stiphodon semoni</i>	MT706724.1	92	99.70

Other species that also show the phenomenon of sexual dichromatism are *S. longifilis* and *S. lagocephalus*. *S. longifilis* males (B1) are green with particular features. They have vertical stripes that sometimes extend to the abdomen, the black pectoral fins with white ribbons on edge, and slightly orange with red and blue lines at the dorsal and ventral tips of the caudal fin. On another side, the female fish are brown and duller (B4 and B5). At the same time, the similar condition is also shown by *S. lagocephalus* species, in which male fish has a very attractive color that is bluish to greenish with a reddish-orange caudal fin (B3). In contrast, the female fish are brownish to grayish and have an orange ribbon on the edge of the pectoral fin, anal fin, and the dorsal and ventral sides of the caudal fin (B2).

Sexual dichromatism is not found in *B. segura* and *B. belobranchus* species, as reported by Keith et al (2012). Both of these species are goby sleepers with the elongated body with anterior parts forming cylindrical and posterior compressed, and having brownish to mahogany body colors. The distinctive characteristic between them is the presence of darker and thicker color that extends from chest to the base of caudal fin along the lateral line in *B. segura* species. Whereas the *B. gyrinoides* species found in one morphological characteristic in this study. This species is also goby sleepers with a typical large body, a similar body shape with *Belobranchus* species, a blackish-brown body color, and having fins with blackish-brown spots and blackish-brown vertical stripes on the caudal fin (Huang & Chen 2008). The relationships between adult species of Nike fish and the species of a composer for Nike fish schools are illustrated through the phylogenetic tree in Figure 2.

The phylogenetic tree (Figure 2) clearly shows the kinship between the species of a composer for Nike fish schools and goby fish which have been reported and found in this study. The whole phylogenetic tree forms two monophyletic clades as family clades. The first monophyletic clade is the Gobiidae Family clade consisting of *S. lagocephalus*, *S. longifilis*, and *Stiphodon semoni*. The second monophyletic clade belongs to Eleotridae family which includes *B. gyrinoides*, *B. segura* and *B. belobranchus*. From nine species of a composer for Nike fish schools, three of them (*S. cynocephalus*, *S. pugnans*, and *S. parvef*) have not been found yet in the adult phase in this study. However, the genetic distance between species found in Bone Bolango River shown in Table 2 will strengthen the results of the phylogenetic tree formed in Figure 2.

Eleotridae is one of the families in the Gobioidi Order known as a sleeper species with prey-waiting characteristics. They have a large blunted head, a torpedo body shape, rounded caudal fins, protruding mandible, protruding pelvic fins that are not connected by membranes, and not bright body color, namely light brown to dark brown (Murdy & Hoese 2002). Eleotrid species usually prefer water habitats where have small flow and they usually stay in river banks. These morphological characteristics have been found in Hundala fish and confirmed through its genetic analysis as species of Eleotridae Family.

Unlike the Eleotridae fish, the species in Gobiidae Family are mostly active swimmers and are very fond of currents. In Reunion Island, adult species of *S. lagocephalus* are known to be rheophilic and like clear habitats with good oxygen levels (Keith 2003). Gobiidae is widely distributed in the sea, estuary, and freshwater. Gobiid species generally have pelvic fins that combined to form something like sucking discs, both partially and wholly. This modification allows quick access to the river's headwaters by attaching itself to the substrate when the species is against the current (Taillebois et al 2014). Also, this species' body color varies greatly from the drab to various bright patterns (Murdy & Hoese 2002; Thacker & Roje 2011). The morphological characteristics are found in Busola and Timudu'o fish, which also confirmed through genetic analysis as species in Gobiidae Family.

Therefore, it has been genetically confirmed that six goby species in the adult phase are the species of a composer for Nike fish schools in the Gorontalo Bay waters. *S. longifilis* species with its female morphological characteristics have been reported by Olii et al (2019) as the adult stage of Nike fish. Besides, *B. belobranthus* species with its morphological features of H1 has also been found by Pasisingi et al (2020a) and confirmed as an adult phase of Nike fish through this study. The discovery of the adult phase of Nike fish has been new and important information for the local community in Gorontalo so that they can actively contribute to the availability of Nike fish resources sustainably in the future. It can be carried out by not overfishing gobies or releasing gobies, which carrying eggs into the river. This discovery also refutes the local community's opinions regarding Nike fish only live in sea waters, they cannot increase their size, and they tend to disappear on the surface.




Figure 2. Phylogenetic tree of adult fish goby and species of a composer for Nike fish schools in the Gorontalo Bay waters.

Table 2

The genetic distance of goby adult phases of species of composers for Nike fish schools

	<i>B1</i>	<i>B2</i>	<i>B3</i>	<i>B4</i>	<i>B5</i>	<i>H1</i>	<i>H2</i>	<i>H3</i>	<i>H4</i>	<i>H5</i>	<i>H6</i>	<i>H7</i>	<i>T1</i>	<i>T2</i>
B1														
B2	0.10													
B3	0.10	0.00												
B4	0.01	0.09	0.10											
B5	0.01	0.09	0.09	0.00										
H1	0.22	0.20	0.21	0.22	0.22									
H2	0.24	0.23	0.23	0.24	0.24	0.13								
H3	0.25	0.23	0.23	0.24	0.24	0.13	0.00							
H4	0.23	0.21	0.21	0.22	0.22	0.01	0.14	0.14						
H5	0.22	0.20	0.21	0.22	0.22	0.01	0.13	0.13	0.00					
H6	0.25	0.23	0.23	0.24	0.24	0.13	0.00	0.00	0.14	0.13				
H7	0.20	0.22	0.22	0.20	0.20	0.17	0.19	0.19	0.17	0.17	0.19			
T1	0.12	0.12	0.12	0.12	0.12	0.21	0.22	0.23	0.21	0.21	0.23	0.20		
T2	0.12	0.12	0.12	0.12	0.12	0.21	0.23	0.23	0.22	0.21	0.23	0.19	0.01	

Conclusions. This research has confirmed that Nike is an amphidromous species which supported by the discovery of the adult phase of six species composers of Nike fish assemblages; *Sicyopterus longifilis*, *S. lagocephalus*, *Belobranchus belobranchus*, *B. segura*, *Bunaka gyrinoides*, and *Stiphodon semoni* in Bone Bolango River based on the DNA COI mitochondria analysis. The sustainability of Nike's resources can be done with proper management of the adult phase of Nike fish in the river, such as not overfishing goby or releasing goby that carries eggs into the river.

Acknowledgements. The authors express their gratitude to the Institute of Research and Community Service (LPPM) Gorontalo State University for funding this research through the 2020 Budget on Penelitian Dasar scheme. The authors also thank Roy Bakari and Thomas Tammu for the technical assistance during the study and all those who contributed to this research.

References

- Ellien C., Werner U., Keith P., 2014 Morphological changes during the transition from freshwater to sea water in an amphidromous goby, *Sicyopterus lagocephalus* (Pallas 1770) (Teleostei). *Ecology of Freshwater Fish* 25(1):48-59.
- Hebert P. D. N., Stoeckle M. Y., Zemplak T. S., Francis C. M., 2004 Identification of birds through DNA barcodes. *PLoS Biology* 2(10):e312.
- Huang J. H., Chen I. S., 2008 A new record of the freshwater gobioid genus, *Bunaka* Herre, 1927 (Pisces: Eleotridae) from Taiwan. *Taiwan Journal of Forest Science* 23(2):183-189.
- Hubert N., Hanner R., Holm E., Mandrak N. E., Taylor E., Burrige M., Watkinson D., Dumont P., Curry A., Bentzen P., Zhang J., April J., Bernatchez L., 2008 Identifying Canadian freshwater fishes through DNA barcodes. *PLoS One* 3(6):e2490.
- Iida M., Kondo M., Tabouret H., Maeda K., Pécheyran C., Hagiwara A., Keith P., Tachihara K., 2017 Specific gravity and migratory patterns of amphidromous gobioid fish from Okinawa Island, Japan. *Journal of Experimental Marine Biology and Ecology* 486:160-169.
- Ivanova N. V., Zemplak T. S., Hanner R. H., Hebert P. D. N., 2007 Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Notes* 7(4):544-548.
- Keith P., 2003 Biology and ecology of amphidromous Gobiidae of the Indo-Pacific and the Caribbean regions. *Journal of Fish Biology* 63(4):831-847.
- Keith P., Hoareau T. B., Lord C., Ah-Yane O., Gimonneau G., Robinet T., Valade P., 2008 Characterisation of post-larval to juvenile stages, metamorphosis and recruitment of an amphidromous goby, *Sicyopterus lagocephalus* (Pallas) (Teleostei: Gobiidae: Sicydiinae). *Marine and Freshwater Research* 59(10):876-889.
- Keith P., Hadiaty R. K., Lord C., 2012 A new species of *Belobranchus* (Teleostei: Gobioidae: Eleotridae) from Indonesia. *Cybio* 36(3):479-484.
- Kumar S., Stecher G., Li M., Knyaz C., Tamura K., 2018 MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35(6):1547-1549.
- Lejeune L., Tabouret H., Taillebois L., Monti D., Keith P., 2016 Larval traits of the Caribbean amphidromous goby *Sicydium punctatum* (Gobioidae: Sicydiinae) in Guadeloupe. *Ecology of Freshwater Fish* 25(2):272-280.
- Linh N. M., Thu P. T., Quan N. V., Chien P. V., Ly D. H., Nhan D. V., Len D. T., 2018 DNA barcoding application of mitochondrial COI gene to identify some fish species of family Gobiidae in Vietnam. *Vietnam Journal of Marine Science and Technology* 18(4):433-451.
- Maeda K., Tan H. H., 2013 Review of *Stiphodon* (Gobiidae: Sicydiinae) from Western Sumatra, with description of a new species. *Raffles Bulletin of Zoology* 61(2):749-761.
- Murdy E. O., Hoesel D. F., 2002 Suborder Gobioidae. In: *The living marine resources of the Western Central Atlantic*. Vol. 3: Bony fishes part 2 (Opistognathidae To Molidae), sea turtles and marine mammals. Carpenter K. E. (ed), FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Special Publication No.5, FAO, Rome, pp. 1778-1798.

- Nurjirana, Haris A., Sahami F. M., Keith P., Burhanuddin A. I., 2019 Preliminary note on the morphological characters of *penja* (amphidromous goby post larvae) in West Sulawesi and Gorontalo Bay. IOP Conference Series: Earth and Environmental Science 370(1):012007.
- Olii A. H., Sahami F. M., Hamzah S. N., Pasingi N., 2017 Preliminary findings on distribution pattern of larvae of Nike fish (*Awaous* sp.) in the estuary of Bone River, Gorontalo Province, Indonesia. AACL Bioflux 10(5):1110-1118.
- Olii A. H., Sahami F. M., Hamzah S. N., Pasingi N., 2019 Molecular approach to identify gobioid fishes, "Nike" and "Hundala" (local name), from Gorontalo waters, Indonesia. Online Journal of Biological Sciences 19(1):51-56.
- Pasingi N., Abdullah S., 2018 [Pattern of Nike fish (Gobiidae) occurrence in the Gorontalo Bay, Indonesia]. Depik 7(2):111-118. [in Indonesian]
- Pasingi N., Habibie S. A., Olii A. H., 2020a Are *Awaous ocellaris* and *Belobranchus belobranchus* the two species of Nike fish schools? Aceh Journal of Animal Science 5(2):87-91.
- Pasingi N., Olii A. H., Habibie S. A., 2020b Morphology and growth pattern of Nike fish (amphidromous goby larvae) in Gorontalo Waters, Indonesia. Tomini Journal of Aquatic Science 1(1):1-7.
- Sahami F. M., Kepel R. C., Olii A. H., Pratasik S. B., 2019a Determination of morphological alteration based on molecular analysis and melanophore pattern of the migrating Nike fish in Gorontalo Bay, Indonesia. AACL Bioflux 12(4):1358–1365.
- Sahami F. M., Kepel R. C., Olii A. H., Pratasik S. B., 2019b What species make up the Nike fish assemblages at the macrotidal estuary in Gorontalo Bay, Indonesia? [version 1; peer review: 2 approved with reservation]. F1000Research 8:1654.
- Sahami F. M., Kepel R. C., Olii A. H., Pratasik S. B., Lasabuda R., Wantasen A., Habibie S. A., 2020 Morphometric and genetic variations of species composers of Nike fish assemblages in Gorontalo Bay Waters, Indonesia. Biodiversitas 21(10):4571-4581.
- Strüder-Kypke M., Lynn D. H., 2010 Comparative analysis of the mitochondrial cytochrome *c* oxidase subunit I (COI) gene in ciliates (Alveolata, Ciliophora) and evaluation of its suitability as a biodiversity marker. Systematics and Biodiversity 8(1):131–148.
- Taillebois L., Maeda K., Vigne S., Keith P., 2012 Pelagic larval duration of three amphidromous Sicydiinae gobies (Teleostei: Gobioidei) including widespread and endemic species. Ecology of Freshwater Fish 21(4):552-559.
- Taillebois L., Castelin M., Lord C., Chabarría R., Dettai A., Keith P., 2014 New *sicydiinae* phylogeny (Teleostei: Gobioidei) inferred from mitochondrial and nuclear genes: insights on systematics and ancestral areas. Molecular Phylogenetics and Evolution 70:260-271.
- Thacker C. E., Roje D. M., 2011 Phylogeny of Gobiidae and identification of gobiid lineages. Systematics and Biodiversity 9(4):329-347.
- Usman M. Y., 2016 [Analysis of genetic variation *penja* indigenous from Polewali Mandar and Nike (*Awaous* sp.) indigenous from Gorontalo]. Undergraduate Thesis, Makassar: Fakultas Sains dan Teknologi, UIN Alauddin Makassar, Makassar, 71 pp. [in Indonesian]
- Viswambharan D., Pavan-Kumar A., Singh D. P., Jaiswar A. K., Chakraborty S. K., Nair J. R., Lakra W. S., 2013 DNA barcoding of gobiid fishes (Perciformes, Gobioidei). Mitochondrial DNA 26(1):15-19.
- Ward R. D., Zemlak T. S., Innes B. H., Last P. R., Hebert P. D. N., 2005 DNA barcoding Australia's fish species. Philosophical Transactions of the Royal Society B - Biological Sciences 360(1462):1847-1857.
- Yamasaki N., Kondo M., Maeda K., Tachihara K., 2011 Reproductive biology of three amphidromous gobies, *Sicyopterus japonicus*, *Awaous melanocephalus*, and *Stenogobius* sp., on Okinawa Island. Cybium 35(4):345-359.

Received: 08 August 2020. Accepted: 05 October 2020. Published online: 26 October 2020.

Authors:

Femy M. Sahami, Faculty of Fisheries and Marine Sciences, Gorontalo State University, Jl. Jendral Sudirman No. 6, 96128, Gorontalo, Indonesia, e-mail: femysahami@ung.ac.id

Sitty Ainsyah Habibie, Faculty of Fisheries and Marine Sciences, Gorontalo State University, Jl. Jendral Sudirman No. 6, 96128, Gorontalo, Indonesia, e-mail: ainsyahabibie@ung.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Sahami F. M., Habibie S. A., 2020 Exploration of adult phase of Nike fish to maintain its sustainability in Gorontalo Bay waters, Indonesia. *AACL Bioflux* 13(5): 2859-2867.