



Digital Receipt

This receipt acknowledges that **Turnitin** received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Margaretha Solang
Assignment title: Cek 1
Submission title: Anadara granosa substitution in fee...
File name: 17__fr-2020-458_solang.pdf
File size: 353.28K
Page count: 8
Word count: 5,976
Character count: 31,315
Submission date: 25-Mar-2021 07:33PM (UTC+1100)
Submission ID: 1541902456

Food Research 5 (1) : 132 - 139 (February 2021)
Journal homepage: <http://www.myfoodresearch.com>

FOOD
RESEARCH

FULL PAPER

Anadara granosa substitution in feed to improve the zinc, protein of the feed, serum albumin, and body weight of malnourished rats

^{1,2}Solang, M. and ²Adriani, M.

¹State University of Gorontalo, Faculty of Mathematics and Natural Sciences, Department of Biology, Jl. Jenderal Sudirman No. 6 Kota Gorontalo, 96128, Gorontalo, Indonesia
²Airlangga University, Faculty of Public Health, Health Science Department, Campus C, Mulyorejo, Kec. Mulyorejo, Surabaya, 6011, Jawa Timur, Indonesia

Article history:
Received: 21 August 2020
Received in revised form: 7 September 2020
Accepted: 21 November 2020
Available Online: 1 December 2020

Keywords:
Anadara granosa,
Feed,
Protein,
Growth,
Malnourished

DOI:
<https://doi.org/10.24055/foodresearch.2021.5.1.148>

Abstract
The objective of this paper was to evaluate zinc and protein feed levels of *Anadara granosa* flour substitution and its effect of feed on serum albumin and body weight of a malnourished rat (*Rattus norvegicus*). This study comprised two stages: 1) analysis of protein and zinc level of the feed, and 2) feed test on rat using separate sample pre-test and post-test control group design. The malnourished rat was treated with dried rice; the dried rice was substituted with 12.5%, 25%, and 50% of *Anadara granosa* flour. Further, at room temperature, the flour can be kept for six months. This study revealed that the *Anadara granosa*-substituted feed had zinc levels between 0.999 ppm - 2.296 ppm and protein levels of 14.81% - 26.39%. On the other hand, the non-substituted feed had 0.791 ppm of zinc, and the protein level was 8.46%. Provision of the feed substituted with *Anadara granosa* flour increased the albumin serum level ($p = 0.000$) and the bodyweight of malnourished rats significantly ($p = 0.002$). This study revealed that substitution of *Anadara granosa* flour in feed could improve the zinc and protein level of the feed, which in turn improved the growth of malnourished rats (as the albumin level and the bodyweight also improved).

1. Introduction
Malnutrition, especially zinc deficit, is largely found on children in developing countries; in childhood stage, children need a higher zinc intake (Yanagisawa, 2004; Parveen and Dipati, 2016). Zinc deficit would influence homeostasis within the biological system (Kaur *et al.*, 2016). A nutrition deficit condition that occurs for a long time might lead to stunting (Chirande *et al.*, 2015; Mardewi *et al.*, 2016). Stunting is one of the prominent health problems in Indonesia as its prevalence is still currently above 20% (Kemenkes, 2018). Provision of zinc supplements is proven to be able to treat stunting (Kusudaryati *et al.*, 2017) since children with stunting have a low level of zinc and albumin serum levels. Thus, the administration of Vitamin A with zinc supplements could reduce infection risk and improve the linear growth of children with stunting (Adriani and Wirjatmadi, 2014).
Gibson and Fergusson (1994) reported that zinc deficiency could be solved by increasing zinc intake. As what has been done in Africa, increasing zinc intake through zinc-rich meats and meals that are economically and culturally acceptable in that area, the deficiency problem could be addressed. Zinc is commonly available in high protein food. Animal-based meals are the main source of zinc in the human diet (Agustian *et al.*, 2009; Kaur *et al.*, 2016). Zinc is one of the most essential elements in nutrition for humans and animals. In the growth process, zinc helps in protein synthesis to form new cells, growth, and bone development (Agustian *et al.*, 2009). Zinc also has several physiological characteristics and activates enzymes within the body (Kaur *et al.*, 2016).
Anadara granosa is one of the seafood that is rich in zinc contents. Fresh *Anadara granosa* contain 19.48% of protein and 13.91 ppm of zinc (Nurjanah *et al.*, 2005). It also contains complex amino acids. The protein contents in the *Anadara granosa* would help absorb zinc and increase protein intake to the body. Therefore, zinc and protein in the *Anadara granosa* will work in synergy to improve zinc levels of malnourished rats (Solang *et al.*, 2013). On top of that, zinc also plays a role in inducing metallothionein (zinc-binding protein); thus, it regulates

*Corresponding author.
Email: margarethasolang@gmail.com

eISSN: 2550-2166 / © 2020 The Authors. Published by Rynnyss Lynn Resources

Anadara granosa substitution in feed to improve the zinc, protein of the feed, serum albumin, and body weight of malnourished rats

by Margaretha Solang

Submission date: 25-Mar-2021 07:33PM (UTC+1100)

Submission ID: 1541902456

File name: 17__fr-2020-458_solang.pdf (353.28K)

Word count: 5976

Character count: 31315

1

Anadara granosa substitution in feed to improve the zinc, protein of the feed, serum albumin, and body weight of malnourished rats^{1,*}Solang, M. and ²Adriani, M.

2

¹State University of Gorontalo, Faculty of Mathematics and Natural Sciences, Department of Biology, Jl. Jendral Sudirman No. 6 Kota Gorontalo, 96128, Gorontalo, Indonesia²Airlangga University, Faculty of Public Health, Health Science Department, Campus C, Mulyorejo, Kec. Mulyorejo, Surabaya, 6011, Jawa Timur, Indonesia**Article history:**

Received: 21 August 2020

Received in revised form: 7

September 2020

Accepted: 21 November 2020

Available Online: 5

December 2020

Keywords:

Anadara granosa,

Feed,

Protein,

Growth,

Malnourished

DOI:[https://doi.org/10.26656/fr.2017.5\(1\).458](https://doi.org/10.26656/fr.2017.5(1).458)**Abstract**

1

The objective of this paper was to evaluate zinc and protein feed levels of *Anadara granosa* flour substitution and its effect of feed on serum albumin and body weight of a malnourished rat (*Rattus norvegicus*). This study comprised two stages: 1) analysis of protein and zinc level of the feed, and 2) feed test on rat using *separate sample pre-test and post-test control group design*. The malnourished rat was treated with dried rice; the dried rice was substituted with 12.5%, 25%, and 50% of *Anadara granosa* flour. Further, at room temperature, the flour can be kept for six months. This study revealed that the *Anadara granosa*-substituted feed had zinc levels between 0.999 ppm - 2.296 ppm and protein levels of 14.81% - 26.39%. On the other hand, the non-substituted feed had 0.791 ppm of zinc, and the protein level was 8.46%. Provision of the feed substituted with *Anadara granosa* flour increased the albumin serum level ($p = 0.000$) and the bodyweight of malnourished rats significantly ($p = 0.002$). This study revealed that substitution of *Anadara granosa* flour in feed could improve the zinc and protein level of the feed, which in turn improved the growth of malnourished rats (as the albumin level and the bodyweight also improved).

1. Introduction

Malnutrition, especially zinc deficit, is largely found on children in developing countries; in childhood stage, children need a higher zinc intake (Yanagisawa, 2004; Parveen and Dipti, 2016). Zinc deficit would influence homeostasis within the biological system (Kaur *et al.*, 2016). A nutrition deficit condition that occurs for a long time might lead to *stunting* (Chirande *et al.*, 2015; Mardewi *et al.*, 2016). *Stunting* is one of the prominent health problems in Indonesia as its prevalence is still currently above 20% (Kemenkes, 2018). Provision of zinc supplements is proven to be able to treat *stunting* (Kusudaryati *et al.*, 2017) since children with *stunting* have a low level of zinc and albumin serum levels. Thus, the administration of Vitamin A with zinc supplements could reduce infection risk and improve the linear growth of children with *stunting* (Adriani and Wirjatmadi, 2014).

Gibson and Ferguson (1994) reported that zinc deficiency could be solved by increasing zinc intake. As what has been done in Africa, increasing zinc intake

through zinc-rich meats and meals that are economically and culturally acceptable in that area, the deficiency problem could be addressed. Zinc is commonly available in high protein food. Animal-based meals are the main source of zinc in the human diet (Agustian *et al.*, 2009; Kaur *et al.*, 2016). Zinc is one of the most essential elements in nutrition for humans and animals. In the growth process, zinc helps in protein synthesis to form new cells, growth, and bone development (Agustian *et al.*, 2009). Zinc also has several physiological characteristics and activates enzymes within the body (Kaur *et al.*, 2016).

Anadara granosa is one of the seafood that is rich in zinc contents. Fresh *Anadara granosa* contain 19.48% of protein and 13.91 ppm of zinc (Nurjanah *et al.*, 2005). It also contains complex amino acids. The protein contents in the *Anadara granosa* would help absorb zinc and increase protein intake to the body. Therefore, zinc and protein in the *Anadara granosa* will work in synergy to improve zinc levels of malnutrition rats (Solang *et al.*, 2013). On top of that, zinc also plays a role in inducing metallothionein (zinc-binding protein); thus, it regulates

*Corresponding author.

Email: margarethasolang9@gmail.com

the amino acid as the precursor for the synthesis of albumin (Tekeli, 2002). The level of albumin plasma could be used as a sensitive indicator of nutrition status that reflects a diet pattern (Kuwahata *et al.*, 2017). The previous studies about ¹ serves as the rationale, as the present study aims to evaluate zinc and protein in *Anadara* ¹ *granosa* flour-substituted feed and the feed's effect on serum albumin levels and body weight of a malnourished rat (*Rattus norvegicus*). This study hypothesized that *Anadara granosa*-substituted feeds can improve zinc and protein feed levels, and the consumption of the feeds can improve albumin levels and body weight of malnourished rat.

2. Material and methods

2.1 Research design and sample

The *Anadara granosa* used as substitute feeds in this study were taken from Gorontalo province, Indonesia. Rats were fed in the form of pellets consisting of standard feed, dried rice, and *Anadara granosa* flour substitution. The standard pellet feed produced by PT Charoen Pokpan contain 13% water, 13-15% protein, 3% fat, 6% ash, and 0.8% calcium. The dried rice feed was used to create malnutrition conditions. The dried rice was made of leftover rice. The rice that will no longer be consumed was set out to dry under the sun. After adding flour, the rice is cooked and stirred until mixed well. The dough is poured into a grinder and was shaped into a pellet. Further, the pellet was set to dry under the sun for the second time.

Further, the processes of making the flour are as follows. First, the whole cockles were boiled until they open. Then, the soft tissues of the cockles were removed. After that, the tissues were cut into small pieces. All the cut tissues were dried in the sun to get a constant dry weight. The tissues, after the sun-drying process, were finely milled and sifted to get the cockle flour. *Anadara granosa* flour contained ¹ 27.26% of protein and 0.7913 ppm of zinc and the flour can be kept at room temperature for six months.

² This study was designed on testing rats using the *Separate Sample Pre-Post Test Control Group Design* (Campbell and Stanley, 1963; Handley *et al.*, 2018). On the first test, 12 male Wistar rats aged 6 weeks with 115-120 g bodyweight were given standard feed (normal group). Meanwhile, 36 male Wistar rats aged 6 weeks with a bodyweight of 110-120 g were treated using dried rice pellet feed with 8% of protein (nutrition deficit group). In the 8th weeks, 4 rats from the normal control group and 4 rats from the nutrition deficit group were slaughtered to test their albumin level. The rat with albumin serum less than 3.3 mg/dL was regarded to have

nutrition deficit (Giknis and Clifford, 2008; Susanto *et al.*, 2010). In the second phase, normal control rats (8 rats) were treated with standard pellet feed. Thirty-two malnutrition rat were randomly distributed into 4 groups; 8 rats were fed with dried rice, 8 rats were fed with 12.5% *Anadara granosa* -substituted dried rice (*Anadara granosa* flour 12.5% + 87.5% dried rice), 8 rats were fed with 25% *Anadara granosa* -substituted dried rice (*Anadara granosa* flour 25% + 75% dried rice), and 8 rats were fed with 50% *Anadara granosa* -substituted dried rice (*Anadara granosa* flour 50% + 50% dried rice) for 8 weeks. The percentage of *Anadara granosa* substitution in dried rice is based on the zinc needs of children per day, which is 10 g (Kartono *et al.*, 2012), then converted to rat weighing 200 g. Food and drink for rats were given through *ad libitum*.

2.2 Ethics test

This study has obtained the ethical certificate No: 11-KEPK from the Faculty of Public Health, Universitas Airlangga. The implementation methods were presented in front of the ethical committee and have obtained its approval.

2.3 Procedure

Proximate analysis of dried rice feed and feed substituted with *Anadara granosa* flour were carried out at the Laboratory Unit of Veterinary Faculty of Universitas Airlangga. Acclimatization, cultivation, and surgery of the test animals were carried out at the Veterinary Laboratory of Biochemical Department of Medical Faculty of Universitas Airlangga. Protein content test was carried out with the micro Kjeldahl method (AOAC, 1995). The measurement of protein content in the blood cockle flour was conducted at the Animal Feed Laboratory, Faculty of Veterinary Medicine, Universitas Airlangga. Whereas the Soxhlet method was employed to test the fat level (AOAC, 1995). In addition, the zinc level of the feed was measured using Atomic Absorbant Spectrophotometer (AAS) with the Zenit 700 tool. Further, the analysis of the zinc content of the flour was carried out at the Center for Health Laboratory, Surabaya.

As much as 3 mL of a blood sample from the tested rats ² were taken from the heart and stored within the *Blood collection tube* EDTA. To obtain the blood serum, the speed was set into 3000 rpm for 15 minutes centrifuged the blood sample. The serum was then separated into Eppendorf ² bes. Further, the Albumin level was tested by automatic chemical analysis *Prestige 24i*. Cat. No. 4-238 with *Bromocresol Green* (BCG) method (Dumas and Peters, 2009). The

albumin level was stated in g/dL. The bodyweight of the tested rats was measured using a digital scale of Camry brand with the maximum capacity of 500 g, division 0.1 g with the accuracy level of two numbers behind the coma.

2.4 Statistical analysis

The zinc and protein levels were analyzed in descriptive manner. Meanwhile, the serum albumin level of the tested rat was analyzed using One way ANOVA parametric test in the significance level of 95% and the least significant difference (LSD). Further, the body weight was analyzed using the *Kruskal – Wallis* test on the significance level of 95% and followed by an advance test of Mann Whitney (Steel and Torrie, 1980; McDonald, 2014).

3. Results

3.1 Zinc and Protein level of *Anadara granosa* flour-substituted feed

The analysis of the dried rice feed used to create malnutrition conditions and *Anadara granosa* substituted feed showed a different protein and zinc levels. The dried feed rice had a protein level of 8.462% and zinc 0.7913 ppm. The *Anadara granosa* flour had a protein level of 14.81% and zinc 0.9995 ppm. Meanwhile, dried rice feed substituted with 25% of *Anadara granosa* flour had a protein level of 18.74% and zinc 1.151 ppm. In addition, the 50% substitution of *Anadara granosa* flour had a protein level of 26.394% and zinc 2.296 ppm (Table 1). The level of zinc in *Anadara granosa*'s flour-substituted feed increased along with the increase of *Anadara granosa* composition within the feed (Table 2). The increase of zinc level percentage on feed substituted with 12.5%, 25%, and 50 *Anadara granosa*'s feed-in the sequence were 26.31%, 45.46%, and 58.35% respectively. Meanwhile, the increase of protein level of the feed substituted with 12.5%, 25%, and 50% of *Anadara granosa*'s flour were 75%, 121%, and 211% in consecutive order. This showed that the substitution of *Anadara granosa*'s flour could increase the zinc and protein level of the feed.

Table 1. Zinc and protein level of dried rice feed and *Anadara granosa* flour-substituted dried rice feed

Types of feed	Zinc level (ppm)	Rough protein level (%)
Dried rice	0.791	8.462
Dried rice with 12.5% <i>Anadara granosa</i> 's flour substitution	0.999	14.81
Dried rice with 25% <i>Anadara granosa</i> 's flour substitution	1.151	18.74
Dried rice with 50% <i>Anadara granosa</i> 's flour substitution	2.296	26.39

3.2 Albumin level

The results revealed that the *Anadara granosa* flour substitution feed significantly increased the albumin level of the malnutrition rats ($p = 0.000$) (Table 3). The average albumin level of malnutrition rats was 16.16% lower than a normal rat (rat fed with standard feed). In the present work, the percentage was employed to describe the decline in the albumin content due to the differences in the smaller, last number (the level of low-nutrient albumin) and the initial, greater number (the normal group of albumin level). On the other hand, the use of the g/dL unit in measuring the decline and increase in the albumin level will only hamper the process of predicting the increase or decline of the albumin content. Moreover, this percentage also applies to the increase in the albumin level; it takes into account the differences in the recent level of the albumin content of the poor-nourished mice (that had been provided with blood cockle flour) and the level of albumin in the normal group mice). The poor-nourished mice have a higher albumin level than the normal group. Another point worth considering is that the percentage provides more significant data (representing the level of albumin content).

The average albumin level of nutrition deficit rat who were treated with *Anadara granosa* flour substituted feed increased along with the increase of *Anadara granosa*'s composition within the feed. The albumin level of the malnutrition rat treated with 25% and 50% of *Anadara granosa*'s substituted feed similarly experienced an increase of 22.59%. The substitution of

Table 2. Percentage of zinc and protein increase in the *Anadara granosa* flour-substituted feed

No.	Sample	Zinc level (ppm)	Zinc level increase (%)	Rough protein level (%)	Increase of rough protein level (%)
1	Dried rice (kg)	0.791	0.00	8.46	0.00
2	Dried rice feed substituted with 12.5% of <i>Anadara granosa</i> 's flour	0.999	26.31	14.81	75
3	Dried rice feed substituted with 25% of <i>Anadara granosa</i> 's flour	1.151	45.46	18.74	121
4	Dried rice feed substituted with 50% of <i>Anadara granosa</i> 's flour	1.253	58.35	26.39	211

Table 3. The average level of albumin (g/dL) level of malnutrition rat fed with *Anadara granosa*'s flour-substituted feed.

No.	Treatment	Average albumin level (g/dL)	The decrease in albumin level (%)	Increase of albumin level (%)	P-value
1	Rat treated with standard feed	3.59±0.27 ^a	0	0	0
2	Rat treated with dried rice feed	3.01±0.13 ^b	16.16	-	
3	Rat treated with dried rice substituted with 12.5% <i>Anadara granosa</i> 's flour	3.64±0.15 ^a	-	20.93	
4	Rat treated with dried rice substituted with 25% <i>Anadara granosa</i> 's flour	3.69±0.22 ^a	-	22.59	
5	Rat treated with dried rice substituted with 50% <i>Anadara granosa</i> 's flour	3.69±0.42 ^a	-	22.59	

Notes: Average value followed by different superscript letter showed significance difference

Table 4. Average body weight (g) of malnutrition rat treated with *Anadara granosa*'s flour- substituted feed.

No.	Treatment	Average body weight (g)	The decrease in body weight (%)	Increase of body weight (%)	P-value
1	Rat treated with standard feed	257.43±19.28 ^a	0	-	0.002
2	Rat treated with dried rice feed	112.57±21.94 ^b	56.27	-	
3	Rat treated with dried rice substituted with 12.5% <i>Anadara granosa</i> 's flour	215.43±15.38 ^c	-	47.75	
4	Rat treated with dried rice substituted with 25% <i>Anadara granosa</i> 's flour	225.86± 19.96 ^{cd}	-	50.16	
5	Rat treated with dried rice substituted with 50% <i>Anadara granosa</i> 's flour	236.29±10.63 ^d	-	50.16	

Notes: Average value followed by different superscript letter showed significance difference

12.5% of *Anadara granosa*'s flour had improved the albumin level of malnutrition rat into the normal level.

3.3 Bodyweight

The result showed that the malnutrition rats fed with *Anadara granosa* flour substitution experienced a significant increase in body weight ($p = 0.002$) (Table 4). The average bodyweight of the malnutrition rats was 56.27% lower than normal rat treated with standard feed. Further, the average body weight of malnutrition rat treated with 25% and 50% of *Anadara granosa* flour substitution increased by 50.16%.

Moreover, the bodyweight of malnutrition rat treated with feed substituted with 50% *Anadara granosa*'s flour was yet to achieve the maximum result, as indicated by the rats' below normal bodyweight.

4. Discussion

4.1 The level of zinc and protein in the *Anadara granosa* flour-substituted feed

The feed substituted with *Anadara granosa* flour had a higher zinc content due to the natural zinc. Similar studies by Solang *et al.* (2017) showed that the zinc level in *cireng* snack made with flour substituted with *Anadara granosa* flour increased. The increase of zinc in food made with *Anadara granosa* substitution flour indicates the *Anadara granosa* flour's potential as an alternative source of zinc. This is shown in several studies, where *Anadara granosa* are found to contain

zinc level below the maximum allowed level, which is under 100 ppm as stipulated in Malaysian Food Regulation 1985 (Ministry of Health Malaysia, 1985). In the meantime, the fresh *Anadara granosa* from Boalemo, Gorontalo Province, Indonesia is found to contain 13.91 ppm zinc, while the dried *Anadara granosa* contain 54.27 ppm zinc the boiled *Anadara granosa* contain 12.99 ppm zinc, and the dried, boiled *Anadara granosa* contain 37.86 ppm of zinc (Nurjanah *et al.*, 2005). On top of that, the *Anadara granosa* from Pohuwato regency of Gorontalo Province Indonesia was found to contain 2.70 – 2.82 ppm (Solang *et al.*, 2013). In addition, the *Anadara granosa* taken from Tanjung Mas and Wedung water of Semarang, Indonesia contains zinc 68.13 – 94.22 mg/kg of wet weight (Taurusiana *et al.*, 2014).

This present study also revealed that substitution of *Anadara granosa*'s flour could increase the protein level. This signifies that the substitution of *Anadara granosa*'s flour is one of the alternatives to improve the level of protein within the meal. This present finding supports Subaryono *et al.* (2003) who found that chips added with *Anadara granosa* had the protein level of 16.51%. Solang *et al.* (2017) also described that supplementation of *Anadara granosa* increased the level of protein in *Cireng* snack with a percentage of 5.05% - 54.49%. Several other studies showed that *Anadara granosa* were considered as important source of protein in tropical, subtropical, and warm climate regions (Broom, 1985; Ibrahim, 1995; Nurnadia *et al.*, 2011). The protein level of *Anadara granosa* was 19.8 % (Nurjanah *et al.*, 2005).

This present study has pointed out that the higher the composition of *Anadara granosa* flour into a meal, the meal's zinc and protein level will also increase. Kaji and Nishi (2006) also found that main meals composed of eggs, milk, poultry, and fish have lower zinc: protein ratio compared to meals made of cockles, beef, and other red meats.

4.2 Albumin serum level

This present study shows that the malnourished rat given the dried rice were low in zinc level by 0.791 ppm, in protein level by 8.46%, as well as low in albumin level by 46%. It is also shown that the substitution of *Anadara granosa* flour increased the albumin level on the tested rat. The malnourished rats had an albumin level of 3.01 g/dL. The level is considered below the normal level of albumin for a rat, which was 3.3 g/dL (Giknis and Clifford, 2008; Susanto *et al.*, 2010). Meanwhile, the provision of feed substituted with *Anadara granosa*'s flour could increase the albumin level of rat into the normal range, about 3.64 -3.69 g/dL (Table 3). This finding is similar to Giknis *et al.* (2008) who found that normal male rat had the range of albumin level between 3.4 – 4.8 g/dL. This present study is also similar to Kuwahata *et al.*, (2017) who discovered that albumin concentration on rats decreased due to the decrease of protein intake and it returned to normal when the malnutrition rats were given protein intake by 20% of casein through ad libitum method. Khasanah *et al.* (2015), Abdullahi *et al.* (2018), and Gounden *et al.* (2018) stated that protein deficiency in food intake could lead to lack of various essential amino acid in blood serum, which needed to develop cells (synthesis) and for metabolism process as an amino acid is the precursor for albumin synthesis. The lack of amino acid in this serum would lead to a lack of albumin liver production (protein).

Provision of feed substituted with *Anadara granosa* flour could improve the level of albumin in malnutrition rat. This is assumed to correlate with the increase of zinc and protein level on the substituted feed (Tables 1 and 2). Choundhary (2013), described that the provision of zinc could increase protein serum. In turn, the protein contents could increase the absorption and transportation of zinc. Moreover, the presence of protein could increase the availability of amino acid as the precursor for albumin synthesis. Synthesis of albumin depends on the adequate amino acid supply (Marshall, 2012). Meanwhile, Shidhu *et al.* (2004) showed that the provision of zinc on protein-deficient rats could help increase the level of hepatic protein. This zinc ability is linked to its role to induce metallothionein (zinc-binding protein). Thus, it regulates the amino acid as the

precursor for albumin synthesis (Tekeli, 2002).

According to the results, the *Anadara granosa* flour contains the amino acid, glutamate, aspartate, serine, histidine, glycine, arginine, alanine, tyrosine, methionine, valine, phenylalanine, isoleucine, leucine, and lysine. In the present study, the analysis of cysteine and proline was not performed since there was no standard for examining these amino acids. The common amino acid found in the study is glutamate. Azis (2007) reports that *Anadara granosa* contain cysteine. The amino acids that contribute to the metabolism of zinc are cysteine and histidine (Snedeker and Greger, 1983; Pace and Weerapana, 2014). Both of these amino acids function to transport zinc by facilitating the process of the formation of zinc-histidine and zinc-cysteine complex.

Types of amino acids that serve as a precursor of albumin synthesis are lysine, tryptophan, and isoleucine. Isoleucine and tryptophan also play a role in increasing the value of albumin synthesis (Kelman *et al.*, 1972). Hutson *et al.* (1987, as cited in Harp *et al.*, 1991) argue that the deficit of essential amino acid content, such as lysine, tryptophan, and isoleucine, can decrease the albumin release. Therefore, zinc and protein in *Anadara granosa* were assumed to work simultaneously to increase the availability of amino acid as a precursor for albumin synthesis; hence, increase the albumin level of malnutrition rat.

4.3 Bodyweight

This present study showed that the provision of dried rice feed could decrease the bodyweight of the test rat, while the *Anadara granosa* flour substitution feed was proven to increase the bodyweight of the tested rat. A decrease in the bodyweight of the tested rats was assumed to be caused by the low zinc and protein contents in the dried rice feed. The zinc level of dried rice was below 1 ppm (Choundhary, 2013). Meanwhile, the protein level of dried rice was 8%; thus, it was considered unsuitable for the rats' protein needs at 12% (complete protein with 20 amino acids) (Smith and Mangkoewidjojo, 1987). Moreover, Shidhu *et al.* (2004) reported that protein deficiency decreases body weight. Meanwhile, a decrease in body weight could also blame zinc deficiency (Rossi *et al.*, 2001; Ishikawa *et al.*, 2008; Parveen and Dipti, 2016). The provision of *Anadara granosa* substituted feed could significantly improve bodyweight. This was suspected due to the increased zinc and protein level in the feed (Tables 1 and 2). Parveen and Dipti (2016) explained that zinc supplementation causes a quick bodyweight increase in the malnutrition rehabilitation phase. Budiastutik *et al.* (2011) also described that supplementation of zinc

phosphate and biscuit increases body weight and height. Meanwhile, an increase of protein on the feed also regulates the bodyweight through regulating the mechanism of thermogenesis and body composition, food intake, as well as protein synthesis (Westerterp-Plantenga, 2003; Greco *et al.*, 2017). Further, Westerterp-Plantenga (2003) also explained that animal-based protein-induced higher thermogenesis than plant-based protein.

The increase of body weight in malnutrition rat fed with *Anadara granosa* substituted feed was assumed to correlate with the increase of albumin serum as observed in this study. The increase of albumin levels would optimize the zinc absorption of the feed. Marshall (2012) explained that albumin functions to transport zinc. As the zinc has been appropriately absorbed, the zinc content within the body was assumed to increase. This availability of zinc within the body would accelerate growth and cell differentiation (Parveen and Dipti, 2016).

5. Conclusion

This study has shown that the provision of feed substituted with *Anadara granosa* flour could improve the zinc level and protein of the feed. The increase of zinc and protein could improve the availability of zinc and protein within the body; thus, it will, in turn, improve the growth of malnutrition rats through the mechanism of albumin level improvement, which later increases growth, particularly through the increase of body weight.

Conflict of interest

The authors do not have any conflicts of interest regarding the content of the present work.

Acknowledgments

We would like to acknowledge the contribution of DP2M DIKTI for funding this research through the Competitive Grant Scheme in 2015.

References

Abdullahi, S.M., Yakubu, A.M., Bugaje, M.A. and Akuyam, S.M. (2018). Serum total protein and albumin levels among malnourished children aged 6-59 months in Zaria. *Nigerian Journal of Paediatrics*, 45(1), 15-18. <https://doi.org/10.4314/njp.v45i1.4>

Adriani, M. and Wirjatmadi, B. (2014). The effect of adding zinc to vitamin A on IGF-1, bone age, and linear growth in stunted children. *Journal of Trace Elements in Medicine and Biology*, 28(4), 431-435.

<https://doi.org/10.1016/j.jtemb.2014.08.007>

Agustian, L., Sembiring, T. and Ariani, A. (2009). Peran Zinkum Terhadap Pertumbuhan Anak/ The role of zinc on child's growth. *Sari Pediatri*, 11(4), 244-249. <https://doi.org/10.14238/sp11.4.2009.244-9>

AOAC [Association of Official Analytical Chemist]. (1995). Official methods of analysis of the association of official analytical chemist. 18th ed. Gaithersburg, USA: AOAC International.

Broom, M.J. (1985). The biology and culture of marine bivalve mollusks of the genus *Anadara*. Philippines: International Center for Living Aquatic Resources Management.

Budiastutik, I., Wirjatmadi, B. and Adriani, M. (2011). Pengaruh suplementasi zinc sulfat dan biscuit terhadap konsentrasi zinc rambut balita (program MP ASI biskuit di Kertosono Jawa Timur). *Buletin Penelitian Sistem Kesehatan*, 14(3), 270-281. [In Bahasa Indonesia].

Campbell, D.T. and Stanley, J.C. (1963). *Experimental and quasi-experimental designs for research*. Chicago: Rand McNally College Publishing Company.

Doumas, B.T. and Peters, T.Jr. (2009). Origin of the dye-binding method for measuring serum albumin. *Clinical Chemistry*, 55(3), 583-584. <https://doi.org/10.1373/clinchem.2008.112854>

Ministry of Health Malaysia (1985). Malaysian Food Regulations 1985. Retrieved September 2019, from https://extranet.who.int/nutrition/gina/sites/default/files/MYS%201985%20Food%20Regulations_0.pdf

Gibson, R.S. and Ferguson, E.L. (1994). Dietary Strategy for preventing iron and zinc deficiency in Africa children presented at the Nutrition in a sustainable environment: Proceedings of the XVth International Congress of Nutrition, p. 301-303. Adelaide, Australia: Smith-Gordon.

Giknis M.L.A. and Clifford, C.B. (2008). Clinical laboratory parameters for CrI: WI (Han). Wilmington, MA: Charles River Laboratory International.

Gounden, V., Vashisht, R. and Ishwarlal, J. (2018). Hypoalbuminemia. Treasure Island, Florida, USA: Stat Pearls Publishing LLC.

Greco, E., Winquist, A.Lee, T.J., Collins, S. and Lebovic, Z. (2017). The Role of Source of Protein in Regulation of Food Intake, Satiety, Body Weight, and Body Composition. *Journal of Nutritional Health and Food Engineering*, 6(6), 186-193. <https://doi.org/10.15406/jnhfe.2017.06.00223>

Handley, M.A., Lyles, C.R., McCulloch, C. and Cattamanchi, A. (2018). Selecting and Improving

- Quasi-Experimental Designs in Effectiveness and Implementation Research. *The Annual Review of Public Health*, 39, 5-25. <https://doi.org/10.1146/annurev-publhealth-040617-014128>
- Harp, J.B., Goldstein, S. and Phillips, L.S. (1991). Molecular regulation of IGF-I by amino acid availability in Cultured hepatocytes. Nutrition and somatomedin. *Diabetes*, 40(1), 95-101. <https://doi.org/10.2337/diab.40.1.95>
- Chirande, L., Charwe, D., Mbwana, H., Victor, R., Kimboka S, Issaka, A.I, Baines, S.K., Dibly, M.J. and Agho, K.E. (2015). Determinants of Stunting and Severe Stunting Among Under-fives in mid-Tanzania: Evidence from the 2010 cross-sectional household survey. *BMC Pediatrics*, 15(165), 1-13. <https://doi.org/10.1186/s12887-015-0482-9>
- Choundhary, D. (2013). Influence of dietary zinc deficiency on serum zinc and protein. *Indian Journal of Fundamental and Applied Life Sciences*, 3(1), 143-148.
- Ibrahim, N. (1995). The trace element content of Malaysian cockles (*Anadara granosa*). *Food Chemistry*, 54(2), 133-135. [https://doi.org/10.1016/0308-8146\(94\)00149-Y](https://doi.org/10.1016/0308-8146(94)00149-Y)
- Ishikawa, Y., Kudo, H., Suzuki, S., Nemoto, N., Sassa, S. and Sakamoto, S. (2008). Downregulation by a low- zinc diet in gene expression of rat prostatic thymidylate synthase and thymidine kinase. *Nutrition and Metabolism*, 5(12), 1-5. <https://doi.org/10.1186/1743-7075-5-12>
- Kaji, M. and Nishi, Y. (2006). Growth and minerals: Zinc. *Growth, Genetics and Hormones (GGH)*, 22 (1), 1-10.
- Kartono, D.J., Hardinsyah, Jahari, A.B., Sulaeman, A., Astuti, M., Soekatri, M. and Riyadi H. (2012). Penyempurnaan Kecukupan Gizi Untuk Orang Indonesia. *Widyakarya Nasional Pangan Dan Gizi X.20-21 November 2012*. Jakarta. Retrieved September, 2019, from <https://www.dropbox.com/s/zj1wbi5ld0qwul/akg%202012.pdf>. [In Bahasa Indonesia].
- Kaur, K., Gupta, R., Saraf, S.A. and Saraf, S.K. (2014). Zinc: The Metal of Life. *Comprehensive Reviews in Food Science and Food Safety*, 13(4), 358- 376. <https://doi.org/10.1111/1541-4337.12067>
- Kelman L., Saunders S.J., Wicht S., Frith L., Corrigan A., Kirsch R.E. and Terblanche J. (1972). The Effects of Amino Acids on Albumin Synthesis by the Isolated Perfused Rat Liver. *Biochemical Journal*, 129(4), 805-809. <https://doi.org/10.1042/bj1290805>
- Kemenkes. (2018). Riset Kesehatan Daerah. Jakarta: Badan penelitian dan pengembangankesehatan. Indonesia: Kementerian Kesehatan, Republik Indonesia. [In Bahasa Indonesia].
- Khasanah, Y., Ratnayani, Ariani, D., Angwar, M. and Nuraeni, T. (2015). *In Vivo* Study on Albumin and Total Protein in White Rat (*Rattus norvegicus*) after Feeding of Enteral Formula from Tempe and Local Food. *Procedia Food Science*, 3, 274-279. <https://doi.org/10.1016/j.profoo.2015.01.030>
- Kusudaryati, D.P.D., Muis, S.F. and Widajanti, L. (2017). Pengaruh suplementasi Zn terhadap perubahan indeks TB/U anak stunted usia 24-36 bulan. *Jurnal Gizi Indonesia*, 5(2), 98-104. <https://doi.org/10.14710/jgi.5.2.98-104>
- Kuwahata, M., Mari, H., Yukiko, K., Yasuaki, W. and Yasuhiro, K. (2017). An oxidized/reduced state of plasma albumin reflects malnutrition due to an insufficient diet in rats. *Journal of Clinical Biochemistry and Nutrition*, 60(1), 70-75. <https://doi.org/10.3164/jcbrn.16-33>
- Mardewi, K.W., Sidiartha, I.G.L. and Gunawijaya, E. (2016). Low serum zinc and short stature in children. *Paediatrica Indonesiana*, 56(3), 171-175. <https://doi.org/10.14238/pi56.3.2016.171-5>
- Marshall, W. (2012). Albumin (*serum, plasma*). Association for Clinical Biochemistry. Retrieved on September 2019 from website: <http://www.acb.org.uk/Nat%20Lab%20Med%20Hbk/Albumin.pdf>
- McDonald, J.H. (2014). Handbook of Biological Statistics. 3rd ed. Baltimore, Maryland: Sparky House Publishing.
- Nurjanah, Zulhamsyah and Kustiyariyah. (2005). Kandungan mineral dan proksimatkerendahan (*Anadara granosa*) yang diambil dari kabupaten Boalemo, Gorontalo. *Buletin Teknologi Hasil Perikanan*, 8(2), 15-24. <https://doi.org/10.17844/jphpi.v8i2.1012>. [In Bahasa Indonesia].
- Nurnadia, A.A., Azrina, A. and Amin, I. (2011). Proximate composition and energetic value of selected marine fish. *International Food Research Journal*, 18, 137-148.
- Pace, N.J. and Weerapana, E. (2014). Zinc-Binding Cysteines: Diverse Functions and Structural Motifs. *Biomolecules*, 4(2), 419-434. <https://doi.org/10.3390/biom4020419>
- Parveen, M. and Dipti, B. (2016). Role of Zinc in Malnutrition. *Annals of Nutritional Disorders and Therapy*, 3(1), 1-4.
- Rossi, L., Migliaccio, S., Corsi, A., Marzia, M., Bianco, P., Teti, A., Gambelli, L., Cianfarani, S., Paoletti, F. and Branca, F. (2001). Reduced growth and skeletal

- changes in zinc-deficient growing rats are due to Impaired growth plate activity and inanition. *American Society for Nutrition Sciences*, 131(4), 1142-1148. <https://doi.org/10.1093/jn/131.4.1142>
- Shidhu, P., Garg, M.L. and Dhawan, D.K. (2004). Protective effects of zinc on oxidative stress enzymes in the liver of protein-deficient rats. *Nutricion Hospitalaria*, 19(6), 341-347.
- Smith, J.B. and Mangkoewidjojo, S. (1987). The care, breeding, and management of experimental animals for research in the tropics. Canberra: International Development Program of Australia Universities and Collages (IDP).
- Snedeker, S.M. and Greger, J.L. (1983). Metabolism of Zinc, Copper and Iron As Affected by Dietary Protein, Cysteine and Histidine. *The Journal of Nutrition*, 113(3), 644-652. <https://doi.org/10.1093/jn/113.3.644>
- Solang, M., Adriani, M. and Wirjatmadi, B. (2013). The Analysis of Blood Cockles (*Anadara granosa*) Flour Supplementasi on The Concentration of Zinc, IGF-I, and Ephifiseal Plate Width of Femur Malnourished Male Rats (*Rattus norvegicus*). *IEESE International Journal of Science and Technology*, 2(4), 12-24.
- Solang, M., Lamondo, D. and Kumaji, S.S. (2017). Zinc, calcium, protein, lead, mercury, and the sensorics quality of cireng snacks supplemented with blood cockle (*Anadara granosa*). *Nusantara Bioscience*, 9 (4), 385-39. <https://doi.org/10.13057/nusbiosci/n090408>
- Steel, R.G.D. and Torrie, J.H. (1980). Principles of Statistics for University. 2nd ed., p. 168-214. California: McGraw Hill.
- Subaryono, Irianto, H.E. and Indriati, N. (2003). Fortifikasi ikan pada emping mlijo. *Journal penelitian perikanan Indonesia*, 9(5), 85 -94. <https://doi.org/10.15578/jppi.9.5.2003.85-94>
- Susanto, H., Maslikah, S.I. and Hernowati, T.E. (2010). The nutritional effects of Moringa leaf powder (*Moringa oleifera*) varieties of East Nusa Tenggara in albumin levels of protein-energy malnutrition of Wistar rats (*In vivo* study of *Moringa* as a candidate for supplementation therapy in cases of severe malnutrition). Prosiding Seminar Nasional MIPA: "Peran MIPA dalam Pengembangan Teknologi dan Pendidikan Berkarakter Menuju Bangsa Mandiri". Malang, Indonesia: Universitas Negeri Malang.
- Taurusiana, S., Afiati, N. and Widyorini, N. (2014). Kajian Kandungan Logam Berat Besi (Fe) Dan Seng (Zn) Pada Jaringan Lunak Kerang Darah (*Anadara granosa* (L.) Di Perairan Tanjung Mas, Semarang Dan Perairan Wedung, Demak. *Diponegoro Journal* of *Maquares Management of Aquatic Resources*, 3 (1), 143-150. <https://doi.org/10.14710/marj.v3i1.4431> [In Bahasa Indonesia].
- Tekeli, S.K. (2002). The study of effects on serum glucose, total lipid, total protein, and albumin levels of orally zinc in rats. *Trace Elements and Electrolytes*, 19(1), 6-10.
- Westerterp-Plantenga, M.S. (2003). The significance of protein in food intake and body weight regulation. *Current Opinion in Clinical Nutrition and Metabolic Care*, 6(6), 635-638. <https://doi.org/10.1097/00075197-200311000-00005>
- Yanagisawa, H. (2004). Zinc Deficiency and Clinical Practice. *Japan Medical Association Journal*, 47(8), 359-364. <https://doi.org/10.1248/yakushi.128.333>

Anadara granosa substitution in feed to improve the zinc, protein of the feed, serum albumin, and body weight of malnourished rats

ORIGINALITY REPORT

14%	14%	0%	1%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	10%
2	biosains.mipa.uns.ac.id Internet Source	3%
3	Submitted to Program Pascasarjana Universitas Negeri Yogyakarta Student Paper	1%

Exclude quotes	Off	Exclude matches	< 1%
Exclude bibliography	On		