



ISTANBUL  
UNIVERSITY  
PRESS

Indexed in  
Web of Science



# Istanbul Journal of Pharmacy

## Original Articles

**Evaluation of drug-drug interactions and their clinical importance in a pediatric hematopoietic stem cell transplantation unit**  
Neslişül Özdemir, Ayçe Çeliker, Bülent Barış Kuşkonmaz, Duygu Uçkan Çetinkaya

**Alpha lipolic acid bioequivalence study redesigned: a candidate for highly variable drugs**  
Burcu Bulut, Nagehan Sarraçoğlu, Onur Pinarbaşı

**Development and validation an HPLC - UV method for determination of esomeprazole and pirfenidone simultaneously in rat plasma: application to a drug monitoring study**  
Emrah Dural, Sema Tülay Köz, Süleyman Köz

**Preparation and *in vitro* studies of fixed-dose tablet combination of repaglinide and metformin**  
Sinem Şahin, Burcu Mesut, M. Ezgi Durgun, Esher Özçelik, Yıldız Özsoy

**The formulation and characterization of water-soluble snakehead fish (*Ophiocephalus striatus*) dry extract in nanoemulsion using permeation and *in vivo* study**  
Robert Tungadi, Widysusanti Abdulkadir, Munafri Tahir

**Pseudo ternary phase diagrams: a practical approach for the area and centroid calculation of stable microemulsion regions**  
Murat Sami Berkman, Kadri Güleç

**Rational design and synthesis of quinazoline derivatives**  
Fatimah AlKharaz, Zead Abudayeh, Qais Abualassal, Loay Hassounah

**Molecular and crystal structure of 1-methyl-5-trifluoromethoxy-1*H*-indole-2,3-dione 3-[4-(4-methoxyphenyl)thiosemicarbazone]**  
Özge Soylu Eter, Zeliha Atioğlu, Mehmet Akkurt, Cem Cüneyt Ersanlı, Nilgün Karalı

**Synthesis, characterization and antibacterial evaluation of new pyridyl-thiazole hybrids of sulfonamides**  
Zafer Şahin, Sevd Nur Biltekin, Leyla Yurttaş, Şeref Demirayak

**The effects of vanadyl sulfate on glutathione, lipid peroxidation and nonenzymatic glycosylation levels in various tissues in experimental diabetes**  
Sevim Tunali, Refiye Yanardağ

**Short-term adaptive metabolic response of *Escherichia coli* to ciprofloxacin exposure**  
Engin Koçak, Ceren Özkul

***In vitro* urease and trypsin inhibitory activities of some sulfur compounds**  
Eda Dağsuyu, Refiye Yanardağ

**Quality of turmeric powder in herbal stores: pharmacognostical investigations on turmeric powders obtained from herbal stores in Istanbul, Turkey**  
Ebru Kuruldağ, Fatmanur Yılmaz, Gizem Gülsoy Toplan, Berna Özbek Çelik, Afife Mat

**Isolation of potential liver x receptor alpha agonist and antioxidant compounds from *Hypericum microcalycinum* Boiss. & Heldr.**  
Seçil Sarıkaya Aydın, Vahap Murat Kutluay, Toshiaki Makino, Makoto Inoue, Ümmühan Şebnem Harput, İclal Saraçoğlu

***In vitro* evaluation of antimicrobial activity of *Distemonanthus benthamianus* chewing stick extract mouthwash**  
Mbang Nyong Femi-Oyewo, Olatayo Ademola Adeleye, Caroline Olufunke Babalola, Olufemi Babatunde Banjo, Modupe Nofisat Adebawale, Florence Olubola Odeleye

**Phytochemical screening, phenolic content and antioxidant activity of *Lavandula* species extracts from Algeria**  
Farah Haddouchi, Tarik Mohammed Chaouche, Meriem Saker, Imane Ghellai, Ouhiba Boudjemai

**Antimicrobial activities of some narrow endemic gypsophytes**  
Esma Ocak, Şule İnci, Derviş Öztürk, Sanem Akdeniz Şafak, Ebru Özdeniz, Sevd Kırbağ, Ahmet Harun Evren, Latif Kurt

***Chelidonium majus* L. (Papaveraceae) morphology, anatomy and traditional medicinal uses in Turkey**  
Golshan Zare, Neziha Yağmur Diker, Zekiye Ceren Arıtuluk, İffet İrem Tatlı Çankaya

**The anatomical properties of endemic *Hypericum kotschyannum* Boiss.**  
Onur Altınbaşak, Gülay Ecevit Genç, Şükran Kültür

## Review Articles

**Fungal homoserine transacetylase: a potential antifungal target**  
Esra Seyran

**Bioavailability of berberine: challenges and solutions**  
Asha Thomas, Seema Kamble, Sanjeevani Deshkar, Lata Kothapalli, Sohan Chitlange

**Evaluation of toxic effects of statins and their possible role in treatment of cancer**  
Aysun Ökçesiz, Ülkü Ünüğer Bucurğat



# Istanbul Journal of Pharmacy

## INDEXING AND ABSTRACTING

Web of Science - Emerging Sources Citation Index (ESCI)

TÜBİTAK-ULAKBİM TR Index



# Istanbul Journal of Pharmacy

## OWNER

Prof. Dr. Erdal CEVHER

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Technology, İstanbul, Turkey

## RESPONSIBLE MANAGER

Asst. Prof. Bahar GÜRDAL ABAMOR

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, İstanbul, Turkey

## CORRESPONDENCE ADDRESS

Istanbul University, Faculty of Pharmacy,  
Department of Pharmaceutical Botany,  
Beyazıt, 34116, Fatih / İstanbul, Turkey

Phone: +90 212 440 02 75

Fax: +90 212 440 02 52

E-mail: [akaline@istanbul.edu.tr](mailto:akaline@istanbul.edu.tr)

<https://dergipark.org.tr/tr/pub/iujp>

<https://iupress.istanbul.edu.tr/tr/journal/ijp/home>

## PUBLISHER

Istanbul University Press

Istanbul University Central Campus,  
34452 Beyazıt, Fatih / İstanbul, Turkey

Phone: +90 212 440 00 00

---

Authors bear responsibility for the content of their published articles.

The publication languages of the journal is English.

This is a scholarly, international, peer-reviewed and open-access journal published triannually in April, August and December.

---

**Publication Type:** Periodical

---



# Istanbul Journal of Pharmacy

## EDITORIAL MANAGEMENT

### Editor-in-Chief

Prof. Dr. Emine AKALIN

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, İstanbul, Turkey – [akaline@istanbul.edu.tr](mailto:akaline@istanbul.edu.tr)

### Co-Editors in Chief

Prof. Dr. B. Sönmez UYDEŞ DOĞAN

Istanbul University, Faculty of Pharmacy, Department of Pharmacology, İstanbul, Turkey – [sonmezdo@istanbul.edu.tr](mailto:sonmezdo@istanbul.edu.tr)

Prof. Dr. Nilgün KARALI

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, İstanbul, Turkey – [karalin@istanbul.edu.tr](mailto:karalin@istanbul.edu.tr)

### Section Editors

Prof. Dr. Nuriye AKEV

Istanbul University, Faculty of Pharmacy, Department of Biochemistry, İstanbul, Turkey – [nakev@istanbul.edu.tr](mailto:nakev@istanbul.edu.tr)

Prof. Dr. Nilgün KARALI

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, İstanbul, Turkey – [karalin@istanbul.edu.tr](mailto:karalin@istanbul.edu.tr)

Prof. Dr. B. Sönmez UYDEŞ DOĞAN

Istanbul University, Faculty of Pharmacy, Department of Pharmacology, İstanbul, Turkey – [sonmezdo@istanbul.edu.tr](mailto:sonmezdo@istanbul.edu.tr)

Prof. Dr. Sevgi GÜNGÖR

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Technology, İstanbul, Turkey – [sgungor@istanbul.edu.tr](mailto:sgungor@istanbul.edu.tr)

Asst. Prof. Bahar GÜRDAL ABAMOR

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, İstanbul, Turkey – [bahar.gurdal@istanbul.edu.tr](mailto:bahar.gurdal@istanbul.edu.tr)

### Language Editors

Elizabeth Mary EARL

Istanbul University, Department of Foreign Languages, İstanbul, Turkey – [elizabeth.earl@istanbul.edu.tr](mailto:elizabeth.earl@istanbul.edu.tr)

Alan James NEWSON

Istanbul University, Department of Foreign Languages, İstanbul, Turkey – [alan.newson@istanbul.edu.tr](mailto:alan.newson@istanbul.edu.tr)

### Statistics Editor

Prof. Dr. Abdülbari BENER

Istanbul University-Cerrahpasa, Cerrahpasa Faculty of Medicine, Department of Biostatistics and Medical Informatics, İstanbul, Turkey – [abdulbari.bener@iuc.edu.tr](mailto:abdulbari.bener@iuc.edu.tr)

### Scientific Secretariat

Asst. Prof. Gülsev ÖZEN

Istanbul University Faculty of Pharmacy, Department of Pharmacology, İstanbul, Turkey – [gulsevozen@istanbul.edu.tr](mailto:gulsevozen@istanbul.edu.tr)



## EDITORIAL BOARD

### **Afife MAT**

Istanbul University, Faculty of Pharmacy, Department of Pharmacognosy, İstanbul, Turkey – [afifemat@gmail.com](mailto:afifemat@gmail.com)

### **Berna ÖZBEK-ÇELİK**

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Microbiology, İstanbul, Turkey – [berna.ozbek@istanbul.edu.tr](mailto:berna.ozbek@istanbul.edu.tr)

### **Bilge ŞENER**

Gazi University, Faculty of Pharmacy, Department of Pharmacognosy, Ankara, Turkey – [bilgesener11@gmail.com](mailto:bilgesener11@gmail.com)

### **Carsten EHRHARDT**

Trinity College Dublin, School of Pharmacy and Pharmaceutical Sciences and Trinity Biomedical Sciences Institute, Dublin, Ireland – [ehrhards@tcd.ie](mailto:ehrhards@tcd.ie)

### **Claudiu T. SUPURAN**

University of Florence, Section of Pharmaceutical and Nutriceutical Sciences, Neurofarba Department, Florence, Italy – [claudiu.supuran@unifi.it](mailto:claudiu.supuran@unifi.it)

### **Domenico Vittorio DELFINO**

University of Perugia, Department of Medicine and Surgery, Perugia, Italy – [domenico.delfino@unipg.it](mailto:domenico.delfino@unipg.it)

### **Erden BANOĞLU**

Gazi University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Ankara, Turkey – [banoglu@gazi.edu.tr](mailto:banoglu@gazi.edu.tr)

### **Fatma AKAR**

Gazi University, Faculty of Pharmacy, Department of Pharmacology, Ankara, Turkey – [fakar@gazi.edu.tr](mailto:fakar@gazi.edu.tr)

### **Gianniantonio DOMINA**

University of Palermo, Food and Forest Sciences, Department of Agricultural, Palermo, Italy – [gianniantonio.domina@unipa.it](mailto:gianniantonio.domina@unipa.it)

### **İlkay KÜÇÜKGÜZEL**

Marmara University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, İstanbul, Turkey – [ikucukguzel@marmara.edu.tr](mailto:ikucukguzel@marmara.edu.tr)

### **Johan Van de VOORDE**

Ghent University, Department of Pharmacology, Gent, Belgium – [johan.vandevoorde@ugent.be](mailto:johan.vandevoorde@ugent.be)

### **Melih ALTAN**

Bezmialem University, Faculty of Pharmacy, Department of Pharmacology, İstanbul, Turkey – [vmaltan@bezmialem.edu.tr](mailto:vmaltan@bezmialem.edu.tr)

### **Meral ÖZALP**

Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Microbiology, Ankara, Turkey – [mozalp@hacettepe.edu.tr](mailto:mozalp@hacettepe.edu.tr)

### **Müberra KOŞAR**

Eastern Mediterranean University, Faculty of Pharmacy, Department of Pharmacognosy, Famagusta, Northern Cyprus – [muberra.kosar@emu.edu.tr](mailto:muberra.kosar@emu.edu.tr)

### **Nilüfer YÜKSEL**

Ankara University, Faculty of Pharmacy, Department of Pharmaceutical Technology, Ankara, Turkey – [nyuksel@pharmacy.ankara.edu.tr](mailto:nyuksel@pharmacy.ankara.edu.tr)

### **Nurşen BAŞARAN**

Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Toxicology, Ankara, Turkey – [nbasaran@hacettepe.edu.tr](mailto:nbasaran@hacettepe.edu.tr)

### **Oya ALPAR**

Altınbaş University, Faculty of Pharmacy, Department of Pharmaceutical Technology, İstanbul, Turkey and Department of Pharmaceutical Technology, UCL, UK – [oya.alpar@altinbas.edu.tr](mailto:oya.alpar@altinbas.edu.tr)

### **Özlem Nazan ERDOĞAN**

Istanbul University, Faculty of Pharmacy, Department of Pharmacy Management, İstanbul, Turkey – [nazan.erdogan@istanbul.edu.tr](mailto:nazan.erdogan@istanbul.edu.tr)

### **Sıdıka TOKER**

Istanbul University, Faculty of Pharmacy, Department of Analytical Chemistry, İstanbul, Turkey – [serturk@istanbul.edu.tr](mailto:serturk@istanbul.edu.tr)

### **Sibel ÖZDEN**

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Toxicology, İstanbul, Turkey – [stopuz@istanbul.edu.tr](mailto:stopuz@istanbul.edu.tr)

### **Stephen R. DOWNIE**

University of Illinois, Department of Plant Biology, Urbana, Illinois, USA – [sdownie@illinois.edu](mailto:sdownie@illinois.edu)

### **Tao CHEN**

Medical College of Soochow University, School of Public Health, Department of Toxicology, Suzhou, China – [tchen@suda.edu.cn](mailto:tchen@suda.edu.cn)

### **Ufuk KOLAK**

Istanbul University, Faculty of Pharmacy, Department of Analytical Chemistry, İstanbul, Turkey – [kolak@istanbul.edu.tr](mailto:kolak@istanbul.edu.tr)

### **Zeliha YAZICI**

Biruni University, Faculty of Medicine, Department of Medical Pharmacology, İstanbul, Turkey – [zyazici@biruni.edu.tr](mailto:zyazici@biruni.edu.tr)



# Istanbul Journal of Pharmacy

## AIMS AND SCOPE

Istanbul Journal of Pharmacy (Istanbul J Pharm) is an international, scientific, open access periodical published in accordance with independent, unbiased, and double-blinded peer-review principles. The journal is the official publication of İstanbul University Faculty of Pharmacy and it is published triannually on April, August, and December. The publication language of the journal is English.

Istanbul Journal of Pharmacy (Istanbul J Pharm) aims to contribute to the literature by publishing manuscripts at the highest scientific level on all fields of pharmaceutical sciences. The journal publishes original articles, short reports, letters to the editor and reviews.

The target audience of the journal includes specialists and professionals working and interested in all disciplines of pharmaceutical, also medicinal, biological and chemical sciences.

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Committee of Medical Journal Editors (ICMJE), World Association of Medical Editors (WAME), Council of Science Editors (CSE), Committee on Publication Ethics (COPE), European Association of Science Editors (EASE), and National Information Standards Organization (NISO). The journal is in conformity with the Principles of Transparency and Best Practice in Scholarly Publishing ([doaj.org/bestpractice](http://doaj.org/bestpractice)).

Istanbul Journal of Pharmacy is currently indexed in Web of Science-Emerging Sources Citation Index, TUBITAK ULAKBIM TR Index and CAS database.

Processing and publication are free of charge with the journal. No fees are requested from the authors at any point throughout the evaluation and publication process. All manuscripts must be submitted via the online submission system, which is available at <http://dergipark.gov.tr/iujp>. The journal guidelines, technical information, and the required forms are available on the journal's web page.

All expenses of the journal are covered by the İstanbul University Faculty of Pharmacy. Potential advertisers should contact the Editorial Office. Advertisement images are published only upon the Editor-in-Chief's approval.

Statements or opinions expressed in the manuscripts published in the journal reflect the views of the author(s) and not the opinions of the İstanbul University Faculty of Pharmacy, editors, editorial board, and/or publisher; the editors, editorial board, and publisher disclaim any responsibility or liability for such materials.

Istanbul Journal of Pharmacy is an open access publication and the journal's publication model is based on Budapest Open Access Initiative (BOAI) declaration. Journal's archive is available online, free of charge at <http://ijp.istanbul.edu.tr>. İstanbul Journal of Pharmacy's content is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



**Editor in Chief:** (Prof. Dr.) Emine AKALIN

**Address:** İstanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Beyazıt, 34116, Fatih İstanbul

**Phone:** +90 212 440 02 75

**Fax:** +90 212 440 02 52

**E-mail:** [akaline@istanbul.edu.tr](mailto:akaline@istanbul.edu.tr)

**Publisher:** İstanbul University Press

**Address:** İstanbul University Central Campus, 34452 Beyazıt, Fatih / İstanbul - Turkey

**Phone:** +90 212 440 00 00



## INSTRUCTIONS TO AUTHORS

### Context

Istanbul Journal of Pharmacy (Istanbul J Pharm) is an international, scientific, open access periodical published in accordance with independent, unbiased, and double-blinded peer-review principles. The journal is the official publication of İstanbul University Faculty of Pharmacy and it is published triannually on April, August, and December. The publication language of the journal is English.

Istanbul Journal of Pharmacy (Istanbul J Pharm) aims to contribute to the literature by publishing manuscripts at the highest scientific level on all fields of pharmaceutical sciences. The journal publishes original articles, short reports, letters to the editor and reviews.

### Editorial Policy

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Council of Medical Journal Editors (ICMJE), the World Association of Medical Editors (WAME), the Council of Science Editors (CSE), the Committee on Publication Ethics (COPE), the European Association of Science Editors (EASE), and National Information Standards Organization (NISO). The journal conforms to the Principles of Transparency and Best Practice in Scholarly Publishing ([doaj.org/bestpractice](http://doaj.org/bestpractice)).

Originality, high scientific quality, and citation potential are the most important criteria for a manuscript to be accepted for publication. Manuscripts submitted for evaluation should not have been previously presented or already published in an electronic or printed medium. The journal should be informed of manuscripts that have been submitted to another journal for evaluation and rejected for publication. The submission of previous reviewer reports will expedite the evaluation process. Manuscripts that have been presented in a meeting should be submitted with detailed information on the organization, including the name, date, and location of the organization.

### Peer-Review Policy

Manuscripts submitted to İstanbul Journal of Pharmacy will go through a double-blind peer-review process. Each submission will be reviewed by at least two external, independent peer reviewers who are experts in their fields in order to ensure an unbiased evaluation process. The

editorial board will invite an external and independent editor to manage the evaluation processes of manuscripts submitted by editors or by the editorial board members of the journal. The Editor in Chief is the final authority in the decision-making process for all submissions.

### Ethical Principles

An approval of research protocols by the Ethics Committee in accordance with international agreements (World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects," amended in October 2013, [www.wma.net](http://www.wma.net)) is required for experimental, clinical, and drug studies. If required, ethics committee reports or an equivalent official document will be requested from the authors. For manuscripts concerning experimental research on humans, a statement should be included that shows that written informed consent of patients and volunteers was obtained following a detailed explanation of the procedures that they may undergo. For studies carried out on animals, the measures taken to prevent pain and suffering of the animals should be stated clearly. Information on patient consent, the name of the ethics committee, and the ethics committee approval number should also be stated in the Materials and Methods section of the manuscript. It is the authors' responsibility to carefully protect the patients' anonymity. For photographs that may reveal the identity of the patients, signed releases of the patient or of their legal representative should be enclosed.

### Plagiarism

All submissions are screened by a similarity detection software (iThenticate by CrossCheck) at any point during the peer-review or production process. Even if you are the author of the phrases or sentences, the text should not have unacceptable similarity with the previously published data.

When you are discussing others' (or your own) previous work, please make sure that you cite the material correctly in every instance.

In the event of alleged or suspected research misconduct, e.g., plagiarism, citation manipulation, and data falsification/fabrication, the Editorial Board will follow and act in accordance with COPE guidelines



# Istanbul Journal of Pharmacy

## Authorship

Each individual listed as an author should fulfill the authorship criteria recommended by the International Committee of Medical Journal Editors (ICMJE - [www.icmje.org](http://www.icmje.org)). The ICMJE recommends that authorship be based on the following 4 criteria:

1. Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND
2. Drafting the work or revising it critically for important intellectual content; AND
3. Final approval of the version to be published; AND
4. Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

In addition to being accountable for the parts of the work he/she has done, an author should be able to identify which co-authors are responsible for specific other parts of the work. In addition, authors should have confidence in the integrity of the contributions of their co-authors.

All those designated as authors should meet all four criteria for authorship, and all who meet the four criteria should be identified as authors. Those who do not meet all four criteria should be acknowledged in the title page of the manuscript.

Istanbul Journal of Pharmacy requires corresponding authors to submit a signed and scanned version of the authorship contribution form (available for download through [http://ijp.istanbul.edu.tr/en/\\_](http://ijp.istanbul.edu.tr/en/_)) during the initial submission process in order to act appropriately on authorship rights and to prevent ghost or honorary authorship. If the editorial board suspects a case of "gift authorship," the submission will be rejected without further review. As part of the submission of the manuscript, the corresponding author should also send a short statement declaring that he/she accepts to undertake all the responsibility for authorship during the submission and review stages of the manuscript.

## Conflict of Interest

Istanbul Journal of Pharmacy requires and encourages

the authors and the individuals involved in the evaluation process of submitted manuscripts to disclose any existing or potential conflicts of interests, including financial, consultant, and institutional, that might lead to potential bias or a conflict of interest. Any financial grants or other support received for a submitted study from individuals or institutions should be disclosed to the Editorial Board. To disclose a potential conflict of interest, the ICMJE Potential Conflict of Interest Disclosure Form should be filled in and submitted by all contributing authors. Cases of a potential conflict of interest of the editors, authors, or reviewers are resolved by the journal's Editorial Board within the scope of COPE and ICMJE guidelines.

The Editorial Board of the journal handles all appeal and complaint cases within the scope of COPE guidelines. In such cases, authors should get in direct contact with the editorial office regarding their appeals and complaints. When needed, an ombudsperson may be assigned to resolve cases that cannot be resolved internally. The Editor in Chief is the final authority in the decision-making process for all appeals and complaints.

## Copyright and Licensing

Istanbul Journal of Pharmacy requires each submission to be accompanied by a Copyright Agreement Form (available for download at [http://ijp.istanbul.edu.tr/en/\\_](http://ijp.istanbul.edu.tr/en/_)). When using previously published content, including figures, tables, or any other material in both print and electronic formats, authors must obtain permission from the copyright holder. Legal, financial and criminal liabilities in this regard belong to the author(s). By signing the Copyright Agreement Form, authors agree that the article, if accepted for publication by the Istanbul Journal of Pharmacy, will be licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC-BY-NC).

## Disclaimer

Statements or opinions expressed in the manuscripts published in Istanbul Journal of Pharmacy reflect the views of the author(s) and not the opinions of the editors, the editorial board, or the publisher; the editors, the editorial board, and the publisher disclaim any responsibility or liability for such materials. The final responsibility in regard to the published content rests with the authors.



## MANUSCRIPT PREPARATION

The manuscripts should be prepared in accordance with ICMJE-Recommendations for the Conduct, Reporting, Editing, and Publication of Scholarly Work in Medical Journals (updated in December 2015 - <http://www.icmje.org/icmje-recommendations.pdf>). Authors are required to prepare manuscripts in accordance with the CONSORT guidelines for randomized research studies, STROBE guidelines for observational original research studies, STARD guidelines for studies on diagnostic accuracy, PRISMA guidelines for systematic reviews and meta-analysis, ARRIVE guidelines for experimental animal studies, and TREND guidelines for non-randomized public behavior.

Manuscripts can only be submitted through the journal's online manuscript submission and evaluation system, available at [http://ijp.istanbul.edu.tr/en/\\_](http://ijp.istanbul.edu.tr/en/_). Manuscripts submitted via any other medium will not be evaluated.

Manuscripts submitted to the journal will first go through a technical evaluation process where the editorial office staff will ensure that the manuscript has been prepared and submitted in accordance with the journal's guidelines. Submissions that do not conform to the journal's guidelines will be returned to the submitting author with technical correction requests.

Authors are required to submit the following:

- Copyright Agreement Form
- Author Form
- Title Page

during the initial submission.

The manuscript should be prepared in MS Word format by using Times New Roman font (12 pt) and double-spaced on one side of the paper with adequate margins (2.5 cm).

### Preparation of the Manuscript

**Title page:** A separate title page should be submitted with all submissions and this page should include:

- The full title of the manuscript as well as a short title (running head) of no more than 50 characters,

- Name(s), affiliations, and highest academic degree(s) and ORCID ID(s) of the author(s),
- Grant information and detailed information on the other sources of support,
- Name, address, telephone (including the mobile phone number) and fax numbers, and email address of the corresponding author,
- Acknowledgment of the individuals who contributed to the preparation of the manuscript but who do not fulfill the authorship criteria.

**Abstract:** An structured abstract should be submitted with Original Articles (Background and Aims, Methods, Results, Conclusion). Please check Table 1 below for word count specifications.

**Keywords:** Each submission must be accompanied by a minimum of three to a maximum of six keywords for subject indexing at the end of the abstract. The keywords should be listed in full without abbreviations. The keywords should be selected from the National Library of Medicine, Medical Subject Headings database (<https://www.nlm.nih.gov/mesh/MBrowser.html>).

### Manuscript Types

**Original Articles:** This is the most important type of article since it provides new information based on original research. The main text of original articles should be structured with Introduction, Materials and Methods, Results, Discussion, and Conclusion subheadings. Results and Discussion sections can be combined under "Result and Discussion" heading. Please check Table 1 for the limitations for Original Articles.

Statistical analysis to support conclusions is usually necessary. Statistical analyses must be conducted in accordance with international statistical reporting standards (Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. *Br Med J* 1983; 7; 1489-93). Information on statistical analyses with specified statistical software and descriptive details of the chemical used should be provided with a separate subheading under the Materials and Methods section.

Units should be prepared in accordance with the International System of Units (SI).



**Editorial Comments:** Editorial comments aim to provide a brief critical commentary by reviewers with expertise or with high reputation in the topic of the research article published in the journal. Authors are selected and invited by the journal to provide such comments. Abstract, Keywords, and Tables, Figures, Images, and other media are not included.

**Review Articles:** Reviews prepared by authors who have extensive knowledge on a particular field and whose scientific background has been translated into a high volume of publications with a high citation potential are welcomed. These authors may even be invited by the journal. Reviews should describe, discuss, and evaluate the current level of knowledge of a topic in clinical practice and should guide future studies. Please check Table 1 for the limitations for Review Articles.

**Short Papers:** Please check Table 1 for the limitations for Short Papers.

**Letters to the Editor:** This type of manuscript discusses important parts, overlooked aspects, or lacking parts of a previously published article. Articles on subjects within the scope of the journal that might attract the readers' attention, particularly educative cases, may also be submitted in the form of a "Letter to the Editor." Readers can also present their comments on the published manuscripts in the form of a "Letter to the Editor." Abstract, Keywords, and Tables, Figures, Images, and other media should not be included. The text should be unstructured. The manuscript that is being commented on must be properly cited within this manuscript.

## Tables

Tables should be included in the main document, pre-

sented after the reference list, and they should be numbered consecutively in the order they are referred to within the main text. A descriptive title must be placed above the tables. Abbreviations used in the tables should be defined below the tables by footnotes (even if they are defined within the main text). Tables should be created using the "insert table" command of the word processing software and they should be arranged clearly to provide easy reading. Data presented in the tables should not be a repetition of the data presented within the main text but should be supporting the main text.

## Figures and Figure Legends

Figures, graphics, and photographs should be submitted as separate files (in TIFF or JPEG format) through the submission system. The files should not be embedded in a Word document or the main document. When there are figure subunits, the subunits should not be merged to form a single image. Each subunit should be submitted separately through the submission system. Images should not be labeled (a, b, c, etc.) to indicate figure subunits. Thick and thin arrows, arrowheads, stars, asterisks, and similar marks can be used on the images to support figure legends. Like the rest of the submission, the figures too should be blind. Any information within the images that may indicate an individual or institution should be blinded. The minimum resolution of each submitted figure should be 300 DPI. To prevent delays in the evaluation process, all submitted figures should be clear in resolution and large in size (minimum dimensions: 100 × 100 mm). Figure legends should be listed at the end of the main document.

All acronyms, abbreviations, and symbols used in the manuscript must follow international rules and should be defined at first use, both in the abstract and in the

**Table 1. Limitations for each manuscript type**

Type of manuscript	Word limit	Abstract word limit	Table limit	Figure limit
Original Article	3500	250 (Structured)	6	7 or total of 15 images
Review Article	5000	250 (Unstructured)	6	10 or total of 20 images
Short Paper	1000	200	No tables	10 or total of 20 images
Letter to the Editor	500	No abstract	No tables	No media



# Istanbul Journal of Pharmacy

main text. The abbreviation should be provided in parentheses following the definition.

For plant materials, herbarium name (or acronym), number, name and surname of the person who identified the plant materials should be indicated in the Materials and Methods section of the manuscript.

When a drug, product, hardware, or software program is mentioned within the main text, product information, including the name of the product, the producer of the product, and city and the country of the company (including the state if in USA), should be provided in parentheses in the following format: "Discovery St PET/CT scanner (General Electric, Milwaukee, WI, USA)"

All references, tables, and figures should be referred to within the main text, and they should be numbered consecutively in the order they are referred to within the main text. Limitations, drawbacks, and the shortcomings of original articles should be mentioned in the Discussion section before the conclusion paragraph.

## REFERENCES

### Reference Style and Format

Istanbul Journal of Pharmacy complies with APA (American Psychological Association) style 6th Edition for referencing and quoting. For more information:

- American Psychological Association. (2010). Publication manual of the American Psychological Association (6<sup>th</sup> ed.). Washington, DC: APA.
- <http://www.apastyle.org>

Accuracy of citation is the author's responsibility. All references should be cited in text. Reference list must be in alphabetical order. Type references in the style shown below

### Citations in the Text

Citations must be indicated with the author surname and publication year within the parenthesis.

If more than one citation is made within the same parenthesis, separate them with (;).

### Samples:

#### **More than one citation;**

(Esin et al., 2002; Karasar, 1995)

#### **Citation with one author;**

(Akyolcu, 2007)

#### **Citation with two authors;**

(Sayiner & Demirci, 2007)

#### **Citation with three, four, five authors;**

First citation in the text: (Ailen, Ciambune, & Welch, 2000) Subsequent citations in the text: (Ailen et al., 2000)

#### **Citations with more than six authors;**

(Çavdar et al., 2003)

### Citations in the Reference

All the citations done in the text should be listed in the References section in alphabetical order of author surname without numbering. Below given examples should be considered in citing the references.

### Basic Reference Types

#### **Book**

##### **a) Turkish Book**

Karasar, N. (1995). *Araştırmalarda rapor hazırlama* (8<sup>th</sup> ed.) [Preparing research reports]. Ankara, Turkey: 3A Eğitim Danışmanlık Ltd.

##### **b) Book Translated into Turkish**

Mucchielli, A. (1991). *Zihniyetler* [Mindsets] (A. Kotil, Trans.). İstanbul, Turkey: İletişim Yayınları.

##### **c) Edited Book**

Ören, T., Üney, T., & Çölkesen, R. (Eds.). (2006). *Türkiye bilişim ansiklopedisi* [Turkish Encyclopedia of Informatics]. İstanbul, Turkey: Papatya Yayıncılık.

##### **d) Turkish Book with Multiple Authors**

Tonta, Y., Bitirim, Y., & Sever, H. (2002). *Türkçe arama motorlarında performans değerlendirme* [Performance evaluation in Turkish search engines]. Ankara, Turkey: Total Bilişim.

##### **e) Book in English**

Kamien R., & Kamien A. (2014). *Music: An appreciation*. New York, NY: McGraw-Hill Education.

##### **f) Chapter in an Edited Book**

Bassett, C. (2006). Cultural studies and new media. In G. Hall & C. Birchall (Eds.), *New cultural studies: Adventures in theory* (pp. 220-237). Edinburgh, UK: Edinburgh University Press.



## **g) Chapter in an Edited Book in Turkish**

Erkmen, T. (2012). Örgüt kültürü: Fonksiyonları, öğeleri, işletme yönetimi ve liderlikteki önemi [Organization culture: Its functions, elements and importance in leadership and business management]. In M. Zencirkıran (Ed.), *Örgüt sosyolojisi* [Organization sociology] (pp. 233–263). Bursa, Turkey: Dora Basım Yayın.

## **h) Book with the same organization as author and publisher**

American Psychological Association. (2009). *Publication manual of the American psychological association* (6<sup>th</sup> ed.). Washington, DC: Author.

## **Article**

### **a) Turkish Article**

Mutlu, B., & Savaşer, S. (2007). Çocuğu ameliyat sonrası yoğun bakımda olan ebeveynlerde stres nedenleri ve azaltma girişimleri [Source and intervention reduction of stress for parents whose children are in intensive care unit after surgery]. *Istanbul University Florence Nightingale Journal of Nursing*, 15(60), 179–182.

### **b) English Article**

de Cillia, R., Reisigl, M., & Wodak, R. (1999). The discursive construction of national identity. *Discourse and Society*, 10(2), 149–173. <http://dx.doi.org/10.1177/0957926599010002002>

### **c) Journal Article with DOI and More Than Seven Authors**

Lal, H., Cunningham, A. L., Godeaux, O., Chlibek, R., Diez-Domingo, J., Hwang, S.-J. ... Heineman, T. C. (2015). Efficacy of an adjuvanted herpes zoster subunit vaccine in older adults. *New England Journal of Medicine*, 372, 2087–2096. <http://dx.doi.org/10.1056/NEJMoa1501184>

### **d) Journal Article from Web, without DOI**

Sidani, S. (2003). Enhancing the evaluation of nursing care effectiveness. *Canadian Journal of Nursing Research*, 35(3), 26–38. Retrieved from <http://cjr.mcgill.ca>

### **e) Journal Article with DOI**

Turner, S. J. (2010). Website statistics 2.0: Using Google Analytics to measure library website effectiveness. *Technical Services Quarterly*, 27, 261–278. <http://dx.doi.org/10.1080/07317131003765910>

## **f) Advance Online Publication**

Smith, J. A. (2010). Citing advance online publication: A review. *Journal of Psychology*. Advance online publication. <http://dx.doi.org/10.1037/a45d7867>

## **g) Article in a Magazine**

Henry, W. A., III. (1990, April 9). Making the grade in today's schools. *Time*, 135, 28–31.

## **Doctoral Dissertation, Master's Thesis, Presentation, Proceeding**

### **a) Dissertation/Thesis from a Commercial Database**

Van Brunt, D. (1997). *Networked consumer health information systems* [Doctoral dissertation]. Available from ProQuest Dissertations and Theses database. (UMI No. 9943436)

### **b) Dissertation/Thesis from an Institutional Database**

Yaylı-Yıldız, B. (2014). *University campuses as places of potential publicness: Exploring the political, social and cultural practices in Ege University* [Doctoral dissertation]. Retrieved from <http://library.iyte.edu.tr/hizli-erisim/iyte-tez-portali>

### **c) Dissertation/Thesis from Web**

Tonta, Y. A. (1992). *An analysis of search failures in online library catalogs* [Doctoral dissertation, University of California, Berkeley]. Retrieved from <http://yunus.hacettepe.edu.tr/~tonta/yayinlar/phd/ickapak.html>

### **d) Dissertation/Thesis abstracted in Dissertations Abstracts International**

Appelbaum, L. G. (2005). Three studies of human information processing: Texture amplification, motion representation, and figure-ground segregation. *Dissertation Abstracts International: Section B. Sciences and Engineering*, 65(10), 5428.

### **e) Symposium Contribution**

Krinsky-McHale, S. J., Zigman, W. B., & Silverman, W. (2012, August). Are neuropsychiatric symptoms markers of prodromal Alzheimer's disease in adults with Down syndrome? In W. B. Zigman (Chair), *Predictors of mild cognitive impairment, dementia, and mortality in adults with Down syndrome*. Symposium conducted at the meeting of the American Psychological Association, Orlando, FL.

### **f) Conference Paper Abstract Retrieved Online**

Liu, S. (2005, May). *Defending against business crises with the help of intelligent agent based early warning solutions*. Paper presented at the Seventh



International Conference on Enterprise Information Systems, Miami, FL. Abstract retrieved from [http://www.iceis.org/iceis2005/abstracts\\_2005.htm](http://www.iceis.org/iceis2005/abstracts_2005.htm)

**g) Conference Paper - In Regularly Published Proceedings and Retrieved Online**

Herculano-Houzel, S., Collins, C. E., Wong, P., Kaas, J. H., & Lent, R. (2008). The basic nonuniformity of the cerebral cortex. *Proceedings of the National Academy of Sciences*, 105, 12593–12598. <http://dx.doi.org/10.1073/pnas.0805417105>

**h) Proceeding in Book Form**

Parsons, O. A., Pryzwansky, W. B., Weinstein, D. J., & Wiens, A. N. (1995). Taxonomy for psychology. In J. N. Reich, H. Sands, & A. N. Wiens (Eds.), *Education and training beyond the doctoral degree: Proceedings of the American Psychological Association National Conference on Postdoctoral Education and Training in Psychology* (pp. 45–50). Washington, DC: American Psychological Association.

**i) Paper Presentation**

Nguyen, C. A. (2012, August). *Humor and deception in advertising: When laughter may not be the best medicine*. Paper presented at the meeting of the American Psychological Association, Orlando, FL.

**Other Sources**

**a) Newspaper Article**

Browne, R. (2010, March 21). This brainless patient is no dummy. *Sydney Morning Herald*, 45.

**b) Newspaper Article with no Author**

New drug appears to sharply cut risk of death from heart failure. (1993, July 15). *The Washington Post*, p. A12.

**c) Web Page/Blog Post**

Bordwell, D. (2013, June 18). David Koepp: Making the world movie-sized [Web log post]. Retrieved from <http://www.davidbordwell.net/blog/page/27/>

**d) Online Encyclopedia/Dictionary**

Ignition. (1989). In *Oxford English online dictionary* (2<sup>nd</sup> ed.). Retrieved from <http://dictionary.oed.com>

Marcoux, A. (2008). Business ethics. In E. N. Zalta (Ed.). *The Stanford encyclopedia of philosophy*. Retrieved from <http://plato.stanford.edu/entries/ethics-business/>

**e) Podcast**

Dunning, B. (Producer). (2011, January 12). *inFact: Conspiracy theories* [Video podcast]. Retrieved from <http://itunes.apple.com/>

**f) Single Episode in a Television Series**

Egan, D. (Writer), & Alexander, J. (Director). (2005). Failure to communicate. [Television series episode]. In D. Shore (Executive producer), *House*; New York, NY: Fox Broadcasting.

**g) Music**

Fuchs, G. (2004). Light the menorah. On *Eight nights of Hanukkah* [CD]. Brick, NJ: Kid Kosher.

**REVISIONS**

When submitting a revised version of a paper, the author must submit a detailed “Response to the reviewers” that states point by point how each issue raised by the reviewers has been covered and where it can be found (each reviewer’s comment, followed by the author’s reply and line numbers where the changes have been made) as well as an annotated copy of the main document. Revised manuscripts must be submitted within 30 days from the date of the decision letter. If the revised version of the manuscript is not submitted within the allocated time, the revision option may be cancelled. If the submitting author(s) believe that additional time is required, they should request this extension before the initial 30-day period is over.

Accepted manuscripts are copy-edited for grammar, punctuation, and format. Once the publication process of a manuscript is completed, it is published online on the journal’s webpage as an ahead-of-print publication before it is included in its scheduled issue. A PDF proof of the accepted manuscript is sent to the corresponding author and their publication approval is requested within 2 days of their receipt of the proof.

**Editor in Chief:** Emine AKALIN

**Address:** İstanbul University Faculty of Pharmacy, İstanbul, Turkey

**Phone:** +90 212 440 02 75

**Fax:** +90 212 440 02 52

**E-mail:** [jfacpharm@istanbul.edu.tr](mailto:jfacpharm@istanbul.edu.tr)

**Publisher:** Istanbul University Press

**Address:** İstanbul University Central Campus, 34452 Beyazıt, Fatih / İstanbul - Turkey

**Phone:** +90 212 440 00 00



## CONTENTS

### ORIGINAL ARTICLES

- Evaluation of drug-drug interactions and their clinical importance in a pediatric hematopoietic stem cell transplantation unit ..... 1  
Nesligül Özdemir, Ayçe Çeliker, Bülent Barış Kuşkonmaz, Duygu Uçkan Çetinkaya
- Alpha lipoic acid bioequivalence study redesigned: a candidate for highly variable drugs ..... 8  
Burcu Bulut, Nagehan Sarraçoğlu, Onur Pinarbaşı
- Development and validation an HPLC - UV method for determination of esomeprazole and pirfenidone simultaneously in rat plasma: application to a drug monitoring study ..... 16  
Emrah Dural, Sema Tülay Köz, Süleyman Köz
- Preparation and *in vitro* studies of fixed-dose tablet combination of repaglinide and metformin ..... 26  
Sinem Şahin, Burcu Mesut, M. Ezgi Durgun, Esher Özçelik, Yıldız Özsoy
- The formulation and characterization of water-soluble snakehead fish (*Ophiocephalus striatus*) dry extract in nanoemulsion using permeation and *in vivo* study ..... 35  
Robert Tungadi, Widysusanti Abdulkadir, Munafri Tahir
- Pseudo ternary phase diagrams: a practical approach for the area and centroid calculation of stable microemulsion regions ..... 42  
Murat Sami Berkman, Kadri Güleç
- Rational design and synthesis of quinazoline derivatives ..... 50  
Fatimah AlKharaz, Zead Abudayeh, Qais Abualassal, Loay Hassouneh
- Molecular and crystal structure of 1-methyl-5-trifluoromethoxy-1*H*-indole-2,3-dione 3-[4-(4-methoxyphenyl)thiosemicarbazone] ..... 59  
Özge Soylu Eter, Zeliha Atioğlu, Mehmet Akkurt, Cem Cüneyt Ersanlı, Nilgün Karalı
- Synthesis, characterization and antibacterial evaluation of new pyridyl-thiazole hybrids of sulfonamides ..... 67  
Zafer Şahin, Sevede Nur Biltekin, Leyla Yurttaş, Şeref Demirayak
- The effects of vanadyl sulfate on glutathione, lipid peroxidation and nonenzymatic glycosylation levels in various tissues in experimental diabetes ..... 73  
Sevim Tunalı, Refiye Yanardağ
- Short-term adaptive metabolic response of *Escherichia coli* to ciprofloxacin exposure ..... 79  
Engin Koçak, Ceren Özkul
- In vitro* urease and trypsin inhibitory activities of some sulfur compounds ..... 85  
Eda Dağsuyu, Refiye Yanardağ
- Quality of turmeric powder in herbal stores: pharmacognostical investigations on turmeric powders obtained from herbal stores in Istanbul, Turkey ..... 92  
Ebru Kuruldak, Fatmanur Yılmaz, Gizem Gülsoy Toplan, Berna Özbek Çelik, Afife Mat



## CONTENTS

### ORIGINAL ARTICLES

- Isolation of potential liver x receptor alpha agonist and antioxidant compounds from *Hypericum microcalycinum* Boiss. & Heldr. ....98  
Seçil Sarıkaya Aydın, Vahap Murat Kutluay, Toshiaki Makino, Makoto Inoue, Ümmühan Şebnem Harput, İclal Saraçoğlu
- In vitro* evaluation of antimicrobial activity of *Distemonanthus benthamianus* chewing stick extract mouthwash..... 105  
Mbang Nyong Femi-Oyewo, Olutayo Ademola Adeleye, Caroline Olufunke Babalola, Olufemi Babatunde Banjo, Modupe Nofisat Adebawale, Florence Olubola Odeleye
- Phytochemical screening, phenolic content and antioxidant activity of *Lavandula* species extracts from Algeria.....111  
Farah Haddouchi, Tarik Mohammed Chaouche, Meriem Saker, Imane Ghellai, Ouhiba Boudjemai
- Antimicrobial activities of some narrow endemic gypsophyte.....118  
Esma Ocak, Şule İnci, Derviş Öztürk, Sanem Akdeniz Şafak, Ebru Özdeniz, Sevda Kırbağ, Ahmet Harun Evren, Latif Kurt
- Chelidonium majus* L. (Papaveraceae) morphology, anatomy and traditional medicinal uses in Turkey.....123  
Golshan Zare, Neziha Yağmur Diker, Zekiye Ceren Arıtuluk, İffet İrem Tatlı Çankaya
- The anatomical properties of endemic *Hypericum kotschyanum* Boiss. ....133  
Onur Altınbaşak, Gülay Ecevit Genç, Şükran Kültür

### REVIEW ARTICLES

- Fungal homoserine transacetylase: A potential antifungal target .....137  
Esra Seyran
- Bioavailability of berberine: challenges and solutions.....141  
Asha Thomas, Seema Kamble, Sanjeevani Deshkar, Lata Kothapalli, Sohan Chitlange
- Evaluation of toxic effects of statins and their possible role in treatment of cancer .....154  
Aysun Ökçesiz, Ülkü Ündeğer Bucurgat

# The formulation and characterization of water-soluble snakehead fish (*Ophiocephalus striatus*) dry extract in nanoemulsion using permeation and *in vivo* study

Robert Tungadi<sup>1</sup> , Widysusanti Abdulkadir<sup>1</sup> , Munafri Tahir<sup>1</sup> 

<sup>1</sup>State University of Gorontalo, Faculty of Sport and Health, Department of Pharmacy, Gorontalo, Indonesia

**ORCID IDs of the authors:** R.T. 0000-0003-2141-2402; W.A. 0000-0002-8975-134X; M.T. 0000-0002-9351-2843

**Cite this article as:** Tungadi, R., Abdulkadir, W., & Tahir, M. (2021). The formulation and characterization of water-soluble snakehead fish (*Ophiocephalus striatus*) dry extract in nanoemulsion using permeation and *in vivo* study. *Istanbul Journal of Pharmacy*, 51(1), 35-41.

## ABSTRACT

**Background and Aims:** The study was conducted to determine the optimal concentration of water-soluble snakehead fish dry extract (SFDE) in nanoemulsion and the amount of albumin required to penetrate the skin in order to accelerate the wound healing process.

**Methods:** The snakehead fish (SF) was extracted using an atomizer while the nanoemulsion basis was optimized using oleic acid, Tween 80, and propylene glycol. The developed SFDE in nanoemulsion was characterized based on droplet size, PDI, and zeta potential. The ability of the mixture to penetrate the snakeskin was tested using Franz diffusion cells. The effectiveness of the nanoemulsion was evaluated by dividing the rabbits used for experiment into 6 treatment groups including SFDE F1 0.25%, F2 0.5%, F3 1%, F4 SF 2% cream, F5 nanoemulsion basis, and F6 no treatment.

**Results:** The SFDE nanoemulsion produced a particle size of 147.5 nm with acceptable PDI (0.23) and zeta potential (+13.38 mV). The most effective SFDE to accelerate the healing of open wounds in rabbits was a concentration of 1%, which was found to have dried and closed the wound on the 3rd day.

**Conclusion:** The permeation study and the effectiveness test showed the 1% SFDE nanoemulsion is the best concentration in accelerating the wound healing process and ensuring the highest albumin penetration into the skin.

**Keywords:** Snakehead fish, nanoemulsion, albumin, wound, water-soluble, rabbit

## INTRODUCTION

Snakehead fish (SF) (*Ophiocephalus striatus*) is an economically valuable fish widely used for processed products. According to Suprayitno (2003), it has a protein content estimated to be 25.1% compared to the 6.224% found in albumin and is higher than the values obtained from other animal sources used for patients with hypoalbuminemia (i.e. low albumin) and wounds. This is important because albumin has been discovered in medical science to have the ability of accelerating the recovery of broken body cell tissues due to surgery (Suprayitno, 2003; Ulandari, Kurniawan, & Putri, 2011).

Albumin is the largest type of protein in plasma with 60% content and also has the ability to synergize with zinc mineral needed for the development and formation of new cell tissues in wounds. Zinc has been reported to have the ability to functions as an antioxidant to protect cells, accelerate the wound healing process, and regulate expression of lymphocytes and proteins (Mustafa, Widodo, & Kristianto, 2012; Maryanto, 2004). Moreover, the chemical compounds of Snakehead fish dry extract (SFDE), including albumin and amino acids (glycine and lysine), have been discovered to be soluble in water based on chemical analysis tests from

## Address for Correspondence:

Robert TUNGADI, e-mail: robert.tungadi@ung.ac.id

Submitted: 10.05.2020  
Revision Requested: 06.07.2020  
Last Revision Received: 02.09.2020  
Accepted: 21.09.2020

This work is licensed under a Creative Commons Attribution 4.0 International License.



LIPI conducted using spectrophotometry and HPLC methods. It is, therefore, the mix of these elements with nanoemulsion (NE) that is needed to obtain a homogenous system through the emulsification method (Zhang, Zhang, Fan, Liu, & Meng, 2019).

According to Tungadi, Susanty, Wicita, & Pido (2018), snake-head fish with 2% cream was found to have accelerated the healing process of rabbit skin's open wound in an *in vivo* study, but the cream was observed to be physically unstable after 3 months of storage. This was associated with the mixture of snakehead fish dry extract (SFDE) with macro emulsion, which causes effortless damage due to the strength of oil and water phase and storage temperature. A solution has, however, been reported which involves reducing the particle size of snake-head fish dry extract and stabilizer using nanoemulsion formulation through appropriate utilization of surface-active agents, co-surfactant, and oil (Tungadi, 2011; Devarajan, & Ravichandran, 2011). It is also possible to formulate the SFDE into the emulsion because it contains hydrophilic and hydrophobic compounds with the nanoemulsion discovered to be useful for transdermal drug delivery such as the penetration of active compounds due to stratum corneum deformability (Tungadi, Susanty, Wicita, & Pido, 2018).

Meanwhile, Tungadi, R. (2016) showed snakehead fish cream containing only 50% albumin has the ability to penetrate the skin membrane using penetrant enhancers such as propylene glycol. This, according to an *in vivo* study, has been reported to accelerate the healing of open wounds (treatment group) due to the increase in the rate of diffusing albumin into the stratum corneum. However, a low percentage of albumin is produced without the use of a penetrant enhancer (Tungadi & Hasan, 2016).

This shows a nanoemulsion system is suitable for the drug delivery through the skin due to its large surface area, which makes the penetration of active substances faster. It is also useful because its manufacturing process is very easy and efficient (Chuesiang, Siripatrawan, Sanguandeekul, McLandsborough, & McClements, 2018; Laxmi, Bhardwaj, Mehta, & Mehta, 2015) as observed in the formation of SFDE into dosage forms in Winda's research. This involved the optimization of nanoemulsion basis as a carrier for SF nanoemulsion preparation and later characterization by particle size, polydispersity index, and zeta potential with the results found to be 147.5 nm, 0.234, and +13.38 mV respectively (Tungadi, Moo, & Mozin, 2017). Therefore, this current study was conducted to determine the effectiveness of different concentrations of SFDE at 0.25, 0.5, and 1% in accelerating the healing of open wounds on rabbits dorsum and the amount of albumin required to penetrate their skin using the Franz diffusion cell.

## MATERIALS AND METHODS

### Materials

Snakehead fish dry extract was obtained from PT. Ismut Medical Pharmaceutical, Indonesia. The Rabbits were purchased from the animal market. The nanoemulsion basis (tween 80, propylene glycol, and oleic acid) was purchased from PT. Brataco Chemical. Other materials, such as propylparaben,

methylparaben, isopropyl myristate, lanolin, cetyl alcohol, paraffin liquid, and BHT were purchased from PT. Sentana Chemical. A UV-Vis Spectrophotometer (USA), Delsa™ Nano (UK), pH meter (Systronics model EQMK), sonicator (Specta Lab), hot air oven (Mettler), and the Franz diffusion cell (Intalab) were used.

Albino rabbits (2 kg) were obtained from the animal laboratory center of LIPI, Serpong, Indonesia. The experimental procedure was conducted according to the Institutional Animal Ethics Committee based on the recommendations of the Health Ethics Committee, The Faculty of Medicine, Hasanuddin University, Indonesia Government with registration No. UH08060042

### The optimization and characterization of SFDE nanoemulsion basis

The nanoemulsion basis was optimized by comparing different concentrations of oil (oleic acid), co-surfactant (propylene glycol), and surfactant (Tween 80) using five formulas including F1 (1:2:4), F2 (1:3:4), F3 (1:3:5), F4 (1:3:6), and F5 (1:3:7). The Tween 80 and propylene glycol were mixed collectively using a magnetic stirrer for 30 minutes at 250 rpm. For the first mixture, the oleic acid was introduced during the stirring process. Water containing 0.25%, 0.5%, and 1% of SFDE was added drop by drop then other adjuvants, such as methylparabens and propylparabens (preservatives) as well as BHT (antioxidant) were added. After that, sonication of the mixtures at 20 KHz was performed for 10 minutes at 25°C to complete the process. The same procedure was performed for all the formulations with different concentrations of Tween 80, propylene glycol, and oleic acid. All formulations were characterized using a particle size analyzer to measure the size of droplets, zeta potential, and PDI.

### Permeation study

*In vitro* permeation, conducted using Franz diffusion cell, has been described as a dependable technique to predict the transport of drugs in the skin (Zhu et al., 2009) and, for this study, an excised python skin (*Python reticulatus*) was used.

This process involved the separation of the skin from abundant fats and the elimination of connective tissue using a scalpel. The excised skin was washed with NaCl 0.9% and examined for integrity before it was hooked up on the diffusion cell with an effective diffusion area. Moreover, the stratum corneum facet was focused on the donor while the dermal layer was on the receiver compartment consisting of 47 ml phosphate buffer of pH 7.4 as the receptor fluid agitated at 100 rpm and maintained at 37±0.5°C during the experiments with 1 g of the nanoemulsion used in every diffusion cell. Approximately 2 ml of the samples were withdrawn for evaluation at 0, 30, 60, 90, 120, 150, 180, 210, and 240 min after the experiment has commenced and changed immediately with an equal volume of fresh diffusion medium (Tungadi et al., 2018).

### Skin irritation study

Skin inflammation was evaluated using 12 healthy rabbits without any injuries or skin disorders. They were grouped into three with n=3 of albino male rabbits weighing 1.5-2 kg; positive control (2% w/w SFDE, commercial product), and negative control (nanoemulsion basis) also with n=3 on the 2 cm<sup>2</sup> dor-

sal facet of the shaven skin of the rabbits. The treatment was eliminated after 72 h to check for any symptoms of erythema and edema (Tungadi et al., 2018; Barot, Parejiya, Patel, Mehta, & Shelat, 2012; Lala, & Awari, 2014). Undesirable skin changes such as coloration and morphology were examined at 1h, 24 h, 48 h, and 72 h intervals. The reactions obtained were recorded and compared with a control group (n=3).

### Effectiveness of the SFDE *in vivo* study

#### Preparation and grouping of test animals

The implementation stage started with the preparation of 12 male white rabbits randomly divided into 6 groups of treatments, each consisting of 2 rabbits, each of which were placed in individual cages and acclimated for 5 days. The Treatment Group contained SFDE varied at G1 0.25%, G2 0.5% and G3 1% of SFDE.

#### Testing of SFDE on test animals

The dorsal back of each test animal was shaved and cleaned with 70% alcohol after which they were locally anesthetized with 0.2 mL lidocaine and the wounds created by slicing off 4 cm<sup>2</sup> of skin and smearing the wounds with the SFDE treatments. The average change in length and the condition of the wounds were observed and documented every day for 10 days.

#### Measurement of the open wound area

The average length of the open wound was calculated using a ruler while pictures were also taken from day 0 to 10 to determine the healing process. The values measured in each day were converted to amount of contraction to determine the reduction effect of SFDE in different concentrations.

#### Statistical analysis

All the experimental measurements were recorded in triplicate and the final values were expressed as mean value±standard deviation (SD). The statistical evaluation of the permeation *in vitro* for the predetermined intervals was conducted using One-way ANOVA SPSS 16 with a degree of significance of P cost <0.05\* and <0.01\*\*.

## RESULTS AND DISCUSSION

### The formulation and optimization of nanoemulsion basis

There are several challenges to the application of nanoemulsion as a transdermal system to successfully deliver drugs via the skin (Kong, Chen, Kweon, & Park, 2011) and some of the important ones include the small particle-sized formulation and rheology properties. Therefore, it is necessary to understand the best formula to improve the introduction of snakehead fish dry extract (SFDE) into nanoemulsion using appropriate oil, surfactant, and co-surfactant (Tungadi et al., 2018).

The best optimization for nanoemulsion basis was found to be Formula 5 (F5) with oleic acid, tween eighty and propylene glycol (1:10) based on its viscosity, clarity, and stability as shown in Table 1.

Formula 5 was also observed to be physically stable by not segregating after being centrifuged at 3800 rpm for 5 hours while Formulas 1 to 4 produced a cloudy appearance and segregated. The stability was associated with the use of Tween 80 as a nonionic surfactant considering its excessive hydrophilic and lipophilic balance estimated at 15 which made it steady in an emulsion formulation with oil in water (Brandelero, Yamashita, & Grossmann, 2010).

Surfactant plays important roles in the nanoemulsion basis due to the fact it has a large surface area to decrease interfacial and surface tension, which further leads to its absorption in the interface phase. This means it has the ability to reduce the surface free energy by disintegrating a globule into smaller parts (Natalia, 2012). However, most surfactants are unable to decrease interfacial tension in the emulsion. Therefore, there is a need to add co-surfactant such a propylene glycol to improve the solubility of nonpolar agencies (Swarbrick, 2007), intensify the flexibility of surfactant film and fluidity of the emulsion phase to shield compounds from adverse environmental conditions, and enhance their balance (Madene, Jacquot, Scher, & Desobry, 2006; Kumar, Bishnoi, Shukla, & Jain, 2019).

**Table 1. The optimization of nanoemulsion basis.**

Materials	Formula %				
	F1	F2	F3	F4	F5
	1:2:4	1:3:4	1:3:5	1:3:6	1:3:7
Oleic acid	5	5	5	5	5
Tween 80	18	20	23	25	27.5
Propylene glycol	12	15	17	20	22.5
Distilled water	100	100	100	100	100
Observation	cloudy	cloudy	cloudy	cloudy	clear
Stability tests:					
pH	6.5±0.3	6.2±0.5	6.0±0.7	5.8±0.2	5.5±0.1
Viscosity (cP)	385.6±1.3	267.8±2.5	200.3±2.1	187.5±3.2	178.2±1.4
Transmittance (%)	75.65±1.5	82.34±0.9	87.35±1.1	90.58±1.8	98.75±0.8

### Characterization of snakehead fish nanoemulsion

Nanoemulsion systems can be used to deliver drugs through trans-mucosal and transdermal routes and this means they have the ability to effectively enhance bioavailability (Kumar et al., 2019; Rehman et al., 2017). The polydispersity index (PDI) of the SFDE produced good results in the three replications, 0.205, 0.215, and 0.284 respectively, and the 147.5 nm average droplet size shown in Table 2.

**Table 2. The characterization of Snakehead Fish Nanoemulsion.**

Sample	Particle size (nm)	Average of Size (nm)	Zeta potential (mV)	Polydispersity index (PDI)	Average of PDI
1% SFDE	111 ± 0.2	147.5 ± 0.53	+ 13.38	0.205 ± 0.1	0.23 ± 0.26
	233 ± 0.5			0.215 ± 0.2	
	98.6 ± 0.9			0.284 ± 0.5	

As shown in Table 2, the average size of the droplet of SFDE nanoemulsion was 147.5 nm showing that SFDE meets the criteria of nanostructures, which require a particle size range between 1 – 100 nm or 2 – 500 nm (Shah, Bhalodia, & Shelat, 2011). Meanwhile, the zeta potential value was +13.38 mV and this indicates it has a good degree of stability. This is associated with the standard that nanoparticles with values above or below  $\pm 30$  mV indicate a physically stable colloidal system due to their ability to ensure the magnitude of the charged particle prevents particle aggregation (Singh, & Lillard, 2014; Hadian, Sahari, & Moghimi, 2014). Meanwhile, smaller values have been reported to cause particles to aggregate and flocculate due to van der Waals attractive forces acting on them, thereby, leading to physical instability. Furthermore, the average polydispersity index was recorded to be 0.234 and this means SFDE has a uniform particle size and homogeneous dispersion because this value is below 0.25 (Winterhalter, & Lasic, 2013).


The solubility of active compounds is very important in drug formulation due to its ability to increase bioavailability through oral, topical, and parenteral formulations. SFDE contains water-soluble active compounds such as albumin and amino acids and water-insoluble ones such as polyunsaturated fatty acids, vitamins, and amino acids. This study made use of only the albumin and amino acid contents to ensure easy formulation into the nanoemulsion. Therefore, solubility is one of the important parameters to achieve the appropriate concentration of drug in systemic circulation and appropriate pharmacological response (Vemula, Lagishetty, & Lingala, 2010).

### Permeation study of SFDE in nanoemulsion

Ex vivo permeation studies were also conducted using snake-skin as the membrane and the drugs from G1 (0.25% SFDE), G2 (0.5% SFDE), G3 (1% SFDE), and G4 (2% SF cream; commercial product) were found have produced  $62.80 \pm 1.45\%$ ,  $69.30 \pm 2.34\%$ ,  $72.30 \pm 1.22\%$ , and  $50.80 \pm 0.50\%$  permeation, respectively in 4h as shown in Figure 1.

Figure 1 shows 2% SF cream had the lowest percentage of albumin permeation into the skin with approximately 50.80%

compared to all other concentrations and this is associated with the formulation of SFDE containing albumin into cream o/w to produce the big particle size in SF cream due to the macroemulsion. Its introduction to nanoemulsion produced a small particle size estimated to be 147.5 nm and water-soluble compounds with the ability to increase the loading capacity of albumin to penetrate the skin easily. This is consistent with the findings of previous research on the formulation of SFDE into



**Figure 1.** The amount of albumin penetrated into the skin in 4 h; \*P<0.05; One Way Anova Test.

liposome which showed solubility and particle size to be the most important factors to increase the loading capacity and bioavailability of drugs. SFDE into liposome was discovered to have a smaller particle size, 121 nm, compared to nanoemulsion and this led to the production of the highest entrapment efficiency of albumin recorded to be 85.75% (Tungadi, Abdulkadir, Ischak, & Rahim, 2019).

The biggest impediment to the transdermal drug transport is usually associated with the stratum corneum as observed in the 10-20  $\mu\text{m}$  thick tissue layer which has a remarkably composed lipid/protein matrix structure (Ceve, 2004). According to Tungadi (2011), a study of SFDE cream containing penetrant enhancer such as propylene glycol is expected to accelerate the wound healing process through skin permeation, but the cumulative albumin penetration into rat skin membrane was recorded to be 50%. This study found SFDE nanoemulsion to have the ability to enhance the permeation of drug through the skin as observed from the cumulative percentage of SFDE permeation of F3 which was found to be the highest with  $72.30 \pm 1.22\%$  using a snakeskin membrane while the positive control, SF cream 2%, had  $50.80 \pm 0.50\%$ . This, therefore, means nanoemulsion formulation acts as drug reservoirs in the transdermal delivery systems affecting the release of drugs from the inner to the outer phase and similarly to the skin (Tungadi

et al., 2018; Mou et al., 2008). These release mechanisms, however, rely on the composition of the network surfactant chains and the Crosslink density (Tungadi et al., 2018; Bernard, 2012). Moreover, the capacity of a drug to penetrate the skin and release the therapeutic agent effectively is affected by its affinity to diffuse out from the vehicle and permeate through the barrier (Tungadi et al., 2018; Alves, Scarrone, Santos, Pohlmann, & Guterren 2007).

In the current permeation study using Franz diffusion cell, *Phyton* snakeskin was used as a membrane to facilitate the penetration of the test substances compared to the use of extracts of stratum corneum isolated from the skins. This method was used in the study by Lin and colleagues (1992) and the permeability values in snakeskin (*Phyton molurus*) were found to be 2 to 4 times higher than in isolated stratum corneum for sodium diclofenate, theophylline, and benzoic acid (2 mg/mL or 0.2% in aqueous solution). The use of *phyton* snakeskin in studying SFDE nanoemulsion as a promoter of skin penetration for hydrophilic substances such as albumin required the consideration of the lower permeability coefficient (3.3 to 6.1 times) of these membranes for such compounds, thereby, causing an extension of the time needed for the experiments. Meanwhile, lipophilic compounds have been reported to have permeability coefficients close to those obtained from human skin membranes (0.9 to 1.8 times and 3.3 to 6.1 times) (Tungadi et al., 2019).


### Skin irritation test of snakehead fish nanoemulsion

The results from the skin irritation study including erythema and edema on the rabbit skin after 1 h, 24 h, 48 h, and 72 h post-treatment of positive control, negative control, F1, F2, and F3 are represented in Table 3. The results showed no proof of inflammation, erythema, or edema; based on visible inspection after the application of all formulations of nanoemulsion on the rabbit skin during the three days of observation. This, therefore, means they were all non-sensitizing and safe for topical use.


### Percentage of wound contraction on rabbit's skin

Based on observations, on the 3<sup>rd</sup>, 6<sup>th</sup>, and 10<sup>th</sup>-day, the open wound on the rabbit in group I (F1 0.25% SFDE) was found to have wound contraction percentages between 90% and 46% as shown in Figures 2-6 with the physical appearance marked by the presence of fibrin yarns protecting the open wound as presented in Figures 3-6. In group II (F2 0.5%), the reduction

was found to be 100% to 42% and was discovered to be drying in contrast to the observation made for group I. The results of group III (F3 1%) showed a substantial contraction from 100% to 25% compared to the negative control, which was observed to be faster. This change was characterized by the production of new granulation tissue on the side of the open wound and the fact that it was already dry on the third day. Furthermore, the positive control (F4) containing snakehead fish cream 2% had the change of wound contraction from 100% to 56%. The negative control F5 with nanoemulsion basis and F6 without treatment had the slowest healing process of approximately 15 days and a marked wound contraction exchange from 100% to 75-77% (Figures 2, 3-6).



**Figure 2.** Percentage of wound contraction on rabbit's skin \*P<0.05; \*\*P<0.01; One Way Anova Test.




**Figure 3.** The observation of wound area on the first day, F1: NE 0.25% of SFDE; F2: NE 0.5%; F3: NE 1%; F4: 2% of SF cream; F5: NE basis; F6: no treatment.


**Table 3. Skin irritation study.**

	1 h		24 h		48 h		72 h	
	Erythema	Edema	Erythema	Edema	Erythema	Edema	Erythema	Edema
G1 0.25%	0	0	0	0	0	0	0	0
G2 0.5%	0	0	0	0	0	0	0	0
G3 1%	0	0	0	0	0	0	0	0
Positive Control	0	0	0	0	0	0	0	0
Negative Control	0	0	0	0	0	0	0	0


Positive control: SF cream 2% (w/w); commercial product, negative control: nanoemulsion basis; Erythema scale: 0= none, 1=slight, 2= well-defined, 3= moderate, and 4= scar formation; Edema scale: 0= none, 1= slight, 2= well-defined, 3= moderate, and 4= severe



**Figure 4.** The observation of wound area on the third day, F1: NE 0.25% of SFDE; F2: NE 0.5%; F3: NE 1%; F4: 2% of SF cream; F5: NE basis; F6: no treatment.



**Figure 5.** The observation of wound area on the sixth day, F1: NE 0.25% of SFDE; F2: NE 0.5%; F3: NE 1%; F4: 2% of SF cream; F5: NE basis; F6: no treatment.



**Figure 6.** The observation of wound area on the ninth day, F1: NE 0.25% of SFDE; F2: NE 0.5%; F3: NE 1%; F4: 2% of SF cream; F5: NE basis; F6: no treatment.

The One-Way ANOVA analysis showed the P or Sig value was  $0.022 < 0.05$  and  $0.01$ . This means there was a significant difference between the averages of open wound contraction for all treatment and control groups. However, observation data indicated NE 1% of SFDE had a faster wound area reduction compared to 0.25% and 0.5% nanoemulsion preparations and 2% SF cream.

### Effectiveness of the SFDE *in vivo* study

F3 was found to be the best formula of SFDE nanoemulsion in this *in vivo* study functioning as a transdermal delivery system to ensure a controlled release of substances over a period and improve patient comfort during dosage preparation. Meanwhile, the small droplet size has been reported to have the ability to absorb albumin containing large molecules following the spontaneous size of the globule and surroundings (Lovelyn & Attama, 2011). The percentage of the albumin penetration and wound contraction of F3 were estimated at 72.30% and 25% on the 10<sup>th</sup> day. This was associated with the particle size and zeta potential of SFDE nanoemulsion because its small size of droplets increases the diffusion rate of albumin compared to micro or macro emulsion while the significant stability was due to the PDI and zeta potential.

The SFDE in nanoemulsion was able to accelerate the wound healing process due to the nutritional contents of snakehead fish including 0.003% Zn, 30.2% albumin, and 0.001% glycine (Mansyur, 2010) triggering the formation of Endothelial Progenitor Cells (EPC). The Zn plays a key role and has also been reported to be an important mineral in the structure and function of cell membranes by limiting the damage caused by free radicals during inflammation. Furthermore, it is also involved in the immune system, the defense of the skin, and the regulation of genes in lymphocytes (Tungadi et al., 2019; gawhirunpat, Panomsuk, Opanasopit, Rojanata, & Hatanaka, 2006; Tungadi, & Wicita, 2020).

### CONCLUSION

It is possible to formulate water-soluble snakehead fish dry extract into nanoemulsion with small particles to increase the loading capacity of albumin in penetrating the skin. The permeation study and the effectiveness test showed the 1% SFDE in nanoemulsion is the best concentration compared to others in accelerating the wound healing process and ensuring the highest albumin penetration into the skin.

**Acknowledgments:** The authors are thankful to the Ministry of Research, Technology, and Higher Education of Indonesia, which has funded this research by grant competition (decentralization grant) and are also thankful to PT. Ismut, Pharmaceuticals, Indonesia, for providing snakehead fish powder for this work and PT. NanoTech Herbal Indonesia, LIPI Serpong, Indonesia which has given technical supports. Besides that, the authors also thanks the native proofreading service (NPS) for improving the quality of this paper.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Conception/Design of Study- R.T.; Data Acquisition- R.T., W.A.; Data Analysis/Interpretation- R.T., W.A., M.T.; Drafting Manuscript- R.T., W.A., M.T.; Critical Revision of Manuscript- R.T., W.A.; Final Approval and Accountability- R.T., W.A., M.T.

**Conflict of Interest:** The authors have no conflict of interest to declare.

**Financial Disclosure:** Authors declare having received financial support from The Ministry of Research and Technology (DIKTI).

## REFERENCES

- Alves, M. P., Scarrone, A. L., Santos, M., Pohlmann, A. R., & Guterren S. S. (2007). Human skin penetration and distribution of nimesulide from hydrophilic gels. *International Journal of Pharmaceutics*, 314(1-2), 215-220.
- Barot, B. S., Parejiya, P. B., Patel, H. K., Mehta D. M., & Shelat, P. K. (2012). Microemulsion-based antifungal gel delivery to nail for the treatment of onychomycosis: Formulation, optimization, and efficacy studies. *Drug Delivery and Translational Research*, 2(6), 463-476.
- Bernard, P. B. (2012). *Modern Aspects of Emulsion Science, Emulsions-Recent Advances in Understanding*. UK: Royal Science of Chemistry.
- Brandelero, R. P. H., Yamashita, F., & Grossmann, M. V. E. (2010). The effect of surfactant Tween 80 on the hydrophilic water vapor permeation, and the mechanical properties of cassava starch and poly (butylenes adipate-co-terephthalate) (pbat) blend films. *Carbohydrate Polymers*, 82, 1102-1109.
- Ceve, G. (2004). Lipid vesicles and other colloids as drug carriers on the skin. *Advanced Drug Delivery Review*, 56(5), 675-711.
- Chuesiang, P., Siripatrawan, U., Sanguandeeukul, R., McLandsborough, L., & McClements, D. J. (2018). Optimization of cinnamon oil nanoemulsions using phase inversion temperature method: Impact of oil phase composition and surfactant concentration. *Journal of Colloid and Interface Science*, 514, 208-216.
- Devarajan, V., & Ravichandran, V. (2011). Nanoemulsions: As modified drug delivery tool. *International Journal of Comprehensive Pharmacy*, 2, 1-5.
- Hadian, Z., Sahari, M. A., & Moghimi, H. R. (2014). Formulation, characterization and optimization of liposomes containing EPA and DHA; A methodology approach. *Iranian Journal of Pharmaceutical Research*, 13(2), 393-404.
- Kong, M., Chen, X. G., Kweon, D. K., & Park, H. J. (2011). Investigation on skin hyaluronic acid based on nanoemulsion as transdermal carrier. *Carbohydrate Polymers*, 86(2), 837-843.
- Kumar, M., Bishnoi, R. S., Shukla, A. K., & Jain, P. (2019). Techniques for formulation of nanoemulsion drug delivery system: A review. *Preventive Nutrition and Food Science*, 24(3), 225-234.
- Lala, R., & Awari, N. (2014). Nanoemulsion-based gel formulations of COX-2 inhibitors for enhanced efficacy in inflammatory conditions. *Applied Nanoscience*, 4, 143-151.
- Laxmi, M., Bhardwaj, A., Mehta, S., & Mehta, A. (2015). Development and characterization of nanoemulsion as carrier for the enhancement of bioavailability of artemether. *Artificial Cells Nanomedicine and Biotechnology*, 43(5), 334-344.
- Lovelyn, C., & Attama, A. A. (2011). Current state of nanoemulsions in drug delivery. *Journal of Biomaterials and Nanobiotechnology*, 2(5), 626-639.
- Madene, A., Jacquot, M., Scher, J., & Desobry, S. (2006). Flavour encapsulation and controlled release - a review. *International Journal of Food Science and Technology*, 41, 1-21.
- Mansyur. (2010, July 15). Analysis of snakehead fish dry extract. Indonesian Institute of Sciences Biotechnology Research Center. Retrieved from <https://worldwidescience.org/topicpages/s/snakehead+fish+channa.html>
- Maryanto, A. (2004, June 18). The impact of albumin serum on length of postoperative wound healing process, Faculty of Medicine, University of Gadjah Mada. Retrieved from [http://etd.repository.ugm.ac.id/home/detail\\_pencarian/25247](http://etd.repository.ugm.ac.id/home/detail_pencarian/25247)
- Mou, D., Chen, H., Du, D., Mao, C., Wan, J., Xu, H., & Yang, X. (2008). Hydrogel thickened nanoemulsion system for topical delivery of lipophilic drugs. *International Journal of Pharmaceutics*, 353(1-2), 270-276.
- Mustafa, A., Widodo, A., & Kristianto, Y. (2012). Albumin and zinc content of snakehead fish extract and its role in health. *International Journal of Science and Technology*, 1, 1-8.
- Natalia, M. (2012). The stability and antibacterial activity test of black cumin oil (*nigella sativa* L.) nano-emulsion gel (nano-emulgel). (Master's thesis). Retrieved from <http://lib.ui.ac.id/file?file=digital/20309121-543091-Uji%20stabilitas.pdf>
- Ngawhirunpat, T., Panomsuk, S., Opanasopit, P., Rojanata, T., & Hatanaka, T. (2006). Comparison of the percutaneous absorption of hydrophilic and lipophilic compounds in shed snake skin and human skin. *Pharmazie*, 61(4), 331-335.
- Rehman, F. U., Shah, K. U., Shah, S. U., Khan, I. U., Khan, G. M., & Khan, A. (2017). From nanoemulsions to self-nanoemulsions, with recent advances in self-nanoemulsifying drug delivery systems (SNEDDS). *Expert Opinion on Drug Delivery*, 14(11), 1325-1340.
- Shah, P., Bhalodia, D., & Shelat, P. (2011). Nanoemulsion : A pharmaceutical review. *Systematic Reviews in Pharmacy*, 1, 24-32.
- Singh, R., & Lillard, J. W. (2014). Nanoparticle-based targeted drug delivery. *Experimental Molecular and Pathology*, 86(3), 215-223.
- Suprayitno, E. (2003). Snakehead Fish (*Ophiocephalus striatus*) albumin as functional food to overcome future nutrition problems. *Faculty of Fisheries, Brawijaya University*, 5(3), 32-36.
- Swarbrick, J. (2007). *Encyclopedia of pharmaceutical technology*. New York: Informa Healthcare USA Press, pp 1548-1565.
- Tungadi, R. (2011). The acceleration of wound healing of snakehead fish cream towards rabbit's skin wound histopathologically. *Indonesian Pharmaceutical Journal*, 9, 91-97.
- Tungadi, R., & Hasan, A. M. (2016). The effect of penetrant enhancer combination towards the diffusion rate of snakehead fish (*Ophiocephalus striatus*) cream in vitro and vivo. *International Journal of PharmTech Research*, 9(6), 508-13.
- Tungadi, R., Moo, D. R., & Mozin, W. R. (2017). Characterization and physical stability evaluation of snakehead fish (*Ophiocephalus striatus*) powder nanoemulsion. *International Journal of Pharmaceutical Sciences and Research*, 8(6), 2720-4.
- Tungadi, R., Susanty, W., Wicita, P., & Pido, E. (2018). Transdermal delivery of snakehead fish (*Ophiocephalus striatus*) nanoemulgel containing hydrophobic powder for burn wound. *Pharmaceutical Sciences*, 24(4), 313-323.
- Tungadi, R. (2019). Potential of snakehead fish (*Ophiocephalus striatus*) in accelerating wound healing. *Universal Journal of Pharmaceutical Research*, 4(5), 40-44.
- Tungadi, R., Abdulkadir, W., Ischak, N. I., & Rahim, B. R. (2019). Liposomal formulation of snakehead fish (*Ophiocephalus striatus*) powder and toxicity study in zebrafish (*Danio rerio*) model. *Pharmaceutical Sciences*, 25(2), 145-153.
- Tungadi, R., & Wicita, P. (2020). Formulation, optimization, and characterization of snakehead fish (*Ophiocephalus striatus*) powder nanoemulgel. *Brazilian Journal of Pharmaceutical Sciences*, 56, 1-8.
- Ulandari, A., Kurniawan, D., & Putri, A. S. (2011). Potential of snakehead fish protein in preventing kwashiorkor in toddlers in Jambi Province. Faculty of Medicine, Jambi University. Retrieved from <https://adoc.pub/potensi-protein-ikan-gabus-dalam-mencegah-kwashiorkor-pada-b.html>
- Vemula, V. R., Lagishetty, V., & Lingala, S. (2010). Solubility enhancement techniques. *International Journal of Pharmaceutical Science Review and Research*, 5(1), 41-51.
- Winterhalter, M., & Lasic, D. D. (2013). Liposome stability and formation: experimental parameters and theories on the size distribution. *Chemistry and Physics of Lipids*, 64, 35-37.
- Zhang, L., Zhang, F., Fan, Z., Liu, B., & Meng, X. (2019). DHA and EPA nanoemulsion prepared by the low-energy emulsification method: process factors influencing droplet size physicochemical stability. *Food Research International*, 121(7), 359-366.
- Zhu, W., Guo, C., Yu, A., Gao, Y., Cao, F., & Zhai, G. (2009). Microemulsion-based Hydrogel Formulation of penciclovir for topical delivery. *International Journal of Pharmaceutics*, 378(1-2), 152-158.