

IOP Conference Series

Earth and Environmental Science

1st Workshop on Methodology for Agriculture
and Forestry (METRO AGRIFOR)

275

VOLUME 275 – 2019

3–5 October 2019
Alessano, Italy

EDITOR:
Enrico Pimentel Fernandes

The open access journal for conference proceedings
<https://openaccess.onlinelibrary.wiley.com/jps>



ⓘ ×

CUHK

Research Postgraduate Studies

PhD Programmes with Annual Studentship of US\$27,730 at CUHK!

gs.cuhk.edu.hk


OPEN

IOP Conference Series: Earth and Environmental Science

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United Kingdom  Universities and research institutions in United Kingdom	Earth and Planetary Sciences Earth and Planetary Sciences (miscellaneous) Environmental Science Environmental Science (miscellaneous)	IOP Publishing Ltd.	26
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Conferences and Proceedings	17551307 , 17551315	2010-2020	Homepage How to publish in this journal ees@ioppublishing.org
SCOPE			

The open access IOP Conference Series: Earth and Environmental Science (EES) provides a fast, versatile and cost-effective proceedings publication service.

 Join the conversation about this journal



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/widget.php?...>
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new free tool**.

Get it





Source details

IOP Conference Series: Earth and Environmental Science

CiteScore 2020

0.5

[i](#)

Scopus coverage years: from 2010 to Present

ISSN: 1755-1307 E-ISSN: 1755-1315

Subject area: [Environmental Science: General Environmental Science](#)

SJR 2020

0.179

[i](#)[Earth and Planetary Sciences: General Earth and Planetary Sciences](#)[Physics and Astronomy: General Physics and Astronomy](#)

Source type: Conference Proceeding

SNIP 2020

0.436

[i](#)[View all documents >](#)[Set document alert](#) [Save to source list](#) [Source Homepage](#)[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology x

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore [2020](#) ▼

0.5 = $\frac{25,463 \text{ Citations 2017 - 2020}}{49,883 \text{ Documents 2017 - 2020}}$

Calculated on 05 May, 2021

CiteScoreTracker 2021 [i](#)

0.6 = $\frac{39,730 \text{ Citations to date}}{69,454 \text{ Documents to date}}$

Last updated on 04 December, 2021 • Updated monthly

CiteScore rank 2020 [i](#)

Category	Rank	Percentile
----------	------	------------


Environmental Science

General Environmental Science	#183/220	17th
-------------------------------	----------	------

Earth and Planetary Sciences

General Earth and Planetary Sciences	#157/186	15th
--------------------------------------	----------	------

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site <](#)



The open access *IOP Conference Series: Earth and Environmental Science (EES)* provides a fast, versatile and cost-effective proceedings publication service.

Latest published conferences

Vol 948



Go

Conference archive

2021



Go

View forthcoming volumes accepted for publication.

If you would like more information regarding *IOP Conference Series: Earth and Environmental Science* please visit conferenceseries.iop.org, and if you are interested in publishing a proceedings with IOP Conference Series please visit our page for conference organizers.

Conference organizers can use our online form and we will get in touch with a quote and further details.

Most read

Most cited

Latest articles

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



JOURNAL HISTORY

2008-present IOP Conference Series: Earth and Environmental Science

doi:10.1088/issn.1755-1315

Online ISSN: 1755-1315

Print ISSN: 1755-1307



PAPER • OPEN ACCESS

Preface

To cite this article: 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **589** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Tidal inundation \("Rob"\) investigation using time series of high resolution satellite image data and from institu measurements along northern coast of Java \(Pantura\)](#)
Heri Andreas, Usriyah, Hasanuddin Zainal Abidin et al.
- [Majalaya Flood Early Warning System: A Community Based Approach](#)
I Dewa Gede A Junnaedhi, Edi Riawan, Rusmawan Suwarman et al.
- [Hydrothermal alteration and timing of gold mineralisation in the Rumbia Complex, Southeast Arm of Sulawesi, Indonesia](#)
Musri Mawaleda, Emmy Suparka, Chalid Idham Abdullah et al.

Preface

3rd International Conference of the Transdisciplinary Research on Environmental Problems in Southeast Asia (TREPSEA 2018)

1. About TREPSEA 2018

The International conference of the Transdisciplinary Research on Environmental Problems in Southeast Asia (TREPSEA) aims to conduct integrative research of interactions between natural environment and human-social systems in Southeast Asia to solve the environmental problems in Southeast Asia. Its scope thus includes topics of geoscience, environmental science, engineering, medicine, economy, culture, education, and administration.



Transdisciplinary Research (TDR) is defined as research efforts conducted by investigators from different disciplines and non-academic participants working jointly to create new conceptual, theoretical, methodological, and translational innovations. Related stakeholders include sponsoring intuitions, governments, development organizations, business and industries, civil society (inhabitant, NGO's etc.), and the media.

Our vision of TREPSEA conference is to become a unique and leading international conference on transdisciplinary research studies in varieties of environmental problems of Asia's countries especially ASEAN countries. And our Mission is to share, discuss and exchange knowledge, experience and outcome of transdisciplinary research on environmental problems with various stakeholders.

The 3rd international conference of the Transdisciplinary Research on Environmental Problems in Southeast Asia (TREPSEA 2018) was held on August 11 – 12, 2018 at State University of Gorontalo's Hotel TC Damhil UNG in Gorontalo city, Sulawesi, Indonesia.

TREPSEA 2018 conference featured oral and poster presentations and workshop, and participants presented, shared and discussed their experience on the following topics:

4 Main Topics of TREPSEA 2018

<p>1. Disaster Mitigation:</p> <ul style="list-style-type: none"> ○ Volcanic Eruption ○ Flood ○ Earthquake ○ Tsunami ○ Landslide ○ Groundwater 	<p>2. Sustainable Development and Environmental Preservation:</p> <ul style="list-style-type: none"> ○ Heavy Metal Problem ○ Conversion of Waste to Energy ○ Renewable Energy
<p>3. Measure and Improvement to Urban Environmental Problems:</p> <ul style="list-style-type: none"> ○ Urban Management and Community Development ○ Urban Transportation Planning ○ Traffic Control and Surveillance System ○ Garbage Problem ○ Waste Water Problem 	<p>4. Food and Human Security:</p> <ul style="list-style-type: none"> ○ Food ○ Security problems

2. TREPSEA 2018 General Committee

Prof. Masayuki Sakakibara, Ph. D.

Chairperson

*Research Institute for Humanity and Nature (RIHN) &
Ehime University*

Prof. Dr. Fenty Usman Puluhulawa, M.Hum,
Vice-Chairperson
State University of Gorontalo

General Committee Members

Prof. Dr. Dwia Aries Tina Pulubuhu, M.A.
Hasanuddin University

Prof. Dr. Emmy Suparka
Bandung Institute of Technology

Prof. Dr. Syamsu Qamar Badu, M.Pd.
Rector of State University of Gorontalo

Dr. Arif Satria, SP, M.Si
Bogor Agricultural University

Prof. Dr. H. Gufran Darma Dirawan, M. EMD.,
State University of Makassar

Scientific Committee Members

Ir. Benjamin Sapiie, *ITB*
Prof. Budu, *UNHAS*
Dr. Hayati Sari Hasibuan, *UI*
Dr. Heriansyah Putra, *IPB*
Prof. Hiroki Kasamatsu, *EU*
Dr. Eng. Imam Achmad Sadisun, *ITB*
Prof. Katsuya Tanaka, *RCSE, SU*
Prof. Mai Trong Nhuan, *VNUH*
Dr. Lukman AR Laliyo, *UNG*
Dr. rer. nat. Mohamad Jahja, *UNG*
Prof. Dr. Muh. Nadjib Bustan (MD), *UNM*
Dr. Zaenal Abidin, *IPB*

Local Committee Members

Head: Dr. Sc.Yayu Indriati Arifin, *UNG*
Secretary: Deasy Natalia Botutihe, M.Si, *UNG*
Lanto Ningrayati Amali, Ph.D, *UNG*
Novriyanto Napu, Ph.D, *UNG*
Dr. Lilan Dama, *UNG*
Dr. Sunarty Eraku, *UNG*
Raflin Hinelo, M.Si, *UNG*
Intan Novianti Manyoe, MT, *UNG*

Abbreviation

EU: Ehime University
IPB: Bogor Agricultural University
ITB: Bandung Institute of Technology
RCSE, SU: Research Centre for Sustainability and Environment, Shiga University
UI: University of Indonesia
UNG: State University of Gorontalo
UNHAS: Hasanuddin University
UNM: State University of Makassar
VNUH: Vietnam National University, Hanoi

3. Special Guest

- Prof. Dr. Syamsu Qamar Badu, M.Pd.
Rector of State University of Gorontalo (UNG)

4. Keynote and Invited Speakers

- Prof. Dr. Ir. Nelson Pomalingo, M.Pd
The Regent of Gorontalo Regency
- H. Hamim Pou, S.Kom, MM
The Regent of Bone Bolango Regency
- Prof. Dr. Ir. Mahludin H. Baruwadi, M.P
Vice Rector for Academic Affairs
State University of Gorontalo (UNG)
- Ms. Kana Furusawa
Vice Secretary General,
The Japanese Geoparks Network

5. Organizers

- Research Institute for Humanity and Nature (RIHN), Japan
- State University of Gorontalo (UNG), Indonesia
- Ehime University (EU), Japan
- Bandung Institute of Technology (ITB), Indonesia
- Muhammadiyah University of Gorontalo (UMGo), Indonesia

6. Manuscripts for TREPSEA 2018

Although more than 120 abstracts were received and presented in TREPSEA 2018 conference, 52 manuscripts were accepted for reviewing processes. Only 37 manuscripts were finally accepted for the publication process.

7. TREPSEA 2018 Photos



TREPSEA2018's Ice Breaking Dinner



TREPSEA2018's Ice Breaking Dinner



TREPSEA2018's Event Photos



TREPSEA2018's Event Photos



TREPSEA2018's Event Photos



TREPSEA2018's Event Photos



TREPSEA2018's Event Photos



TREPSEA2018's Event Photos

PAPER • OPEN ACCESS

Peer review declaration

To cite this article: 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **589** 011002

You may also like

- [Peer review declaration](#)
- [Peer review declaration](#)
- [Peer review declaration](#)

View the [article online](#) for updates and enhancements.

Peer review declaration

All papers published in this volume of IOP Conference Series: Earth and Environmental Science have been peer reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:** Single-blind / Double-blind / Triple-blind / Open / Other (please describe)

Single-blind

- **Conference submission management system:**

TREPSEA 2018 Conference submission had been managed under the supervision of the Committee of TREPSEA 2018. The Committee of TREPSEA 2018 accepted potential abstracts as well as full papers related to its topics to be presented at the Conference. In order to ensure for publication of the manuscripts, TREPSEA 2018 Committee managed the numerous processes including scope checking, looking for relevant SME (Subject Matter Experts such as University Professors, Scientists, and Researchers) and then requested them as reviewers/referees to review the submitted manuscripts under single-blinded peer-reviewing method. Finally, the manuscripts considered by the recommendation of peer reviewers and editorial board were accepted for publication process. The Committee of TREPSEA 2018 declares that the submitted manuscripts have not been published before nor submitted to another journal for the consideration of publication.

- **Number of submissions received:**

Although more than 120 abstracts were received and presented in TREPSEA 2018 conference, 52 manuscripts were accepted for reviewing processes. Only 37 manuscripts were finally accepted for the publication process.

- **Number of submissions sent for review:**

52 manuscripts for reviewing process.

- **Number of submissions accepted:**

37 manuscripts were finally accepted for the publication process.

- **Acceptance Rate (Number of Submissions Accepted / Number of Submissions Received X 100):**



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

71%

- **Average number of reviews per paper:**

Each manuscript was reviewed by 2 - 3 reviewers, and underwent the additional revising processes approximately for 1 - 3 times based on the feedback of reviewers.

- **Total number of reviewers involved:**

23 reviewers

- **Any additional info on review process:**

NA

- **Contact person for queries:**

Professor Masayuki Sakakibara
Chairperson, TREPSEA 2018
Current Vice – Chairperson, TREPSEA 2021

Project Leader SRIREP Project / Mercury Project
Research Institute for Humanity and Nature (RIHN) &
Ehime University
Email: sakaki@chikyu.ac.jp / sakakibara.masayuki.mb@ehime-u.ac.jp

Myo Han Htun
Research Associate
SRIREP Project / Mercury Project
Research Institute for Humanity and Nature (RIHN)
Email: myo@chikyu.ac.jp / myohanhtun@gmail.com

Please call us during office hours (08:30 – 17:00) except weekend and public holidays
075-707-2357 (Japanese Speaking) 075-707-2344 (English Speaking)

Table of contents

Volume 589

2020

◀ Previous issue Next issue ▶

The 3rd International conference of the Transdisciplinary Research on Environmental Problems in Southeast Asia 11-12 August 2018, Negeri Gorontalo, Indonesia

Accepted papers received: 29 September 2020

Published online: 18 November 2020

Open all abstracts

Preface

OPEN ACCESS	011001	
Preface		
+ Open abstract	View article	 PDF

OPEN ACCESS	011002	
Peer review declaration		
+ Open abstract	View article	 PDF

01 - Disaster Mitigation

OPEN ACCESS	012001	
Study on Influence of Joint Orientation on Rock Engineering Properties for Mining and Infrastructure Design		
Purwanto, Djamaluddin, Sabrianto Aswad, Dedi Eka Wahyuwibowo, Nirmana Fiqra Qaidahiyani and Hideaki Yasuhara		
+ Open abstract	View article	 PDF

OPEN ACCESS	012002	
Geology and 2D modelling of magnetic data to evaluate surface and subsurface setting in Bongongoayu geothermal area, Gorontalo		
I N Manyoe, D A Suriamihardja, U R Irfan, S S Eraku, S S S Napu and D D Tolodo		
+ Open abstract	View article	 PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, [see our Privacy and Cookies policy.](#)

012003 

Sediment material potential of Bone River as a prevention of silting and flood of Bone River estuary, Gorontalo

S S S Napu, T H Salama, I N Manyoe, F C A Usman, I Samir, A W W Badaru and K Sugianti

[+ Open abstract](#)

[View article](#)

[PDF](#)

012004

Flood Points Distribution Mapping for the Determination of Evacuation Routes and Disaster Posts in Gorontalo City

Indrawan and Daud Yusuf

[+ Open abstract](#)

[View article](#)

[PDF](#)

012005

Characteristic of the Mount Colo Volcano, Una-Una Island, Central Sulawesi Province: Tectonic Evolution and Disaster Mitigation

Purnama Sendjaja, Emmy Suparka, Chalid I. Abdullah and IGB Eddy Sucipta

[+ Open abstract](#)

[View article](#)

[PDF](#)

012006

Geochemical Exploration of Geothermal Manifestations: Study Case of Bajawa Volcanic Complex, Flores Island, East Nusa Tenggara Province

H H Herzegovina, M Abdurrachman, M Hafidh and M Divi

[+ Open abstract](#)

[View article](#)

[PDF](#)

012007

Facies Study of Lake Deposits Formation (Qpl) To Determine Deposition Environment of Ancient Limboto Lake: a Preliminary Result

A K M A Amin, M Sakakibara, Y I Arifin and N Akase

[+ Open abstract](#)

[View article](#)

[PDF](#)

012008

Geophysical survey of landslide movement and mechanism in Gorontalo Outer Ring Road, Gorontalo

F C A Usman, I N Manyoe, R F Duwingik and D N P Kasim

[+ Open abstract](#)

[View article](#)

[PDF](#)

012009

Landslide disaster and its prevention works in Shikoku region of Japan

Kazunari Kimura, Tomohiro Nagata and Shuya Kan

[+ Open abstract](#)

[View article](#)

[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).

012009

Volcanostratigraphy of Krakatoa Islands, South Lampung District, Lampung Province

Taufik Ismail, Mirzam Abdurrachman, Yan Rizal and Kumala Hardjawidjaksana

[+ Open abstract](#)

[View article](#)

[PDF](#)

012011

The Influence of Rock Resistance for Geomorphology of Krakatau Islands, South Lampung District, Lampung Province

Taufik Ismail, Mirzam Abdurrachman, Yan Rizal, Kumala Hardjawidjaksana and Rahmawati Arbie

[+ Open abstract](#)

[View article](#)

[PDF](#)

02 - Sustainable Development and Environmental Preservation

OPEN ACCESS

012012

Enhanced removal of heavy metals using sorbent and *Phragmites australis*

Nguyen Quoc Bien and Nguyen Thi Hoang Ha

[+ Open abstract](#)

[View article](#)

[PDF](#)

012013

Mercury exposure from small scale gold mining activities and neurological symptoms on inhabitants and miners: a case study in Bolaang Mongondow, North Sulawesi Province, Indonesia

Yayu Indriati Arifin, Masayuki Sakakibara, Koichiro Sera, Puluhulawa Fenty Usman and Fitryane Lihawa

[+ Open abstract](#)

[View article](#)

[PDF](#)

012014

Distribution of heavy metals in sediment along the Southern coast of Vietnam

Nguyen P C Tu, Nguyen N Ha, Nguyen N Tri and Nguyen V Dong

[+ Open abstract](#)

[View article](#)

[PDF](#)

012015

Socioeconomic and Mercury Exposure to The Goldsmiths in Manggal Subdistrict of Urban Artisanal Gold Mining (UAGM) Area in Makassar, South Sulawesi, Indonesia

Hasriwiani Habo Abbas, Masayuki Sakakibara, Koichiro Sera, Lukmanul Hakim Arma and Mansur Sididi

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012016

Regional Geochemical Map of West Java, Indonesia: Evaluation for Environmental and Mineral Resources

Purnama Sendjaja and Ari Kusniadi

[+ Open abstract](#)

[View article](#)

[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012017

Liquefaction Resistance of Sand Improved with Enzymatically Induced Calcite Precipitation based on Laboratory Investigation

Minson Simatupang

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012018

Development of Bioadsorbent Chitosan from Shrimp Shell Waste to Mercury Absorption Efficiency

Astin Lukum, Yoseph Paramata, Deasy N Botutihe, Jefrin Akume, Kostiawan Sukamto and Arfiani Rizki Paramata

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012019

Temperature dependence for purification of leachate containing heavy metals by phytoremediation using the artificial channel

K Okazaki, T Kurahashi, S Yamazaki and M Sakakibara

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012020

Sociodemographic Attributes and Dependency on Artisanal and Small-scale Gold Mining: the Case of Rural Gorontalo, Indonesia

Satoru Komatsu, Katsuya Tanaka, Masayuki Sakakibara, Yayu Indriati Arifin, Sri Manovita Pateda and Intan Noviantari Manyoe

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012021

Chemical content analysis of coral limestone as prospecting of extractive development in Gorontalo City

W Annisa, I N Manyoe, A K Mubarak, S S S Napu, I G S Pratama and S Fatimah

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012022

Environmental Risk and Health Hazardous Substances in Artisanal Small-Scaled Gold Mining in Sekotong, West Nusa Tenggara, Indonesia

S Novianti, M Abdurrachman, D Claudia and NI Basuki

 [Open abstract](#)

 [View article](#)

 [PDF](#)

OPEN ACCESS

012023

Geodiversity of the Togean Islands National Park, Central Sulawesi Province for

Geopark Assessment By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).

Purnama Sendjaja, Emmy Suparka and M U Botjing



[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012024

The nature of ultramafic rocks from Sulawesi, Indonesia and their suitability for CO₂ sequestration

Sufriadin, S Widodo, M Thamrin, A Maulana, A Ito and T Otake

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012025

"Gunung Api Purba Nglangeran" welcomes UNESCO Global Geopark Reassessment in 2019

Nabila Novasari Soviana, Budi Brahmantyo, Mirzam Abdurrachman and Firman Sauqi Nur Sabila

[+ Open abstract](#)

[View article](#)

[PDF](#)

03 - Measure and Improvement to Urban Environmental Problem

OPEN ACCESS

012026

Eco-Biger (Ecopreneur Biogester) is Right Digester Innovation for Household to Increase People's Welfare in Indonesia

A K Syahroni, W T Roby and T Rahmanda

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012027

Customary Practices of Musyawarah Mufakat: An Indonesian Style of Consensus Building

Citra Anggita and Tsuyoshi Hatori

[+ Open abstract](#)

[View article](#)

[PDF](#)

04 - Food and Human Security

OPEN ACCESS

012028

The Development of Earthquake Risk Reduction Curriculum for 4-6 Years Old Children in Gorontalo

P P Ardni, I N Arifin, H Syahputra and B Pupala

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012029

Measuring the Relationship among Personal Network with Japanese Language Skill and Fit Well with Life in Japan

Vita Fajriani Ridwan and Nobuhiko Matsumura

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



OPEN ACCESS

012030

Monitoring and Evaluation of Fish Quality Standard Compliance

Faiza A Dali and Rahim Husain

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012031

Analysis on quality improvement of used cooking oil with young pineapple (*Ananas comosus*) using Taguchi Method

Muhammad Arfan, Wahyu Supartono and Henry Yuliando

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012032

Transformation of Fallow Yard Become Potential for Agribusiness of Dragon Fruit

Dewa Oka Suparwata and Moh. Muchlis Djibrin

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012033

Concentration of Pb, Sn and Fe Metals on Milk Products and Canned Fish in Gorontalo City

Wiwin Kunusa Rewini, Nurain Thomas, Doly Prima Silaban, Hendrik Iyabu and Muhammad Taupik

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012034

Preliminary Study of Clinical Antidiabetic Activity of Salam Leaves (*Eugenia Polyantha*) and Sambiloto Leaves (*Andrographis Paniculata*) In Type 2 Diabetic Patients

N I Ischak and D N Botutihe

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012035

Utilization of Corncob Cellulose Isolate (*Zea mays*) as Adsorbent of Heavy Metal Copper and Cadmium

Ishak Isa, Eka Setiawati, Erni Mohammad and Wiwin Kunusa

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012036

The Effect of Chitosan of Ark Clam Shells to Reduce Pb and Hg Level and Amount of Bacteria in the Blood Cockles Meatball

Margaretha Solang, Djuna Lamondo, Syam S Kumaji and Novri Y Kandowangko

[+ Open abstract](#)[View article](#)[PDF](#)This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).

012037

The potential of isolates endophytic fungi from the leaves of henna plant (*Lawsonia inermis*) toward the growth of *Candida albicans* and *Staphylococcus aureus*

A M Hasan, W D Uno and S Baruwadi

 Open abstract

 View article

 PDF

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)



This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



PAPER • OPEN ACCESS

Geology and 2D modelling of magnetic data to evaluate surface and subsurface setting in Bongongoayu geothermal area, Gorontalo

I N Manyoe¹, D A Suriamihardja², U R Irfan³, S S Eraku⁴, S S S Napu¹ and D D Tolodo¹

Published under licence by IOP Publishing Ltd

IOP Conference Series: Earth and Environmental Science, Volume 589, The 3rd International conference of the Transdisciplinary Research on Environmental Problems in Southeast Asia 11-12 August 2018, Negeri Gorontalo, Indonesia

Citation I N Manyoe *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **589** 012002

intan.manyoe@ung.ac.id

¹ Geological Engineering, Universitas Negeri Gorontalo, B.J. Habibie Street, Bone Bolango, Gorontalo, 96119 Indonesia

² Geophysics, Universitas Hasanuddin, Perintis Kemerdekaan Street, Makassar, South Sulawesi, 90245 Indonesia

³ Geological Engineering, Universitas Hasanuddin, Malino Street, Gowa, South Sulawesi, 92119 Indonesia

⁴ Geography Education, Universitas Negeri Gorontalo, B.J. Habibie Street, Bone Bolango, Gorontalo, 96119 Indonesia

<https://doi.org/10.1088/1755-1315/589/1/012002>

Buy this article in print

 Journal RSS

Sign up for new issue notifications

Create citation alert

PDF

Help

Abstract

Bongongoayu is one of the regions in Indonesia that has geothermal potential. Bongongoayu requires surface and subsurface data to support the preliminary data. This research aims to determine surface and subsurface data conducted by geology and magnetic method. The surface

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy 

data, including geomorphology, lithology, hydrology and manifestation. The subsurface data have taken by the magnetic method. The result showed that the geothermal manifestation of Bongongoayu is a hot pool. The surface temperature is 43 to 59 °C. The geomorphology units is composed of volcanic hills unit and lake plains unit. The lithology of the research area is composed of granite and alluvial deposits. Based on petrographic analysis, the level of alteration in granite rocks is 65% and is classified as moderate alteration. The recharge area is in the north and southwest of the research area. Discharge area is in the central area. Based on 2D magnetic modeling, there are two subsurface layers. The first layer is alluvial and the second layer is granite. Rocks that are under the alluvial layer and have been altered are interpreted as a cap rock. 2D magnetic models show normal faults in the research area as a controlling factor for geothermal fluid.

Export citation and abstract

BibTeX

RIS

◀ Previous article in issue

Next article in issue ▶



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

You may also like

JOURNAL ARTICLES

PDF

Controlled vapor growth of 2D magnetic Cr₂Se₃ and its magnetic proximity effect in heterostructures Help

Viscous effects on plasmoid formation from nonlinear resistive tearing growth in a Harris sheet

Subsurface structure identification uses derivative analyses of the magnetic data in Candi Umbul-

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

see our [Privacy and Cookies policy](#).



Rational Design of Two-Dimensional Magnetic Chromium Borides Based on First-Principles Calculation

PDF

Help



PAPER • OPEN ACCESS

Geology and 2D modelling of magnetic data to evaluate surface and subsurface setting in Bongongoayu geothermal area, Gorontalo

To cite this article: I N Manyoe *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **589** 012002

View the [article online](#) for updates and enhancements.

You may also like

- [Controlled vapor growth of 2D magnetic Cr₂Se₃ and its magnetic proximity effect in heterostructures](#)

Danliang Zhang, , Chen Yi et al.

- [Viscous effects on plasmoid formation from nonlinear resistive tearing growth in a Harris sheet](#)

Nisar AHMAD, Ping ZHU, et al.

- [Subsurface structure identification uses derivative analyses of the magnetic data in Candi Umbul-Telomoyo geothermal prospect area](#)

U Septyasari, S W Niasari and P D Maghfira

Geology and 2D modelling of magnetic data to evaluate surface and subsurface setting in Bongongoayu geothermal area, Gorontalo

I N Manyoe¹, D A Suriamihardja² U R Irfan³ S S Eraku⁴ S S S Napu¹ and D D Tolodo¹

¹Geological Engineering, Universitas Negeri Gorontalo, B.J. Habibie Street, Bone Bolango, Gorontalo, 96119 Indonesia

²Geophysics, Universitas Hasanuddin, Perintis Kemerdekaan Street, Makassar, South Sulawesi, 90245 Indonesia

³Geological Engineering, Universitas Hasanuddin, Malino Street, Gowa, South Sulawesi, 92119 Indonesia

⁴Geography Education, Universitas Negeri Gorontalo, B.J. Habibie Street, Bone Bolango, Gorontalo, 96119 Indonesia

Corresponding Author: intan.manyoe@ung.ac.id

Abstract. Bongongoayu is one of the regions in Indonesia that has geothermal potential. Bongongoayu requires surface and subsurface data to support the preliminary data. This research aims to determine surface and subsurface data conducted by geology and magnetic method. The surface data, including geomorphology, lithology, hydrology and manifestation. The subsurface data have taken by the magnetic method. The result showed that the geothermal manifestation of Bongongoayu is a hot pool. The surface temperature is 43 to 59 °C. The geomorphology units is composed of volcanic hills unit and lake plains unit. The lithology of the research area is composed of granite and alluvial deposits. Based on petrographic analysis, the level of alteration in granite rocks is 65% and is classified as moderate alteration. The recharge area is in the north and southwest of the research area. Discharge area is in the central area. Based on 2D magnetic modeling, there are two subsurface layers. The first layer is alluvial and the second layer is granite. Rocks that are under the alluvial layer and have been altered are interpreted as a cap rock. 2D magnetic models show normal faults in the research area as a controlling factor for geothermal fluid.

1. Introduction

Indonesia affected by three major plates, i.e. SE-Asian Plate, the Indo-Australian Plate, and the Pacific Plate [1,2]. Sulawesi is formed in the convergence zone of the Eurasian, Pacific, Australian plate complexes [3] and affected by Philippines plate [4]. Northern part of the North Arm Sulawesi is the Sulawesi Sea Subduction. Sulawesi Sea Subduction was thought to have been active since the Early Tertiary and produced Tertiary volcanic arcs that stretch from around Toli-Toli to Manado [5].

The reconstruction in Sulawesi Sea shows that the volcanic arc system surrounding the Sulawesi Sea has been established since the Late Miocene [6]. Volcanism on North Arm Sulawesi is related to

Sulawesi Sea Subduction [3]. Geothermal associated with an area of active volcanic [7]. Besides being associated with active volcanoes, geothermal is also associated with faults [8,9] and intrusions [10].

The volcano-tectonic conditions of Indonesia cause Indonesia, especially the North Arm of Sulawesi, to have big geothermal potential. This big geothermal potential is being developed by the government to support national energy. Based on a recent survey from the Geological Agency, the Ministry of Energy and Mineral Resources [10], there are 331 potential areas that have been identified consisting of 11,073 MW of resources and 17,506 MW of reserves spread across 30 provinces.

Areas identified as having geothermal potential will be explored through scientific survey methods. The scientific survey method can be in the form of geological and geochemical studies. In addition to more detailed geological and geochemical studies, various geophysical techniques can be used, including geomagnetic methods [11].

Geological surveys are preliminary surveys carried out in geothermal exploration by analyzing rocks, determining morphological characteristics, hydrological conditions, and geothermal systems. Geophysical survey in the form of magnetic, electrical and other subsurface surveys.

The geomagnetic method utilizes the contrast magnetization of subsurface rocks [12]. In geothermal areas, geothermal fluids can cause massive changes in the chemical and physical properties of subsurface rocks. Another change is the magnetic properties of rocks will be down or lost due to heat generated [13].

Geological surveys have been conducted in several geothermal areas in Gorontalo, one of which is in the Lombongo geothermal area [14]. Surveys that have been conducted in Bongongoayu are electrical [15] and magnetic [16] surveys, except that they have not provided surface geology and subsurface magnetic models. Based on the explanation above, it is very interesting to study the Bongongoayu geothermal area. This study aims to determine the characteristics of surface and subsurface in the Bongongoayu geothermal area based on geological and magnetic data.

2. Data and Method

The study area is located at coordinates 0.6230 – 0.7890 N and 122.6940 – 122.4510 E, administratively included in the Bongongoayu area (also known as Diloniyohu area), Boliyohuto District, Gorontalo Regency, Indonesia (Figure 1). The research area is located in a depressed area that extends in the central part of Gorontalo.




Figure 1. Geologic map of research area [17].

The instruments used in the study are GPS receivers, Proton Precession Magnetometer GSM 19-T v7.0, Nikon ci-POL Microscope, infrared thermometer, geological hammer and geological compass. The GPS receiver is to determine the location of base station, mobile stations and observation stations. One set of Proton Precession Magnetometer GSM 19-T v7.0 is to measure and determine the distribution of magnetic anomalies. Nikon ci-POL Microscope is to determine mineral composition of rocks. Infrared thermometer to measure the temperature of geothermal manifestation. Geological hammer is to take rock samples and a geological compass is to determine direction.

2.1 Data collection and acquisition

Based on the regional geological map, rocks formation in the research area are composed of Bone Diorite (Tmb), Boliyohuto Diorite (Tmbo), Dolokapa Formation (Tmd), Pinogu Volcanics (Tqpv), and the Lake Deposit (Qpl). The hot pool is located in the Lake Deposit (Figure 1). The geological data collection is carried out to determine rock units in the research area. Rock samples were taken in the research area with the aim to determine the composition of rock minerals. the geological data collection was also carried out to determine the geomorphological and hydrological data of the research area. Determination of geomorphological units by conducting field surveys of landforms.

The acquisition data of magnetic is carried out to obtain primary data. A set of Proton Precession Magnetometer GSM 19-T v7.0 is used for measurements the magnetic observation data of base station and mobile stations. Acquisition data of magnetic is carried out with a closed loop system (A, B, A), i.e. measurements always start and end at the same point.

2.2 Processing, analysis and interpretation data

Thin section of rock samples was analyzed in a laboratory using a Nikon Ci-POL Microscope. Analysis petrography using IUGS (1979) classification. Geomorphology data analysis using Van Zuidam classification (1985). Hydrological analysis of the research area was conducted to determine the recharge area and discharge area. This analysis, using field data, topographic data, and lithology.


Magnetic data processing begins by correcting the measurement data in base station and mobile stations with diurnal variation and International Geomagnetic Reference Field (IGRF) correction. Magnetic modeling of subsurface data is carried out after data correction. Modeling of subsurface magnetic data is a continuation of magnetic studies that were not carried out in previous studies [16]. Previous studies have only mapped total magnetic field anomalies without subsurface modeling. Interpretation of magnetic data uses rock susceptibility values as provided by Telford (1990).

3. Result and Discussion

3.1. Geomorphology

Geomorphological studies in a geothermal field are intended to systematically organize landforms. Based on Van Zuidam classification [18], the Bongongoayu geothermal area is separated into two geomorphological units. Two geomorphological units in the Bongongoayu geothermal area are volcanic hills (Figure 2) and lake plain (Figure 3).

Volcanic hills units occupy the northern and southwestern part of the research area. The elevation of the volcanic hills unit is 28 to 687.5 meters above sea level. The hills which are located in the north are part of the north mountain which consists of Mount Tamboo, Mount Dolokapa and Mount Boliyohuto. The hills in the southwest are part of the Loba Mountains. This geomorphological unit consists of granite, diorite and soil. On the lower slopes, local people use it as an agricultural area.

**Figure 2.** Volcanic hills unit.**Figure 3.** Lake plain unit.

The geomorphological unit of the lake plain is composed of alluvial. This unit occupies the central to southern part of the research area. The elevation of the lake plain unit is about 25 meters above sea level. Lake plain is part of the depression that stretches from east to west Gorontalo. The unit of landform of the lake deposit is located in the western part of the depression. Land use in this unit is for settlement, agriculture/plantation and fisheries.

3.2. Lithology

Bongongoayu geothermal manifestations are hot pool, found in alluvial units. Alluvial units are located in the southern part of the research area. Alluvial is concentrated around geothermal manifestations extending to the south, east and west of geothermal manifestations. The color of alluvial deposits in the research area is blackish brown. The northern part and southwestern part of the research area is composed of granite units. Granite units are composed of granite and diorite.

Granite is located in the northern part and southwestern part of the research area (Figure 4). Rock texture is holocrystalline and composed of orthoclase, quartz and plagioclase. Secondary minerals found in granite rocks are quartz, clay minerals and opaque minerals. The mineral form is anhedral to subhedral.

**Figure 4.** Granite (photo direction N 70° E).**Figure 5.** Carbonate vein in diorite (photo direction N 40 °E).

Diorite is located in the northern part of the research area on volcanic hills units. Rock texture is holocrystalline and composed of orthoclase and quartz. Secondary minerals found in diorite are clay minerals and opaque minerals. The form of minerals is anhedral and subhedral. Vein Carbonate is found in diorite (Figure 5).

Microscopic analysis of rock samples in the study area shows that rock samples are alteration igneous rocks. Rock structure type is massive. Rocks in the study area are granite based on petrographic analysis of rock samples in the research area.

The mineral composition of rocks in the research area is shown in Table 1. Microscopic analysis of rock samples in the study area showed that the secondary mineral composition was at least 45% and the

highest was 65%. Primary mineral forming rocks in the study area are orthoclase, quartz, and plagioclase. Secondary minerals found in rocks are quartz, clay minerals and opaque minerals.

Table 1. The mineral composition of the rocks in the Bongongoayu geothermal area.

Station	Rocks	Mineral Composition (%)					
		Ort	Qz	Pl	Qz	Cl	Opq
SR.a	Granite	20	10	10	35	15	10
SR.b	Granite	20	10	15	25	15	15
SR.c	Granite	10	20	25	30	10	5
SR.d	Granite	20	15	10	35	15	5
SR.e	Granite	10	25	-	45	15	5
SR.f	Granite	20	10	10	35	15	10

Note: Ort = Orthoclase; Qz = Quartz; Pl = Plagioclase; Cl = Clay Mineral; Opq = Opaque Mineral.

The composition of primary minerals at SR.a is 40%, while the secondary mineral composition is 60%. Primary mineral composition at SR.b is 45%, while secondary mineral composition is 55%. SR.c has 55% of primary minerals while secondary minerals are 45%. SR.d primary mineral composition is 45%, while secondary minerals 55%. SR.e primary mineral composition is 35%, while secondary minerals 65%. SR.f shows the composition of primary minerals is 40%, while secondary minerals is 60%.




Figure 6. Photomicrograph of sample in the study area. Sample SR.a up to SR.f are granite.

Alteration intensity is defined as the level change based on the percentage of secondary minerals. Morrison [19] provides alteration intensity based on the percentage of secondary minerals. Not altered, if no secondary mineral. Weak altered if secondary mineral is 25%. Moderate altered if the secondary mineral percentage is about 25 to 75% and strong altered if secondary mineral is >75%.

If all primary minerals are altered (except quartz, zircon and apatite) but the primary texture is still visible, the intensity of the altered is called intense altered intensity. The total altered intensity is all primary minerals are changed (except quartz, zircon and apatite) and the primary texture is not visible. Microscopic analysis of samples in the research area showed that the secondary mineral composition was at least 45% and the highest was 65% (Figure 6). Based on the classification given by Morrison [19] the intensity of rock altered in the study area is moderate altered.

3.3. Hydrology

Annual rainfall in the research area ranges from 1244 to 2311 mm per year. The highest monthly rainfall in eight years generally occurs in May and the lowest monthly rainfall generally occurs in August. Cumulative rainfall a month with rainfall 90 mm occurred in August and September while the cumulative monthly rainfall >100 mm occurred in October – July [20]. The level of rainfall can affect the height of the water table [21]. Thus, high rainfall in the research area can affect groundwater availability.

The northern part of the Bongongoayu geothermal area is a protected forest area, production forest and Nantu reserved forest [22]. Vegetation in the forest can accelerate the process of rainwater infiltration. Plant roots forest vegetation opens the way for rainwater to be infiltrated. The infiltration process is one of the determinants of water availability.

The next determining factor is the shape of the landscape and permeability. According to Noor [23], water will flow faster on steep slopes than moderate sloping. It means that the rainwater infiltration process takes place in the northern part of the study area. The nature of rock that can carry water is another determining factor for infiltration. The fractures in the rocks in the northern part of the Bongongoayu geothermal area [17] are potential infiltrations. Rainwater infiltrates into the soil through fractures in rocks.

According to Rezky et al. [24] shallow groundwater conditions can be seen from the presence of large and watery rivers throughout the year. The rivers surrounding the Bongongoayu geothermal area are the Bongo River, Diloniyohu River, Buliya River and Paguyaman River. Bongo River, Diloniyohu River and Buliya River are watery rivers throughout the year and head into the Paguyaman River.

Recharge areas in the Bongongoayu geothermal area are in the northern part of the Bongongoayu geothermal area which has moderate slopes and rocks that have fractures. Discharge area is located in the central part, which is located in the lake plain unit. The hydrological conditions in the research area have caused a large amount of surface and groundwater reserves in the Bongongoayu geothermal area.

According to Marshak [21] on a local scale, a recharge area occurred at the top of a hill and discharge area in the valley. This means that on a local scale, the recharge area in the research area is on volcanic hill near from hot pool. Discharge area is in the valley where geothermal manifestations occur.

3.4. Geothermal manifestation

Geothermal manifestation in the Bongongoayu area is hot pool (Figure 7). The hot pool is located at coordinates 0.67785 N and 122.5911 E. The hot pool is located at an altitude of 24 meters above sea level.

The research area consists of granite units and alluvial units. Granite and diorite are included in granite units. Hot springs are located in alluvial units. The northern part of the hot pool is the granite and diorite volcanic hills. The wide area of a hot pool is $\pm 600 \text{ m}^2$.

Measurement of surface temperature in hot pool using an infrared thermometer. The surface temperature of the hot pool ranges from 43 to 59 °C. Physical conditions of hot pool are tasteless and odorless. The appearance of the color of hot water is clear with a neutral pH 6.



Figure 7. Hot pool as a geothermal manifestation in research area.

3.5. Magnetic model

The 2D modelling of magnetic data is shown in Figure 8 and Figure 9. The first model is in the Northeast part of the research area stretching from Northwest-Southeast. The second model is in the Southwestern part of the research area stretching from Northeast-Southwest.




Figure 8. 2D modelling of first anomaly.

There are two layers in the first model consisting of alluvial and granite. The total layer depth in the model is 62.5 m. The first layer has 0.0118 susceptibility value. The first layer is at a depth of 0 to 62.5 m. Based on the value of susceptibility, the first layer is interpreted as alluvial. The second layer has 0.5440 susceptibility value. The second layers are at a depth ranging from 62.5 m. Based on the value of susceptibility, the second layer is interpreted as granite.

In the second model, there are two layers. The first layer has a value of 0.0118 susceptibility value. The first layer is at a depth from 0 to 43.75 m. Based on the value of susceptibility, the first layer is interpreted as alluvial. The second layer has a value of 0.5440 susceptibility value. The depth of the

second layer starts at a depth of 43.75 m. Based on the value of susceptibility, the second layer is interpreted as granite. Lithology and susceptibility value refer to Telford et al. [25].




Figure 9. 2D modeling of second anomaly.

2D magnetic models show that alluvial deposits are surface layers. The layers beneath the alluvial are granite. Some minerals in granite rocks have been altered. The altered part of Granite just below the alluvial layer is interpreted as a cap rock. 2D magnetic models also show the geological structure. Magnetic modeling show that the structure contained in the study area is a normal fault. Geothermal fluids from reservoirs exposed to the surface are controlled by this fault.

4. Conclusion

Geomorphology of the research area is volcanic hills and lake plains. The lithology of the study area is composed of alluvial and granite. The alteration intensity in the research area is included in moderate rock alteration intensity. Recharge area is located in the northern part, while the discharge area is in the valley. The valley is located in the central part of the research area, where hot pool appears. The surface temperature of hot pool ranges from 43 to 59 °C. The physical condition of water in a hot pool is tasteless and odorless. The color appearance of hot pool is clear with neutral pH 6. Based on the 2D model, there are two layers, alluvial with susceptibility value 0.0118 and granite with susceptibility value 0.5440. 2D models show that the normal fault in the research area is a controlling factor for geothermal fluids.

5. References

- [1] Katili J A 1980 *Geotectonics of Indonesia, a modern view* (Jakarta: Directorate General of Mines) pp 271
- [2] Kreemer C, Holt W E, Goes S and Govers R 2000 Active deformation in eastern Indonesia and the Philippines from GPS and seismicity data *J. of Geophys. Res.* **105** 663-680
- [3] Carlile J C, Diggowirogo S and Darius K 1990 Geologic setting, characteristics and regional exploration for gold in the volcanic arcs of North Sulawesi, Indonesia. In: Hedenquist, J.W., White, N.C., Siddeley, G. (Eds.), Epithermal Gold Mineralization of the Circum Pacific: Geology, Geochemistry, Origin and Exploration I. *J. Geochem. Explor.* **35** 105– 140
- [4] Verstappen H T 2010 Indonesian landforms and plate tectonics *Indonesian Journal on Geoscience* **5** 197-207
- [5] Simandjuntak T O 1986 *Sedimentology and tectonics of the collision complex in the East Arm of Sulawesi Indonesia* (United Kingdom: University of London, Royal Holloway and Bedford New College)
- [6] Lee T Y and Lawver L A 1995 Cenozoic plate reconstruction of Southeast Asia *Tectonophysics* **251** 85-138
- [7] Wilson M E and Moss S J 1999 Cenozoic palaeogeographic evolution of Sulawesi and Borneo *Palaeogeography, Palaeoclimatology, Palaeoecology* **145** 303-37
- [8] Sukamto R 1973 *Reconnaissance geologic map of Palu Area, Sulawesi — scale 1:250,000* (Bandung: Geol. Res. and Dev. Cent)

- [9] Boedihardi M, Mulyono A, Ginting A, Mosby M D and Radja V T 1992 Geology, energy potential and development of Indonesia's geothermal prospects. In: Tectonic Framework and Energy Resources of the Western Margin of the Pacific Basin *Bull. Geol. Soc. Malays. Spec. Publ.* **33** 369–385
- [10] Kementerian Energi dan Sumber Daya Mineral 2017 *Potensi panas bumi Indonesia jilid 2* (Jakarta: Kementerian Energi dan Sumber Daya Mineral)
- [11] Árnason K and Gíslason G 2009 Geothermal surface exploration *Makalah disajikan dalam Short Course on Surface Exploration for Geothermal Resources* (El Salvador: United Nations University Geothermal Training Programme and LaGeo) p 17-30
- [12] Gupta H K and Roy S 2006 *Geothermal energy: an alternative resource for the 21st century* (United Kingdom: Elsevier)
- [13] Arsadipura S, Kholid M, Djukardi D 2011 Penyelidikan geofisika terpadu gaya berat, geomagnet dan geolistrik daerah panas bumi Kepahiang, Kecamatan Kaba Wetan, Kabupaten Kepahiang, Provinsi Bengkulu *Proc. Hasil Kegiatan Pusat Sumber Daya Geologi Tahun 2010*
- [14] Manyoe I N 2017 Kajian geologi daerah panas bumi Lombongo Kabupaten Bone Bolango Provinsi Gorontalo *Jurnal Geomine* **5** 10-18
- [15] Manyoe I N 2016 Model inversi data geolistrik untuk penentuan lapisan bawah permukaan daerah panas bumi Bongongoayu, Gorontalo *Jurnal Sainstek* **8** 358-371
- [16] Manyoe I.N, Irvan U R and Suriamiharja D A 2015 Distribusi anomali magnetik daerah panas bumi Bongongoayu, Gorontalo *Proc. Seminar Ilmiah Nasional. Universitas Hasanuddin*
- [17] Bachri S, Sukido and Ratman N 1993 *Peta geologi lembar Tilamuta, Sulawesi* (Bandung: Pusat Penelitian dan Pengembangan Geologi)
- [18] Zuidam R A Van and Zuidam F I van Cancelado 1985 *Aerial photointerpretation in terrain analysis and geomorphologic mapping* (Enschede: ITC)
- [19] Morrison K 1997 *Hydrothermal minerals and their significance* (Auckland: Geothermal and Mineral Service Division of Kingston Morrison Ltd)
- [20] Stasiun Meteorologi Jalaluddin Gorontalo 2014 *Data curah hujan daerah Gorontalo tahun 2005 – 2013* (Gorontalo: Badan Meteorologi, Klimatologi dan Geofisika Stasiun Meteorologi Jalaluddin Gorontalo)
- [21] Marshak S 1955 *Earth portrait of a planet* (New York: W.W Norton and Company)
- [22] Direktorat Bina Rencana Pemanfaatan Hutan Produksi 2010 *Peta sebaran spasial IUPHHK HA/HT/HTR* (Jakarta: Kementerian Kehutanan)
- [23] Noor D 2011 *Geologi untuk perencanaan* (Yogyakarta: Graha Ilmu)
- [24] Rezky A, Hasan A R and Dirasutina S 2005 Penyelidikan geologi daerah panas bumi Suwawa Kabupaten Bone Bolango – Gorontalo *Presented in Kolokium Badan Geologi 2005 Pusat Sumber Daya Geologi Badan Geologi*.
- [25] Telford W M, Geldart L P and Sheriff R E 1990 *Applied geophysics* (Cambridge: Cambridge University Press)

Acknowledgments

We would like to express our deep gratitude to KEMENRISTEKDIKTI for the research fund. We would like to extend our thanks to the staff of the Geological Engineering Laboratory and Geography Laboratory of Universitas Negeri Gorontalo for their help in acquisition data of magnetic.