

Journal of Biological Research

Boletino della Società Italiana di Biologia Sperimentale



www.jbiolres.org

jbr

pagepress

[ABOUT THE JOURNAL](#) [EDITORIAL BOARD](#) [CURRENT ISSUE](#) [ARCHIVES](#) [ANNOUNCEMENTS](#)

Search

[HOME](#) / [Editorial Board](#)

Editorial Board

Editors-in-Chief

Marco Giammanco

Dipartimento di Medicina, Pneumologia, Fisiologia e Nutrizione Umana, Università di Palermo
Palermo, Italy

Gian Luigi Mariottini

Dipartimento di Scienze della Terra, dell'Ambiente e della Vita, Università degli Studi di Genova
Genova, Italy

Associate Editors

Renzo Antolini, Dipartimento di Fisica, Università di Trento, Trento, Italy

Massimo Cocchi, Istituto per la Sintesi Organica e la Fotoreattività, ISOF - CNR, Bologna, Italy

Luigi Pane, Dipartimento di Scienze della Terra, dell'Ambiente e della Vita, Università degli Studi di Genova,
Genova, Italy

Emma Rabino Massa, Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università degli Studi di Torino,
Italy

Editorial Board

James Anthony, Michigan State University, East Lansing, USA
Saeme Asgari, Pasteur Institute, Iran
Han Bao, MSU-DOE Plant Research Laboratory of Michigan State University, USA
Emilia Bellone, University of Genoa, Italy
Maria Grazia Bridelli, University of Parma, Italy
Dario Cantino, University of Turin, Italy
David Caramelli, University of Florence, Italy
Giuseppe Caramia, G. Salesi Hospital, Ancona, Italy
Emilio Carbone, University of Turin, Italy
Brunetto Chiarelli, University of Florence, Italy
Pierluigi Consolo, University of Messina, Italy
Amelia De Lucia, University "Aldo Moro", Bari, Italy
Danila Di Majo, University of Palermo, Italy
Andrea Drusini, University of Padua, Italy
Luciano Fadiga, University of Ferrara, Italy
Vittorio Farina, University of Sassari, Italy
Sara Ferrando, University of Genoa, Italy
William Galanter, University of Illinois, Chicago, USA
Lorenzo Gallus, University of Genoa, Italy
Darren Grice, Institute for Glycomics and School of Medical Science, Griffith University, Nathan, Australia
Stefania Grimaudo, University of Palermo, Italy
Millie Hughes-Fulford, University of San Francisco, USA
Gaetano Leto, University of Palermo, Italy
Gianni Losano, University of Turin, Italy
Mansoor A. Malik, Howard University Hospital, Washington DC, USA
Neville A. Marsh, Queensland University of Technology, Brisbane, Australia
Bruno Masala, University of Sassari, Italy
Alejandro M.S. Mayer, Midwestern University, Downers Grove, USA
Concetta Maria Messina, Department of Earth and Sea Sciences, University of Palermo, Italy
Vincenzo Mitolo, University "Aldo Moro", Bari, Italy
Amir Sasan Mozaffari Nejad, Hamadan University of Medical Sciences, Iran
Werner E.G. Muller, Johannes Gutenberg University, Mainz, Germany
Giuseppe Murdaca, University of Genoa, Italy
Giuseppe Palumbo, University Federico II, Naples, Italy
Gian Luigi Panattoni, University of Turin, Italy
Massimo Pregnolato, University of Pavia, Italy
Mark R. Rasenick, University of Illinois, Chicago, USA

Angela Maria Rizzo, University of Milan, Italy
Giacomo Rizzolatti, University of Parma, Italy
Aldo Rustioni, University of North Carolina, USA
Salvatore Sapienza, University of Catania, Italy
Pietro Scotto Di Vettimo, University of Naples, Italy
Vinicio Serino, University of Siena, Italy
Lynne Christine Weaver, University of Western Ontario, Canada
Ming Wei, Griffith University, Australia
Mario Wiesendanger, University of Friburg, Switzerland
Herbert Ryan Marini, University of Messina, Italy
Angela Marino, University of Messina, Italy

FOR AUTHORS

SUBMIT YOUR PAPER

[Guide for Authors](#)

[Benefits for Authors](#)

[How to write a scientific paper](#)

[How to write a Review article](#)

[Article Processing Charge](#)

FOR REVIEWERS

[Benefits for Reviewers](#)

[How to review](#)

[Thanks to Reviewers](#)

INDEXING

[PubMed](#)

[Scopus](#)

[DOAJ](#)

CAS

ESCI

MOST READ LAST MONTH

93rd National Congress of the Italian Society of Experimental...

 170

86th SIBS National Congress | Palermo, Italy, 24-25 October 2013

 62

87th SIBS National Congress on Peoples, Environment, Health |...

 60

Curcumin effects in inducing mRNA gene cathelicidin antimicrobial...

 38

Antibacterial and anti-inflammatory effects of lime (*Citrus...*

 38

KEYWORDS



Journal of Biological Research

is the official journal of the Società Italiana di Biologia Sperimentale and it is published online by [PAGEPress®](#), Pavia, Italy. All credits and honors to [PKP](#) for their [OJS](#).

eISSN 2284-0230
 pISSN 1826-8838

© PAGEPress 2008-2021

PAGEPress® is a registered trademark property of PAGEPress srl, Italy • VAT: IT02125780185 • [Privacy](#)

[ABOUT THE JOURNAL](#) [EDITORIAL BOARD](#) [CURRENT ISSUE](#) [ARCHIVES](#) [ANNOUNCEMENTS](#)


Search

[HOME](#) / [ARCHIVES](#) / Vol. 93 No. 2 (2020)

2020 Vol.93 No.2

Published: 2020-12-31

ORIGINAL ARTICLES



CURCUMIN EFFECTS IN INDUCING MRNA GENE CATHELIDICIN ANTIMICROBIAL PEPTIDE IN BALB/C MICE INFECTED WITH *SALMONELLA TYPHI*

Ami Febriza, Rosdiana Natzir, Mochammad Hatta, Cuno S.P.M Uiterwaal, Suryani As'ad, Budu, Gemini Alam, Vivien Novarina Kasim, Hasta Handayani Idrus

<https://doi.org/10.4081/jbr.0.8942>

PDF




EXAMINATION OF HEPATO-RENAL FUNCTIONS AND LIPID PANEL AMONG PREGNANT WOMEN WITH PREECLAMPSIA IN SÉTIF, ALGERIA

Romyla Bourouba, Kaouthar Benantar, Chaima Ayati, Lina Hedna, Kenza Gherbi, Farida Djabi

 <https://doi.org/10.4081/jbr.0.9035>

 PDF




ANTIBACTERIAL AND ANTI-INFLAMMATORY EFFECTS OF LIME (*CITRUS AURANTIFOLIA*) PEEL EXTRACT IN BALB/C MICE INFECTED BY *SALMONELLA TYPHI*

Vivien Novarina Kasim, Mochammad Hatta, Rosdiana Natzir, Veni Hadju, Yusminah Hala, Budu, Gemini Alam, Suryani As'ad, Ami Febriza, Hasta Handayani Idrus

 <https://doi.org/10.4081/jbr.0.8951>

 PDF




BIOLOGICAL INVESTIGATIONS ON THE ROLE OF HYDROGEL FORMULATIONS CONTAINING BIOACTIVE NATURAL AGENTS AGAINST SOME COMMON PHYTOPATHOGENS OF *PHASEOLUS VULGARIS* L. AND SEED GERMINATION

Hazem Salaheldin Elshafie, Maria Nuzzaci, Giuseppina Logozzo, Tania Gioia, Ippolito Camele

 <https://doi.org/10.4081/jbr.0.9219>

 PDF




THE EFFECT OF ARTIFICIAL SWEETENERS ON BODY WEIGHT OF MICE

Iyad Ali, Naser Shraim, Wafaa' Atrash, Aisha Sirafi, Huda Abadi

 <https://doi.org/10.4081/jbr.2020.8783>

 PDF




AMINO ACID COMPOSITION AND PROTEIN QUALITY OF TEPARY BEAN GRAINS (*PHASEOLUS ACUTIFOLIUS* L.) AS AFFECTED BY SOWING DATE AND PLANTING PATTERN UNDER SEMI-ARID CONDITION

Tabasom Ghadimian, Hamid Madani, Nourali Sajedi, Masoud Gomarian, Saeid Chavosh

 <https://doi.org/10.4081/jbr.2020.9111>

 PDF

INHIBITORY EFFECT OF PLANT EXTRACTS ON THE CYTOTOXICITY OF EURYTELE NEMATOCYSTS FROM *PELAGIA NOCTILUCA*



Rossana Morabito, Laura Cornara, Giuseppina La Spada, Angela Marino, Gian Luigi Mariottini, Alessia Remigante, Bruno Burlando

<https://doi.org/10.4081/jbr.2020.9136>

 PDF




CHIARI I MALFORMATION IN A WAARDENBURG PHENOTYPE WITH MULTIPLE MALFORMATIONS AND 1Q21.1 MICRODELETION

Agostino Berio, Gian Luigi Mariottini, Marco Frascio, Enrico Calcagno, Giacomo Garlaschi, Giuseppe Mangiante, Attilia Piazzì

<https://doi.org/10.4081/jbr.2020.9259>

 PDF



FIBRINOGENOLYTIC ACTIVITY OF PROTEASE FROM THE CULTURE FLUID OF *PLEUROTUS OSTREATUS*

Yevhenii M. Stohnii, Valeriia V. Sakovich, Volodymyr O. Chernyshenko, Volodymyr I. Chernishov, Tamara M. Chernyshenko, Iryna Kolesnikova, Yevhenii P. Kucheriavyi, Dmitrii D. Zhernossekov

<https://doi.org/10.4081/jbr.0.9006>

 PDF



EVALUATION OF ANTIDIABETIC ACTIVITY OF AQUEOUS EXTRACT OF *ORIGANUM FLORIBUNDUM* MUNBY LEAVES IN ALLOXAN INDUCED DIABETIC RATS WISTAR

M'hamed Nasri, Mohamed Zaouani, Noura Mimoune, Fatma Amira Hani, Mohamed Mahdid, Mohamed Toumi

<https://doi.org/10.4081/jbr.2020.9141>

 PDF



THE *PROCAMBARUS CLARKII* (RED SWAMP CRAYFISH) IN MERJA ZERGA WETLAND IN MOROCCO: DYNAMIC AND IMPACTS ON AQUATIC VEGETATION

SARA Saguem, Abdelouahab Sahli, EL Habib Jdi, Majida El Alami El Moutaouakil

<https://doi.org/10.4081/jbr.2020.8778>

 PDF

REVIEW ARTICLES




ADIPOKINES IN OBESITY AND METABOLIC DISEASES

Marco Giammanco, Herbert R. Marini, Socrate Pallio, Manfredi Marco Giammanco, Giovanni Tomasello, Francesco Carini, Fabio Venturella, Gaetano Leto, Maurizio La Guardia

 <https://doi.org/10.4081/jbr.2020.8915>

 PDF



METABOLIC EFFECTS OF 3,5-DIIDO-L-THYRONINE

Marco Giammanco, Manfredi Marco Giammanco, Gaetano Leto, Herbert R. Marini

 <https://doi.org/10.4081/jbr.2020.9182>

 PDF



ANTITUMORAL ACTIVITY OF CURCUMIN: AN ADJUVANT THERAPEUTIC STRATEGY

Giovanni Tomasello, Marco Giammanco, Margherita Mazzola, Provvidenza Damiani, William Ra', Paola Giammanco, Francesca Rappa, Alessandro Pitruzzella, Dario Saguto, Benedetto Di Trapani, Giovanni Taverna, Mike Carini, Antonio Ciulla, Francesco Carini

 <https://doi.org/10.4081/jbr.2020.8252>

 PDF

FOR AUTHORS

SUBMIT YOUR PAPER

Guide for Authors

Benefits for Authors

How to write a scientific paper

How to write a Review article

Article Processing Charge

FOR REVIEWERS

Benefits for Reviewers

How to review

Thanks to Reviewers

INDEXING

PubMed

Scopus

DOAJ

CAS

ESCI

MOST READ LAST MONTH

93rd National Congress of the Italian Society of Experimental...

 170

86th SIBS National Congress | Palermo, Italy, 24-25 October 2013

 62

87th SIBS National Congress on Peoples, Environment, Health |...

 60

Curcumin effects in inducing mRNA gene cathelicidin antimicrobial...

 38

Antibacterial and anti-inflammatory effects of lime (*Citrus...*

 38

KEYWORDS



Journal of Biological Research

is the official journal of the Società Italiana di Biologia Sperimentale and it is published online by [PAGEPress®](#), Pavia, Italy. All credits and honors to [PKP](#) for their [OJS](#).

eISSN 2284-0230

pISSN 1826-8838

© PAGEPress 2008-2021

PAGEPress® is a registered trademark property of PAGEPress srl, Italy • VAT: IT02125780185 • [Privacy](#)

Antibacterial and anti-inflammatory effects of lime (*Citrus aurantifolia*) peel extract in Balb/c mice infected by *Salmonella typhi*

Vivien Novarina Kasim,¹ Mochammad Hatta,² Rosdiana Natzir,³ Veni Hadju,⁴ Yusminah Hala,⁵ Budu,⁶ Gemini Alam,⁷ Suryani As'ad,⁸ Ami Febriza,⁹ Hasta Handayani Idrus¹⁰

¹Faculty of Medicine, Gorontalo State University, Gorontalo; ²Molecular Biology and Immunology Laboratory, Department of Microbiology; ³Department of Biochemistry, Faculty of Medicine; ⁴Department of Nutrition, Faculty of Public Health, Hasanuddin University, Makassar; ⁵Departement of Biology, Faculty of Mathematics and Natural Science, Makassar State University, Makassar; ⁶Department of Ophthalmology, Faculty of Medicine; ⁷Department of Pharmacognosy, Faculty of Pharmacy; ⁸Department of Clinical Nutrition, Faculty of Medicine, Hasanuddin University, Makassar; ⁹Department of Physiology, Faculty of Medicine, Muhammadiyah University Makassar, Makassar; ¹⁰Department of Microbiology, Indonesian Muslim University, Makassar, Indonesia

Abstract

Lime (*Citrus aurantifolia*) is a traditional plant that is widely used as antibacterial. This study proves the effect of Lime Peel

Extract (LPE) on the colonization and growth, mediated by the activity of IL-6, of bacteria *Salmonella typhi* in Balb/c mice. Mice were divided into four groups; LPE 510 mg/kg body weight (bw), LPE 750 mg/kgbw, and positive and negative control. The examination was carried out 3 times, on the 5th day before the intervention, on the 10th day after the intervention and on the 30th day after maintenance. Intervention of LPE for 5 days can decrease the number of *S. typhi* colonies, even maintenance for 20 days after the intervention showed no bacterial growth. IL-6 pro-inflammatory cytokine activity increased on examination day 5 after *S. typhi* injection and decreased after intervention on day 10, it was significantly different between pre and post at all groups except for negative controls ($p=0.15$). The speed of decrease in IL-6 levels was the greatest at the LPE 750 mg/kgbw (velocity=-5.64%). LPE decreased serum levels of IL-6 and inhibited the growth of *S. typhi* colony in Balb/c mice. LPE has potential for antibacterial and anti-inflammatory.

Correspondence: Vivien Novarina Kasim, Faculty of Medicine, Gorontalo State University, Jl. Sudirman No. 6, Dulalowo, Central City, Gorontalo, Gorontalo, Indonesia 96128.
Tel.: 081342419399, Fax: (0435) 821752.
E-mail: vivienovarina@ung.ac.id

Key words: Lime peel extract; *Citrus aurantifolia*, Interleukin 6; antibacterial; anti-inflammatory.

Acknowledgments: The authors would like to thank Markus, Rommy, Mus, and Wani (Molecular Biology and Immunology Laboratory for Infection Diseases, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia), who helped in the implementation of our research activities. This research was financially supported by the Ministry of Research and Technology, Indonesia or BUDI-DN LDPD.

Funding: Ministry of Research and Technology, Indonesia or BUDI-DN LDPD.

Conflict of interest: The authors declare no potential conflict of interests.

Ethics approval: This research was approved by the Health Medical Ethics Research Committee at the Faculty of Medicine, Hasanuddin University (Makassar, Indonesia) with registration number of 900/H4.8.4.5.31/PP36-KOMETIK/2018 on October 31, 2018.

Received for publication: 7 March 2020.
Accepted for publication: 19 July 2020.

©Copyright: the Author(s), 2020
Licensee PAGEPress, Italy
Journal of Biological Research 2020; 93:8951
doi:10.4081/jbr.2020.8951

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (by-nc 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Introduction

Some antibiotics such as ampicillin, chloramphenicol, tetracycline and co-trimoxazole become ineffective on *Salmonella typhi* and *Salmonella paratyphi*.¹ Multi-Drug Resistance (MDR) leads to new treatments in the form of adjuvant therapy which is a traditional plant that can be used as an antimicrobial. One of which is the lime plant (*Citrus aurantifolia*). From many previous studies even to date, all parts of *C. aurantifolia* have various efficacy.²⁻³ Specifically, it was emphasized that the peel of lime contains of active ingredients such as polyphenols, flavonoids, tannins, saponins, alkaloids, and triterpenoids.⁴ *Salmonella typhi*, is a Gram-negative bacterial pathogen whose transmission almost always occurs through the contaminated food and drinks. From several studies, the resistance of *Salmonella typhi* has begun to be high. The development of antimicrobial resistance is in line with the increasing use of antimicrobial drugs and in line with the discovery of new drugs.

Bacterial resistance and especially that of *Salmonella typhi* brings us to a new treatment in the form of adjuvant therapy, one of which is *Citrus aurantifolia* that can be used as an antimicrobial. Based on the previous research, all parts of *Citrus aurantifolia*; leaves, root stem, bark, and peel contain useful metabolic

compounds⁵ the lime peel has a higher concentration of flavonoids compared to other parts such as seed, fruit and juice.⁶ The existence of the content of flavonoids makes the lime peel has antibacterial and antioxidant power. Previous studies tested the bacteriostatic or bactericidal activity of flavonoids by conducting a time-kill study found that flavonoids do not kill bacterial cells but only induce the formation of bacterial aggregates thereby reducing the amount of Colony Forming Units (CFU) in a decent amount.⁷

When *S. typhi* bacterial infection occurs, there is a bond between the lipopolysaccharide (LPS) of the bacterial wall and the Toll-Like Receptor 4 (TLR 4) of the host, which stimulates innate immune activity. The activity releases a number of cytokines such as TNF α , IL-1, and IL-6. Interleukin 6 (IL-6) mediates the protective systemic effects of inflammation, including the effects of fever, acute-phase protein synthesis by the liver, and increased production of leukocytes by the bone marrow.⁸ This study discusses the effect of lime peel extract on the amount of bacterial colonization and its effect on the activity of IL-6 pro-inflammatory cytokines in Balb/c mice injected with *Salmonella typhi*.

Materials and Methods

An experimental design was used to study the effectiveness of LPE in decreasing pro-inflammatory IL-6 levels and bacterial colonization in Balb/c mice strains induced by *Salmonella typhi*.

Lime peel extract

Lime peel extraction was carried out at the Phytochemical Laboratory, Faculty of Pharmacy, Hasanuddin University, Makassar, Indonesia. Lime peel thick extracts were macerated with 96% Ethanol for 72 hours. The Lime Peel Extract (LPE) used in this study was consisted of two doses: 510 mg/kg body weight (bw) and 750 mg/kgbw, both dissolved with Aquades. LPE was obtained as reported in Kasim *et al.*⁴

Balb/c mice

Balb/c mice age ranging from 8 to 12 weeks, and weight from 30 to 40 grams (n=20), were obtained from Laboratory of Molecular Biology and Immunology, Department of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia. Mice were divided into four groups (n=five/group) based on intervention; LPE dose of 510 mg/kgbw, LPE 750 mg/kgbw, positive control (Levofloxacin dose 98 mg/kgbw),⁹ negative control (Aquades). All groups were injected intraperitoneally with *S. typhi* strain Thy1 (3×10^3 CFU/ml). 3 days after the injection of *Salmonella typhi*, each animal was started to be intervened for 5 days. Animals were treated as reported in Kasim *et al.*⁴

Bacterial colonization

Bacterial colonies were counted from samples taken from peritoneal fluid. Samples were taken three times, on the 5th day after mice were induced by *Salmonella typhi* (pre-intervention), the 10th post-intervention and 30th day after maintenance without treatment. Samples were taken as much as 0.5 mL and placed in 4.5 mL of physiological saline solution (0.9% NaCl). Dilution was carried out three times so that the culture obtained was not too dense or filled up the cup (the culture is too dense will interfere with observation). One mL of suspension was poured into a sterile Petri dish, then poured warm sterile (nutrient agar) media (45°C) then tightly

closed and incubated for 1-2 days at 37°C. The method used was the Plate Count Agar (PCA), a technique to grow microorganisms in agar media by mixing the liquid agar media with bacterial culture stock (agar) so that the cells are evenly distributed and still on the surface of agar or inside of agar.¹⁰

Interleukin 6 (IL-6) examination

Serum samples were taken 4 times, baseline (day 0), after induction of *S. typhi* before intervention (day 5), after intervention (10th day) and after 30 days. IL-6 antibody concentrations were determined by the IL6 ELISA Mouse Sandwich method. IL-6 examination was reported in Kasim *et al.*⁴

Ethics statement

This research was approved by the Health Medical Ethics Research Committee at the Faculty of Medicine, Hasanuddin University (Makassar, Indonesia) with registration number of 900/H4.8.4.5.31/PP36-KOMETIK/2018 on October 31, 2018.

Results

Bacterial colonization

The number of bacterial colonies after being given an intervention decreased in all groups, however on the 10th day of the negative control group, the number of bacterial colonies was seen more than other groups, and there were still bacterial colonies (Figure 1) on the 30th day of negative control group. On the 10th day, the number of bacterial colonies decreased after the intervention of lime peel extract which was given for 5 days of intervention. The number of bacterial colonies continued to decrease until the 30th day post-intervention. The decrease in the number of bacterial colonies after LPE administration had almost the same effect as the decrease in the number of colonies after levofloxacin administration (positive control). The effect of quinolone as a bactericidal,¹¹ in this study showed that the number of bacterial colonies after 30 days continued to decrease even they did not grow in the group given LPE 510, LPE 750, and positive control.

In the LPE group at a dose of 750 mg/kgbw, the number of bacterial colonies on day 10 had no growth, better than positive control group. The average decreased in the number of bacterial colonies in the LPE group 750 mg/kgbw from 17.2×10^3 CFU/mL to 0 or no bacterial growth ($p=0.00$) whereas in the positive control group decreased from 30.4×10^3 CFU/mL to 1.6×10^3 CFU/mL




Figure 1. Dynamics of changes in the number of bacterial colonization of *S. Typhi* by the groups during the observation period; LPE, lime peel extract; CFU, colony forming units.

($p=0.001$). In the negative control group, the number of bacterial colonies decreased from 22.6×10^3 CFU/mL to 6×10^3 CFU/mL, meaning that the number of bacterial colonies in the negative control group was still bigger than the other three groups, even though there was a decrease.

Interleukin 6 (IL-6)

The level of serum IL-6 at 4 times of data collection was analyzed through paired T-test to assess the dynamics of changes in levels of serum with respect to changes in observation time for each group, it can be seen in Table 1. The mean of IL-6 levels (delta) showed that the greatest decrease was found in the LPE 750 intervention group, this suggests that the LPE 750 intervention had a greater impact on decreased IL-6, compared to antibiotics (levofloxacin) intervention. The decrease of IL-6 levels in the LPE 510 intervention group was greater than the placebo group (negative control).

Figure 2 shows the velocity of the decrease of IL-6 averages between the time of observation after the intervention (10th day) and before the intervention (5th day). At the time of observation of the 5th day to the 10th day, it was seen that the velocity of decreasing IL-6 levels was the greatest in the LPE 750: 5.64%, and the smallest velocity of decline was in the negative control group: 4.67%. The velocity of decrease in IL-6 levels in the LPE 750 was more than the positive control group given Levofloxacin. It is assumed that the extract of lime peel in this study is effective as an anti-inflammatory.

Discussion

Bacterial colonization

Various parts of *Citrus aurantifolia* plant (leaves, stems, roots, barks, and peels) are evaluated through phytochemical screening and it is found out that they contain of various metabolic compounds. Extract of *Citrus aurantifolia* leaf has an antibacterial effect that has a high inhibitory zone in Gram-positive and Gram-negative bacteria.^{5,12} When *S. typhi* first enters the body, the bacteria are destroyed by macrophages. It is recognized by various receptors located on the surface of phagocytes. LPS in the *S. typhi* cell wall is a signal for macrophages to carry out the activation.¹³ In previous preliminary studies,⁴ it was found that the metabolic compounds contained in the extract of lime peel were polyphenols, flavonoids, alkaloids, tannins, saponins, and triterpenoids. Each of these compounds has the property to kill bacteria. These compounds attack

bacteria directly, causing bacterial cell death. This is indicated in the calculation of the number of bacterial colonies that are reduced even no growth after 5 days given the intervention of LPE.

Interleukin 6 (IL-6)

One of the earliest responses of the innate immune system to infection and tissue damage is the secretion of cytokines by tissue cells, which is very important for acute inflammation. When *S. typhi* first enters the body, the bacteria will be destroyed by macrophages. Bacteria will be recognized by various receptors located on the surface of phagocytes.⁸ Specific marker molecules for Gram-negative bacteria such as *S. typhi* are LPS, LPS will use TLR-4, a receptor that plays a role in observing and destroying *Salmonella typhi*. Activated TLR-4 will recruit the MyD88 adapter protein. Then MyD88 recruits IRAK4, IRAK1 and IRAK2. IRAK kinase then phosphorylates and activates the TRAF6 protein, allowing NF- κ B to dwell in the cell nucleus and activating transcription and causing induced inflammatory cytokines. Proinflammatory cytokines such as IL-1 β and IL-6, IFN- γ and TNF- α are synthesized and systemic inflammation occurs.^{8,14} It can be said that the inflammatory process by IL-6 cytokines is already running.

The results of this study (Figure 2) showed that LPE therapy gave a greater decrease in IL-6 levels compared to levofloxacin. It is indicated that LPE has an anti-inflammatory effect on IL-6.¹⁵ LPE, in addition to having an antibacterial effect, also has an anti-inflammatory effect, LPE has the potential for a more effective therapy. Researchers assume that the inhibitory effect of IL-6 is due to the activity of the polyphenol compounds contained in the extracts of lime peel. In line with this, Haseeb A. (2017) found that the potential inhibitory effect of polyphenols on IL-6 expression and oxidative stress through molecular mechanisms of activation of NF- κ B.¹⁶

The content of polyphenols and flavonoids contained in LPE of this study reveal that polyphenols are bioactive substances, which are likely to have a potential effect on the inflammatory response. It is known that flavonoids are one of the most extensive groups of secondary metabolite plants, flavonoids are found in many edible fruits and vegetables. The source of polyphenols is mostly contained in citrus fruits.¹⁷ The polyphenol content in lime peel extract in this study is 2.29% and flavonoids are 0.26%. The research by Rossano on *Citrus aurantifolia* in Calabria, Italy revealed that lime peels which were extracted by methanol solvent contained of more phenols (95.6 mg/g) and flavonoids (23.5 mg/g) compared to the leaves. In that study, it was found that there was a significant relationship between phenol and flavonoid with antioxidant activity.³

Table 1. Mean values of IL-6 (delta) differences between all groups at 5th day (pre-intervention), 10th day (post-intervention) and 30th day.

Groups	Level of IL-6 (rg/ml)	
	5 th -10 th day Delta (mean \pm SD)	10 th -30 th day Delta (mean \pm SD)
LPE 510	19,63 \pm 7,06	53,98 \pm 43,38
LPE 750	24,54 \pm 7,71	115,61 \pm 56,92
Control (+)	24,07 \pm 7,06	273,41 \pm 68,26
Control (-)	11,62 \pm 6,33	41,91 \pm 18,87
P value	0,074	0,000

IL-6, interleukin 6; LPE, Lime Peel Extract; Control(+), positive control group; Control(-), negative control group; SD, Standard Deviasi.




Figure 2. The velocity of decrease (%) of IL-6 levels in each group after the intervention; LPE: Lime Peel Extract; Control (+): positive control group; Control (-): negative control group.

Tejada S. (2017) proved that hesperidin (a type of flavonoid) extracted from the citrus genus, has potential as an anti-inflammatory. Hesperidin treatment decreases inflammatory mediators and provides a significant antioxidant effect. Molecular bases for anti-inflammatory effects appear to be mediated by signaling pathways especially the nucleus $\kappa\beta$ factor pathway.¹⁸ As for other opinions, a research conducted by Jorge L.A, found that from 3 types of citrus namely *C. limon*, *C. aurantifolia*, *C. limonia*, Essential Oils (EO) contained in the 3 types of citrus have anti-inflammatory effects, especially the limonia content of essential oils from the citrus fruit skin.¹⁵ During the inflammatory event, there is an increase in cytokine production.

Conclusions

This study showed that LPE decreased serum levels of IL-6 and bacterial colonization in Balb/c mice induced by *Salmonella typhi*. LPE has potential as antibacterial and anti-inflammatory.

References

1. Roka G, Pandaya S, Ferdous MR, et al. Susceptibility to patterns of ciprofloxacin among nalidixic acid-resistant *Salmonella* isolates collected in Banepa, Nepal from enteric fever patients. *J Biol Res – Boll Soc Ital Biol Sper* 2016;89:47–50.
2. Rani N, Bharti S, Krishnamurthy B, et al. Pharmacological properties and therapeutic potential of naringenin: acitrus flavonoid of pharmaceutical promise. *Curr Pharm Des* 2016;22:4341–59.
3. Loizzo MR, Tundis R, Bonesi M, et al. Evaluation of *Citrus aurantifolia* peel and leaves extracts for their chemical composition, antioxidant and anti-cholinesterase activities. *J Sci Food Agric* 2012;92:2960–7.
4. Kasim VNA, Hatta M, Febriza A, et al. Lime peel extract effects in decreasing levels of inteleukin 6 in mice infected with *Salmonella typhi*. Topcuoglu, Bulent, Ahmadi R, eds. International Conference on BioMedical Sciences, ICBMS 2019. Istanbul, Turkey; September 27-28, 2019. pp. 69–78.
5. Pathan RK, Gali PR, Pathan P, et.al. In vitro antimicrobial activity of *Citrus aurantifolia* and its phytochemical screening. *Asian Pacific J Trop Dis* 2012;2:S328–3.
6. Lemes RS, Alves CCF, Estevam EBB, et al. Chemical composition and antibacterial activity of essential oils from *Citrus auranti folia* leaves and fruit peel against oral pathogenic bacteria. *An Acad Bras Cienc* 2018;90:1285–92.
7. Cushnie TPT, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 2005;26:343–56.
8. Abbas AK, Lichtman AH. *Celluler and moleculer immunology*. 8th ed. Philadelphia: W.B. Saunders Company; 2014.
9. Kalra SP, Naithani N, Mehta SR, et.al. Current trends in the management of typhoid fever. *Med J Armed Forces India* 2003;59:130–5.
10. Harley J, Prescott L. *Laboratory exercises in microbiology*. 50th ed. Lab Exerc Microbiol: The McGraw-Hill Companies; 2002.
11. Wald-Dickler N, Holtom P, Spellberg B. Busting the myth of “static vs cidal”: asystemic literature review. *Clin Infect Dis* 2018;66:1470–4.
12. Aibinu I, Adenipekun T, Adelowotan T, et.al. Evaluation of the antimicrobial properties of different parts of *Citrus aurantifolia* (lime fruit) as used locally. *African J Tradit Complement Altern Med* 2007;4:185–90.
13. Abbas AK, Lichtman AH, Pillai S. *Basic immunology functions and immune system disorders*. Indo ed. Singapore: Elsevier (Singapore) Pte td; 2014.
14. Kaarthikeyan G, Balakrishnan A, Jayakumar ND. The link between the genetic polymorphisms of the innate immune signaling molecular factors with periodontitis. *J Biol Res – Boll Soc Ital Biol Sper* 2018;91:53–5.
15. Amorim JL, Simas DLR, Pinheiro MMG, et al. Anti-inflammatory properties and chemical characterization of the essential oils of four citrus species. *PLoS One* 2016;11:e0153643.
16. Pérez-Cano F, Massot-Cladera M, Rodríguez-Lagunas M, Castell M. Flavonoids affect host-microbiota crosstalk through tlr modulation. *Antioxidants*.2014;3:649–70.
17. Ghasemi K, Sciences SA, Ghasemi Y, Ebrahimzadeh MA. Antioxidant activity, phenol and flavonoid contents of 13 *Citrus* species peels and tissues. *Pak J Pharm Sci* 2009;22:277-81.
18. Tejada S, Pinya S, Martorell M, et al. Potential anti-inflammatory effects of hesperidin from the genus *Citrus*. *Curr Med Chem* 2018;25:4929–45.