



## Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.


The first page of your submissions is displayed below.

Submission author: Vivien Novarina Kasim  
Assignment title: Plagiarism Checker 1  
Submission title: The Role of IL-6, TNF- $\alpha$ , and VDR in Inhibiting the Growth of ...  
File name: TOMICROJ-14-65.pdf  
File size: 536.04K  
Page count: 7  
Word count: 5,922  
Character count: 30,349  
Submission date: 03-Jul-2021 10:13PM (UTC-0500)  
Submission ID: 1615416430


1574-2682/20

Send Orders for Reprints to [support@ambdjournal.net](mailto:support@ambdjournal.net)

46



**The Open Microbiology Journal**  
Content list available at: <https://openmicrobiologyjournal.com>



**RESEARCH ARTICLE**  
**The Role of IL-6, TNF- $\alpha$ , and VDR in Inhibiting the Growth of *Salmonella Typhi* in vivo Study**  
Ami Febriza<sup>1</sup>, Rosdiana Natriz<sup>1</sup>, Mochammad Hatta<sup>1</sup>, Suryani A'sad<sup>1</sup>, Budi<sup>1</sup>, Cahyono Kaelan<sup>1</sup>, Vivien Novarina Kasim<sup>1</sup> and Hasta Handayani Idrus<sup>2</sup>

<sup>1</sup>Department of Physiology, Faculty of Medicine, University of Muhammadiyah Makassar, Makassar, Indonesia  
<sup>2</sup>Biochemistry Department, Faculty of Medicine, Hasanudin University, Makassar, Indonesia  
<sup>3</sup>Molecular Biology and Immunology Laboratory, Faculty of Medicine, Hasanudin University, Makassar, Indonesia  
<sup>4</sup>Clinical Nutrition Department, Faculty of Medicine, Hasanudin University, Makassar, Indonesia  
<sup>5</sup>Department of Ophthalmology, Faculty of Medicine, Hasanudin University, Makassar, Indonesia  
<sup>6</sup>Department of Pathology, Faculty of Medicine, Hasanudin University, Makassar, Indonesia  
<sup>7</sup>Faculty of Medicine, Universitas Negeri Gorontalo, Gorontalo, Indonesia  
<sup>8</sup>Department of Microbiology, Faculty of Medicine, University Muslim Indonesia, Makassar, Indonesia

**Abstract:**  
**Background and aim:**  
The prevalence of typhoid fever is reportedly high, especially in Asia. When a pathogen enters the human body, there are markers in the form of molecules that will be known by the innate immune system. Specific molecular markers of gram negative bacteria, which are Lipopolysaccharides (LPS) and flagellin receptors will interact with LPS. The binding between LPS and TLR-4 will give rise to activation signals that will activate innate immune cells. Innate cells will release a number of proinflammatory cytokines, such as TNF- $\alpha$ , IL-1, and IL-6. While Vitamin D Receptors (VDR) are expressed in large amounts in tumor tissue and infected cells. This study aimed to prove the role of IL-6, TNF- $\alpha$ , and VDR in inhibiting bacterial growth in mice that have been infected by *S. Typhi*.

**Methods:**  
This research was a real experimental pre-post test design to investigate the level of IL-6, TNF- $\alpha$  and VDR in suppressing the growth of bacteria in the peritoneal fluid of *S. Typhi* mice, male BALB/c. Mice were divided into three groups consisting of 10 mice each. All mice in groups A and B were intraperitoneally inoculated with *S. Typhi* strain Ty1 in study day 0. Group A was treated with antibiotic Levofloxacin, on study day 4<sup>th</sup>. Another study group, group B, was used as a placebo and received aquades on study day 4<sup>th</sup>. While group C as a control was not inoculated with *S. Typhi*. Blood samples from three groups for the calculation of serum IL-6, TNF- $\alpha$ , and VDR were collected. This examination was taken four times, at baseline, 4<sup>th</sup> day, 10<sup>th</sup> day, and 30<sup>th</sup> day. For the calculation of bacterial colony, peritoneal fluid retrieval was collected three times, which is on 4<sup>th</sup> day, 10<sup>th</sup> day, and 30<sup>th</sup> day.

**Results:**  
A repeated measure ANOVA in group A (antibiotic) and group B (placebo) group showed that mean IL-6, TNF- $\alpha$ , and VDR level differed statistically significant between times (p-value 0.000). There was a strong negative correlation between bacterial colony count and VDR level, which was statistically significant in both groups (group A,  $r = -0.875$ , p-value = 0.000 vs group B,  $r = -0.470$ , p-value = 0.002). IL-6 and TNF- $\alpha$  did not give significant statistical correlation with bacterial colony count.

**Conclusion:**  
VDR, IL-6, and TNF- $\alpha$  play an important role in killing bacteria. From the results of this study, IL-6 level is related to the number of bacterial colonies, the lower the IL-6 level, the less the number of bacterial colonies. Similarly, TNF- $\alpha$  levels have a positive correlation with the number of bacterial colonies. While VDR levels are also related to the number of bacterial colonies, the higher the VDR level, the lower the number of bacterial colonies.

**Keywords:** Bacterial colonies, Colony count, IL-6, TNF- $\alpha$ , *Salmonella*, VDR, *S. Typhi*.

**Article History** Received: January 07, 2020 Revised: March 06, 2020 Accepted: March 08, 2020

DOI: 10.21767/16154164301610065.2020.14.65-71

# The Role of IL-6, TNF- $\alpha$ , and VDR in Inhibiting the Growth of Salmonella Typhi: in vivo Study

*by Vivien Novarina Kasim*

---

**Submission date:** 03-Jul-2021 10:13PM (UTC-0500)

**Submission ID:** 1615416430

**File name:** TOMICROJ-14-65.pdf (536.04K)

**Word count:** 5922

**Character count:** 30349



# The Open Microbiology Journal

Content list available at: <https://openmicrobiologyjournal.com>



## RESEARCH ARTICLE

### The Role of IL-6, TNF- $\alpha$ , and VDR in Inhibiting the Growth of *Salmonella Typhi*: *in vivo* Study

Ami Febriza<sup>1,\*</sup>, Rosdiana Natzir<sup>2</sup>, Mochammad Hatta<sup>3</sup>, Suryani As'ad<sup>4</sup>, Budu<sup>5</sup>, Cahyono Kaelan<sup>6</sup>, Vivien Novarina Kasim<sup>7</sup> and Hasta Handayani Idrus<sup>8</sup>

<sup>1</sup>Department of Physiology, Faculty of Medicine, University of Muhammadiyah Makassar, Makassar, Indonesia

<sup>2</sup>Biochemistry Department, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>3</sup>Molecular Biology and Immunology Laboratory, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>4</sup>Clinical Nutritional Department, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>5</sup>Department of Ophthalmology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>6</sup>Department of Pathology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>7</sup>Faculty of Medicine, University Negeri Gorontalo, Gorontalo, Indonesia

<sup>8</sup>Department of Microbiology, Faculty of Medicine, University Muslim Indonesia, Makassar, Indonesia

#### Abstract:

##### Background and aim:

The prevalence of typhoid fever is reportedly high, especially in Asia. When a pathogen enters the human body, there are markers in the form of molecules that will be known by the innate immune system. Specific molecular markers of gram negative bacteria, which are Lipopolysaccharides (LPS) and Toll-Like receptors-4 will interact with LPS. The binding between LPS and TLR-4 will give rise to activation signals that will activate innate immune cells. Immune cells will release a number of proinflammatory cytokines, such as TNF- $\alpha$ , IL-1, and IL-6. While Vitamin D Receptors (VDR) are expressed in large amounts in tumor tissue and infected cells. This study aimed to prove the role of IL-6, TNF- $\alpha$ , and VDR in inhibiting bacterial growth in mice that have been induced by *S. Typhi*.

##### Methods:

This research was a real experimental pre-post test design to investigate the level of IL-6, TNF- $\alpha$  and VDR in suppressing the growth of bacteria in the peritoneal fluid of *S. Typhi*, male, mice BALB/c. Mice were divided into three groups comprised of 10 mice each. All mice in groups A and B were intraperitoneally inoculated with *S. Typhi* strain Thy1 in study day 0. Group A was treated with antibiotic Levofloxacin, on study day 4<sup>th</sup>. Another study group, group B, was used as a placebo and received aquades on study day 4<sup>th</sup>. While group C as a control was not inoculated with *S. Typhi*. Blood samples from three groups for the calculation of serum IL-6, TNF- $\alpha$ , and VDR were collected. This examination was taken four times; at baseline, 4<sup>th</sup> day, 10<sup>th</sup> day, and 30<sup>th</sup> day. For the calculation of bacterial colony, peritoneal fluid retrieval was collected three times, which is on 4<sup>th</sup> day, 10<sup>th</sup> day, and 30<sup>th</sup> day.

##### Results:

A repeated measure ANOVA in group A (antibiotic) and group B (placebo) group showed that mean IL-6, TNF- $\alpha$ , and VDR level differed statistically significant between times (*p*-value 0.000). There was a strong negative correlation between bacterial colony count and VDR level, which was statistically significant in both groups (group A;  $r = -0.875$ , *p*-value = 0.000 vs group B;  $r = -0.470$ , *p*-value = 0.002). IL-6 and TNF- $\alpha$  didn't give significant statistical correlation with bacterial colony count.

##### Conclusion:

VDR, IL-6, and TNF- $\alpha$  play an important role in killing bacteria. From the results of this study, IL-6 level is related to the number of bacterial colonies, the lower the IL-6 level, the less the number of bacterial colonies. Similarly, TNF- $\alpha$  levels have a positive correlation with the number of bacterial colonies. While VDR levels are also related to the number of bacterial colonies, the higher the VDR level, the lower the number of bacterial colonies.

**Keywords:** Bacterial colonies, Colony count, IL-6, TNF- $\alpha$ , Salmonella, VDR, *S. Typhi*.

#### Article History

Received: January 01, 2020

Revised: March 06, 2020

Accepted: March 08, 2020

## 1. INTRODUCTION

Typhoid fever is a systemic infection due to *Salmonella Typhi* (*S. Typhi*). In 2011, *S. Typhi* was estimated to infect 21.7 million people and caused 217,000 deaths worldwide. High incidence of typhoid fever is found in South Asia, Southeast Asia, and South Africa, as many as 80% of cases come from slums in Bangladesh, China, India, Indonesia, Laos, Nepal, Pakistan, and Vietnam [1].

In Indonesia, the prevalence of typhoid fever based on the diagnosis by health professionals and respondent complaints is 1.60%. A total of 14 provinces have Typhoid prevalence above the national prevalence, namely Nanggroe Aceh Darussalam, Bengkulu, West Java, Central Java, Banten, West Nusa Tenggara, East Nusa Tenggara, South Kalimantan, East Kalimantan, Central Sulawesi, South Sulawesi, Gorontalo, West Papua and Papua [2].

*Salmonella Typhi* is a gram-negative bacillus that causes typhoid fever in humans. This bacterium can survive in the phagosome so that it can escape the body's immune system. Some complications of typhoid fever are ileal perforation, bacteremia, and endovascular infection [3].

When a pathogen enters the human body, there are markers in the form of molecules that will be known by the human defense (innate immune system). Specific molecular markers of gram negative bacteria are lipopolysaccharides (LPS) and Toll-Like Receptors 4 (TLR4) will interact with bacteria (LPS) [4]. The binding between LPS and TLR-4 will give rise to activation signals that will activate innate immune cells. Immune cells will release a number of proinflammatory cytokines, such as TNF- $\alpha$ , IL-1 and IL-6.

IL-6 functions in non-specific and specific immunity produced by mononuclear phagocytes, endothelial cells, vascular, fibroblasts, and other cells in response to other microbes and cytokines. IL-6 is an important member of the cytokine family. It can induce differentiation of T and B lymphocytes. It can also promote the synthesis of acute-phase response proteins by stimulating liver cells. In numerous cytokines, some are very similar in terms of their biological activity [5]. Another study has found that IL-6 levels increase significantly in the early stage of inflammation, which provides evidence for rapid diagnosis and differential diagnosis of early bacterial infection in the clinic [6].

TNF- $\alpha$  plays a role in host defense against bacterial, viral, and parasitic infections. TNF- $\alpha$  is produced by macrophages and is activated by lymphocyte T cells, antigens, NK cells, and mast cells [7]. TNF- $\alpha$  is usually not detected in healthy individuals but is often found in conditions of inflammation and infection in the serum. TNF- $\alpha$  acts on leukocytes and endothelial, inducing acute inflammation at low levels because TNF- $\alpha$  is a strong pyrogen [8, 9].

Vitamin D receptors or VDR are expressed in large amounts in tumor tissue and infected cells. Recent research

shows that VDR and enzymes involved in vitamin D metabolism have damage to the VDR signaling pathway. VDR expression by immune cells shows that vitamin D influences the immune system function. More than 30 different body tissues, such as the brain, liver, and pancreas, lymphatics, skin, gonads and prostate are composed of cells including T and B lymphocytes that express VDR. Vitamin D The receptor binds to 1,25-hydroxyvitamin D, the active form of vitamin D and mediates its biological activity [10]. Based on various previous theories, this study aimed the role of IL-6, TNF- $\alpha$ , and VDR in inhibiting bacterial growth in mice that have been induced by *S. Typhi*.

## 2. MATERIALS AND METHODS

This research is a true experimental pre-post test design to investigate the level of IL-6, TNF- $\alpha$ , and VDR in suppressing the growth of bacteria in the peritoneal fluid of *S. Typhi* strain male mice BALB/c.

### 2.1. Experimental Animals

BALB/c mice (male, aged 8-12 weeks, weighing 30-40 grams; n = 30) were maintained in the Molecular Biology and Immunology Laboratory, Microbiology Department Faculty of Medicine, Hasanuddin University (Makassar, Indonesia). The mice were acclimatized for 8 days. Randomization was used as a sampling technique. Mice were randomly put into three study groups of 10 mice each. Two groups (A and B) were intraperitoneally induced with *S. Typhi* strains thyl (3 mL x 10<sup>3</sup> CFU/mL). Group A was treated with antibiotic levofloxacin once a day for five days in a dose of 1.95 mg/kg, and group B received placebo treatment. Control group C did not receive either *S. Typhi* nor antibiotic.

### 2.2. Levofloxacin

Levofloxacin was obtained from Kimia Farma, Pharmaceutical, Indonesia. Dose 750 mg of Levofloxacin given to mice was obtained from the multiplication with a conversion factor of 0.0026. Based on the result, the positive control group was given a dose of Levofloxacin 1.95 mg/day. Antibiotics was dissolved in distilled water and given through the nasogastric sonde, once a day for five days.

### 2.3. Sampling of Peritoneal Fluid And Bacterial Colony Examination

Mice were fixed in the supine position, the abdomen is cleaned with alcohol 70% and as much as 0.8-1 mL saline was injected into the peritoneal cavity. Mice were then allowed to stand for 1 minute as he rocked slowly. Peritoneal fluid removed from the peritoneal cavity of mice supine position, then fluid aspirated by as much as 0.5 mL syringe. Peritoneal fluid retrieval is performed three times, which is on the fourth day (4<sup>th</sup> day), on the tenth day (10<sup>th</sup> day) and 3 weeks after the mice induced with *S. Typhi* (30<sup>th</sup> day).

Examination of bacterial colonization by using the pour plate method. This method is performed by diluting the peritoneal fluid samples of 0.5 mL in 4.5 mL of saline (0.9% NaCl). Dilution is done three times so that the culture obtained is not too dense or fulfill cup (culture too dense will interfere

\* Address correspondence to this author at the Department of Physiology, Faculty of Medicine, University of Muhammadiyah Makassar, Makassar, Indonesia, Tel: (0411) 840199, Fax: (0411) 865588, Mobile: 085256655987 / 087840485864 (WA) E-mail: amifebriza@med.unismuh.ac.id.

with observations). Approximately, 1 mL of the suspension was poured into a sterile petri dish, followed by pouring the fertilizer medium (nutrient agar) sterile warm (45C), then sealed and incubated for 1-2 days at a temperature of 37C.

#### 2.4. IL-6, TNF- $\alpha$ And VDR Examination

Blood samples from two groups for the determination of serum IL-6, TNF- $\alpha$ , and VDR were collected. This examination was taken four times; at baseline, 4<sup>th</sup> day, 10<sup>th</sup> day and 30<sup>th</sup> day. The examination was done using IL-6 ELISA kit (Sandwich ELISA), TNF- $\alpha$  ELISA Kit (Sandwich ELISA), VDR ELISA Kit (Sandwich ELISA) for mouse, purchased from LSBio.

#### 2.5. Statistical Analysis

The data obtained is processed using SPSS 23 (IBM Corporation, New York, USA) for Macbook. All data are presented in the form of a mean and standard deviation. Before testing the difference hypothesis, first, we test the data normality and homogeneity. One-way ANOVA is used to calculate the mean difference of IL-6, TNF- $\alpha$  and VDR level between antibiotic group, placebo group, and control group at baseline. Because in this study, each group is measured multiple times to see changes to specific intervention, so we also used ANOVA with a repeated measure to calculate the differences means of bacterial colony count, IL-6, TNF- $\alpha$  and VDR level in the same groups in multiple time observation. ANOVA repeated measure would be followed by Bonferroni post hoc test to identify which particular differences pairs of groups means are significant. Considered as statistical significance was  $p$ -value <0.05. A Pearson correlation test was used to determine the correlation between bacterial colony count with VDR, IL-6, and TNF- $\alpha$ . Linear regression analysis was used to determine the most influential variables in determining the bacterial colony count.

### 3. RESULTS

This study consisted of 30 mice, which were divided into 3 groups (n = 5). Data on the characteristics of mice can be seen in Table I. No significant differences were found for age, body weight, VDR, IL-6 and TNF- $\alpha$  level at baseline (Table I).

The result found that at baseline, there was no statistically significant difference between VDR level in three groups. Both IL-6 and TNF- $\alpha$  levels also give no statistically significant differences in mean (Table I).

A repeated measure ANOVA in group A (antibiotic) and group B (placebo) showed that mean VDR level differed statistically significant between times ( $p$ -value 0.000), while in control group C showed no significant differences ( $p$ -value 0.495). There are significant differences in the average increase in VDR levels over time in group A (antibiotic) and group B (placebo). Increased levels of VDR were observed to be greater in the group with antibiotics compared to the placebo group and control group. Post hoc tests using Bonferroni correction revealed that in group A (antibiotic) shows an increase in VDR

level from 4<sup>th</sup> day to 10<sup>th</sup> day (6.95 $\pm$ 1.32 ng/mL vs. 15.04 $\pm$ 2.37 ng/mL,  $p$ -value 0.000), which was statistically significant. In group B (placebo), post hoc tests using Bonferroni correction revealed that in this group shows an insignificant decrease in VDR level from 4<sup>th</sup> day to 10<sup>th</sup> day (7.95 $\pm$ 1.71 ng/mL vs. 7.93 $\pm$ 1.42 ng/mL,  $p$ -value 1.000). Therefore, we can conclude that the group with given antibiotics can induce increase of VDR level more significant than with placebo (Table 2).

A repeated measure ANOVA in group A (antibiotic) and group B (placebo) showed that the mean IL-6 level differed statistically significant between times ( $p$ -value 0.000), while in control group C showed no significant differences ( $p$ -value 0.055). Post hoc tests using Bonferroni correction revealed that in group A (antibiotic) shows an increase in IL-6 level from baseline day to 4<sup>th</sup> day (109.25 $\pm$ 29.21 pg/mL vs 481.72 $\pm$ 44.10 pg/mL,  $p$ -value 0.000), which was statistically significant. The same result was found in the group B (placebo), it shows an increase in IL-6 level from baseline day to 4<sup>th</sup> day (125.71 $\pm$ 40.12 pg/mL vs 248.87 $\pm$ 32.75 pg/mL,  $p$ -value 0.000). Given antibiotics for five days, 4<sup>th</sup> day to 10<sup>th</sup> day, showed nonsignificant decreasing in IL-6 (481.72 $\pm$ 44.10 pg/mL vs 465.58 $\pm$ 45.85 pg/mL,  $p$ -value 1.000). In group B (placebo), post hoc tests using Bonferroni correction revealed that an insignificant decrease in IL-6 level from 4<sup>th</sup> day to 10<sup>th</sup> day (248.87 $\pm$ 32.75 pg/mL vs 222.84 $\pm$ 37.20 pg/mL,  $p$ -value 0.326) is shown in this group. Therefore, we can conclude that a group that induced infection can induce increasing IL-6 level. After the intervention, both groups given antibiotic and placebo showed a decrease in IL-6 (Table 2).

A repeated measure ANOVA in group A (antibiotic) and group B (placebo) showed that the mean TNF- $\alpha$  level differed statistically significant between times ( $p$ -value 0.000). In contrast, in control group C showed insignificant differences ( $p$ -value 0.801). Post hoc tests using Bonferroni correction revealed that in group A (antibiotic) shows an increase in TNF- $\alpha$  level from baseline day to 4<sup>th</sup> day (96.62 $\pm$ 21.90 pg/mL vs 304.88 $\pm$ 38.31 pg/mL,  $p$ -value 0.000). The same result was found in the group B (placebo), it shows an increase in TNF- $\alpha$  level from baseline day to 4<sup>th</sup> day (98.96 $\pm$ 25.95 pg/mL vs 198.46 $\pm$ 34.65 pg/mL,  $p$ -value 0.001). Post hoc tests using Bonferroni correction revealed that in antibiotic group shows a decrease in TNF- $\alpha$  level from 4<sup>th</sup> day to 10<sup>th</sup> day (304.88 $\pm$ 38.31 pg/mL vs 294.01 $\pm$ 24.34 pg/mL,  $p$ -value 1.000), which was statistically not significant. In group B (placebo), post hoc tests using Bonferroni correction revealed that in this group also shows insignificant decrease in TNF- $\alpha$  level from 4<sup>th</sup> day to 10<sup>th</sup> day (198.46 $\pm$ 34.65 pg/mL vs 183.73 $\pm$ 10.80 pg/mL,  $p$ -value 0.929). Therefore, we can conclude that group that induced with infection can induce increasing TNF- $\alpha$  level. After the intervention, both groups were given antibiotics and placebo showed a decrease in TNF- $\alpha$  (Table 2).

Fig. (1) showed that there was an increase in VDR levels in group A (antibiotic) and group B (placebo), with reasonably wide differences in mean. While IL-6 and TNF- $\alpha$  levels increased on the 4<sup>th</sup> day and then decreased after the 10<sup>th</sup> day.






Fig. (1). Dynamics of changes in VDR, IL-6, and TNF- $\alpha$  in all groups during the observation period

Table 1. Characteristics of experimental animals.

Variable	Groups			p-value
	Group A (antibiotic) (n=10)	Group B (placebo) (n=10)	Group C (control) (n=10)	
Strain	BALB/c	BALB/c	BALB/c	-
Sex	male	male	male	-
Age (weeks)	10.00 $\pm$ 1.49	9.60 $\pm$ 1.43	9.90 $\pm$ 1.28	0.804
Body weight (grams)	34.54 $\pm$ 3.45	35.72 $\pm$ 2.23	34.36 $\pm$ 3.65	0.614
VDR level (ng/mL) at baseline	2.98 $\pm$ 1.09	3.82 $\pm$ 1.81	2.94 $\pm$ 1.07	0.275
IL-6 level (pg/mL) at baseline	109.25 $\pm$ 29.21	125.71 $\pm$ 40.12	103.55 $\pm$ 22.98	0.282
TNF- $\alpha$ level (pg/mL) at baseline	96.62 $\pm$ 21.90	98.96 $\pm$ 25.95	95.50 $\pm$ 20.01	0.909

Values are expressed in mean $\pm$ SD, p-value of <0.05 is considered significant for the comparison between groups with One-Way ANOVA test.

Table 2. Comparison in means of VDR, IL-6 and TNF- $\alpha$  level in each group based on the observation time.

Groups	VDR level (ng/mL)				p value*
	Baseline	4 <sup>th</sup> day	10 <sup>th</sup> day	30 <sup>th</sup> day	
Group A (antibiotic)	2.98 $\pm$ 1.09	6.95 $\pm$ 1.32	15.04 $\pm$ 2.37	16.35 $\pm$ 2.27	0.000
Group B (placebo)	3.82 $\pm$ 1.81	7.95 $\pm$ 1.71	7.93 $\pm$ 1.42	9.28 $\pm$ 1.51	0.000
Group C (Control)	2.94 $\pm$ 1.07	3.09 $\pm$ 1.42	2.63 $\pm$ 1.12	3.06 $\pm$ 0.96	0.495
Groups	IL-6 level (pg/mL)				p value*
Group A (antibiotic)	109.25 $\pm$ 29.21	481.72 $\pm$ 44.10	465.58 $\pm$ 45.85	198.90 $\pm$ 31.49	0.000
Group B (placebo)	125.71 $\pm$ 40.12	248.87 $\pm$ 32.75	222.84 $\pm$ 37.20	197.34 $\pm$ 39.89	0.000
Group C (Control)	103.55 $\pm$ 22.98	112.25 $\pm$ 27.96	89.63 $\pm$ 19.68	85.71 $\pm$ 17.27	0.055
Groups	TNF- $\alpha$ level (pg/mL)				p value*
Group A (antibiotic)	96.62 $\pm$ 21.90	304.88 $\pm$ 38.31	294.01 $\pm$ 24.34	168.35 $\pm$ 19.09	0.000
Group B (placebo)	98.96 $\pm$ 25.95	198.46 $\pm$ 34.65	183.73 $\pm$ 10.80	157.77 $\pm$ 21.59	0.000
Group C (Control)	95.50 $\pm$ 20.01	95.60 $\pm$ 21.86	91.13 $\pm$ 16.95	93.98 $\pm$ 15.26	0.801

\* Repeated Measure ANOVA Test, with values are mean $\pm$ SD n=5, p-value < 0.05 is considered significant

**Table 3. Comparison in means of bacterial colony count in each group based on the observation time.**

Groups	Bacterial Colony Count (CFU/mL)			
	4 <sup>th</sup> day	10 <sup>th</sup> day	30 <sup>th</sup> day	<i>p</i> value*
Group A (antibiotic)	20.20 $\pm$ 4.96	0.70 $\pm$ 1.33	0.00 $\pm$ 0.00	0.000
Group B (placebo)	21.90 $\pm$ 5.93	6.10 $\pm$ 2.07	1.10 $\pm$ 1.10	0.000

\* Repeated Measure ANOVA Test, with values are mean $\pm$ SD n=5, *p*-value < 0.05 is considered significant

**Table 4. Correlation between bacterial colony count with VDR, IL-6, and TNF- $\alpha$  in both groups.**

Variable	Group A (antibiotic)		Group B (placebo)	
	Bacterial colony (CFU/mL)		Bacterial colony (CFU/mL)	
	<i>r</i> -value	<i>p</i> value*	<i>r</i> -value	<i>p</i> value*
VDR (ng/mL)	-0.875	< 0.001	-0.470	< 0.05
IL-6 (pg/mL)	-0.069	> 0.05	-0.203	> 0.05
TNF- $\alpha$ (pg/mL)	-0.150	> 0.05	-0.180	> 0.05

\* Data expressed as correlation coefficient (*r*-value) and *p*-value, *p*-value < 0.05 is considered significant. Bacterial colony count was dependent variable.

**Table 5. Linear regression analysis on VDR, IL-6 and TNF- $\alpha$  for bacterial colony count in group A (antibiotic) and B (placebo).**

Variable	Group A (antibiotic)			Group B (placebo)		
	Standard Error	B-Coefficients	<i>p</i> value*	Standard Error	B-Coefficients	<i>p</i> value*
VDR (ng/mL)	0.145	-1.623	0.000	0.752	-0.557	0.005
IL-6 (pg/mL)	0.012	0.016	0.190	0.037	0.066	0.757
TNF- $\alpha$ (pg/mL)	0.023	-0.014	0.557	0.047	0.086	0.681

A repeated measure ANOVA in group A (antibiotic) and group B (placebo) showed that mean bacterial colony count differed statistically significant between times (*p*-value 0.000). Post hoc tests using Bonferroni correction revealed that in group A (antibiotic) shows a decrease in bacterial colony count from 4<sup>th</sup> day to 10<sup>th</sup> day (20.20 $\pm$ 4.96 CFU/mL vs 0.70 $\pm$ 1.33 CFU/mL, *p*-value 0.000), which was statistically significant. While there is a non-significant decrease in bacterial colony count from 10<sup>th</sup> day to 30<sup>th</sup> day (0.70 $\pm$ 1.33 CFU/mL vs 0.00 $\pm$ 0.00 CFU/mL, *p*-value 0.397). In group B (placebo), post hoc tests using Bonferroni correction revealed that in this group, decreasing significant decrease in bacterial colony count from 4<sup>th</sup> day to 10<sup>th</sup> day (21.90 $\pm$ 5.93 CFU/mL vs 6.10 $\pm$ 2.07 CFU/mL, *p*-value 0.000), also between 10<sup>th</sup> day to 30<sup>th</sup> day (6.10 $\pm$ 2.07 CFU/mL vs 1.10 $\pm$ 1.10 CFU/mL, *p*-value 0.000) is observed. Therefore, we can conclude that both groups can show decrease in bacterial colony count, even in the placebo group. We can assume the role of innate immunity that can induce cytokins, such as IL-6 or TNF- $\alpha$  that help to fight against the bacteria (Table 3).

A Pearson correlation was used to determine the relationship between bacterial colony count and other variables. Based on the result, there was a strong negative correlation between bacterial colony count and VDR level, which was statistically significant in both groups (group A; *r* = -0.875, *p*-value = 0.000 vs group B; *r* = -0.470, *p*-value = 0.002). IL-6 and TNF- $\alpha$  didn't give a significant correlation with bacterial colony count (Table 4).

Multiple regression was used to predict the correlation

between bacterial colony count from VDR, IL-6 and TNF- $\alpha$  level in the group given antibiotic and placebo. The result of multiple linear regression analysis showed that VDR plays a role in determining in a decrease in bacterial colony count, while other variables were not significant. The higher VDR level, the lower bacterial colony count in both group (group A (antibiotic); B-Coefficients = -1.623, *p*-value = 0.000 vs group B (placebo); B-Coefficients = 0.557, *p*-value = 0.005) (Table 5).

#### 4. DISCUSSION

When *S. Typhi* first enters the body, bacteria will be destroyed by macrophages. Bacteria will be known by various receptors located on the surface of phagocytes [11]. Specific marker molecules for gram-negative bacteria, such as *S. Typhi* are Lipopolisakarida (LPS). LPS will activate TLR-4, which is a receptor that plays a role in observing and destroying *S. Typhi*. Activated TLR-4 will recruit the MyD88 adapter protein. Then, MyD88 recruited IRAK4, IRAK1 and IRAK2. IRAK kinase then phosphorylates and activates the TRAF6 protein, allowing NF- $\kappa$ B to dwell in the cell nucleus and activating transcription and causing induced inflammatory cytokines. Proinflammatory cytokines, such as IL-1 $\beta$  and IL-6, IFN- $\gamma$  and TNF- $\alpha$  are synthesized and systemic inflammation occurs [12].

In this study, we found a significant increase in IL-6 levels on the 4<sup>th</sup> day compared to baseline because the *S. Typhi* bacteria had been injected since day 1. In this study, a group of mice in the given antibiotic and placebo, showed an increase in mean levels of IL-6 at 4<sup>th</sup> day compared to baseline (Table 2). It

can be concluded, after induction by *S. Typhi*, the inflammatory process by cytokines IL-6 has been running. After cytokines are secreted, the helper-1 (Th1) and T helper-2 (Th2) will be activated. Signals from cytokines triggered by the interaction of host cells and bacteria are crucial in the development of the disease. The balance between proinflammatory and anti-inflammatory cytokines will control the prevention of host damage due to excessive inflammation [13].

After giving antibiotics for 5 days on, the average value of IL-6 levels decreased in the antibiotic group (Table 2). Antibiotics are given that can kill bacteria, so it occurs modulating the LPS binding pathway with TLR-4 by the death of bacteria. LPS in bacterial cell walls is reduced so that the TLR-4 protein is not activated. Then by the mechanism of action to modulate the transcription factor pathways, such as NF- $\kappa$ B, so that the production process of IL-6 inflammatory cytokines becomes inhibited and IL-6 levels decrease.

In this study, we calculated the amount of bacterial colonization through the sampling of peritoneal fluid and found a decrease in mean levels of bacterial colony at 10<sup>th</sup> day compare to 4<sup>th</sup> day (Table 3). *S. Typhi* infection given intraperitoneum will stimulate macrophages to activate and move to the source of infection. LPS content in *S. Typhi* cell walls is a signal for macrophages to activate. Activation of macrophages has a high ability to ingest foreign objects through phagocytosis. These cells will destroy all foreign objects, such as bacteria, damaged cells, tumor cells, colloidal objects, and large molecules [14][15]. The increasing number of macrophages to the site of infection originates from the migration of macrophages to the source of stimulation. In addition, the increase in the number of macrophages is caused by the acceleration of the proliferation and differentiation of macrophages. Proliferation and differentiation, as well as increased migration of macrophages to the source of stimulation cause the number of macrophages in the peritoneum to increase. The acceleration of migration is also due to stimulation by IL-6 produced by monocyte cells or macrophages [15]. The IL-6 level on the 30<sup>th</sup> day showed that the value decreased even more along with the number of bacterial colonization (Fig. 1).

TNF- $\alpha$  is a pro-inflammatory cytokine that is produced most by macrophages through several mechanisms [16][17]. Immunologically, TNF- $\alpha$  is produced through macrophage presentation to the Antigen-Presenting Cell (APC) which then APC will order T-helper 1 to proliferate into IL-12 and IFN- $\gamma$  where IFN- $\gamma$  will produce TNF- $\alpha$  as an antimycobacterial. Other studies also explain the same thing, the process of infection can cause plasma enlargement by endothelial activation, so that infected macrophage will become active and release cytokines including tumor necrosis factor alpha (TNF- $\alpha$ ), interleukin 1 (IL-1), IL-6, and platelet activating factor (PAF) [18]. In this study, group A that given antibiotics showed that the mean TNF- $\alpha$  level differed statistically significant between times (*p*-value 0.000) (Table 2). This is because the antibiotics given to the mice in group A can reduce the number of *S. Typhi* thereby suppressing the infection process in the host directly, causing a reduction in cytokine release by macrophage, one of which is TNF- $\alpha$ . Whereas TNF- $\alpha$

levels in the group B that given placebo, although decreased on 10<sup>th</sup> day and 30<sup>th</sup> day, the mean decrease was lower than in the group with antibiotics.

Vitamin D receptors are expressed in large amounts in the tumor tissue and the infected cells. Recent research has shown VDR and enzymes involved in the metabolism of vitamin D have damage to the VDR signaling pathways. VDR expression by immune cells suggests that vitamin D affects the immune system function. More than 30 different body tissues, such as the brain, liver, and pancreas, lymphatic, skin, gonads, and prostate consists of cells including T and B lymphocytes that express the VDR. Vitamin D receptor binds to the 1,25(OH)<sub>2</sub>D<sub>3</sub>, the active form of Vitamin D and the mediated its biological effects [10]. Dendritic cells are the primary targets for the immunomodulatory activity of 1,25(OH)<sub>2</sub>D<sub>3</sub>, by inhibiting the differentiation and maturation of dendritic cells, suppressing the expression regulation of MHC-II, costimulatory molecules (CD40, CD80 and CD86) and decreased production of IL-12. In addition, 1,25(OH)<sub>2</sub>D<sub>3</sub> increases the production of IL-10 and promote apoptosis dendritic cells. Together, the effect of 1,25(OH)<sub>2</sub>D<sub>3</sub> inhibits the activation of T cells that depend on DC [19]. In this study, a group of mice that given antibiotic in group A, showed an increase in mean levels of VDR on 4<sup>th</sup> day compared to 10<sup>th</sup> day (Table 2). VDR activation process then activates the transcription factor genes antimicrobial natural peptide, cathelicidin, and defensins [20], then later will kill the bacteria. This was proved in the result that there was a strong negative correlation between bacterial colony count and VDR level (Table 4). Mice that do not have the VDR impaired production of Th1 promotive factor and IL-18, decreased Th1 cell proliferation and decreased expression of signal transducer and activation of transcription 4 (STAT 4) (a transcription factor Th1 cells). Taken together, these circumstances indicate that VDR function is essential for the development of Th1 cells. In mice that did not have VDR, will have a decreased proliferative response to stimulation of CD3 [19][21][22].

## CONCLUSION

VDR, IL-6, and TNF- $\alpha$  play an important role in killing bacteria. From the results of this study, IL-6 levels are related to the number of bacterial colonies, the lower the IL-6 level, the less the number of bacterial colonies. Similarly, TNF- $\alpha$  level has a positive correlation with the number of bacterial colonies. While VDR level is also related to the number of bacterial colonies, the higher the VDR level, the lower the number of bacterial colonies.

## ABBREVIATION USED

CFU	=	Colony Forming Units
IL-6	=	Interleukin 6
TNF- $\alpha$	=	Tumor Necrosis Factor Alpha
VDR	=	Vitamin D Receptor

## ETHIC APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Faculty of Medicine,



Hasanuddin University Makassar, Indonesia, with registration number 901 / H4.8.4.5.31 / PP36-KOMETIK / 2018.

#### HUMAN AND ANIMAL RIGHTS

No humans were used. All experiments on animals were in accordance with the Health Medical Research Ethics Committee at the Faculty of Medicine, Hasanuddin University (Makassar, Indonesia) as the local guidelines for the care and use of laboratory animals.

#### CONSENT FOR PUBLICATION

Not applicable.

#### 10 AVAILABILITY OF DATA AND MATERIALS

The authors confirm that the data supporting the findings of this study are available within the article.

#### FUNDING

This research was financially supported by the Ministry of Research and Technology, Indonesia.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

#### ACKNOWLEDGEMENTS

Authors would like to thank Rommy Usman, Mus, Wani, and Mark (Molecular Biology and Immunology Laboratory for Infection Diseases, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia) who helped in the implementation of our research activities.

#### REFERENCES

- Crump JA, Sjölund-Karlsson M, Gordon MA, Parry CM. Epidemiology, Clinical Presentation, Laboratory Diagnosis, Antimicrobial Resistance, and Antimicrobial Management of Invasive Salmonella Infections. *Clin Microbiol Rev* 2015; 28(4): 901-37. [http://dx.doi.org/10.1128/CMR.00002-15] [PMID: 26180063]
- Yushi Rohana I. (Departement of Epidemiology, University Airlangga, Surabaya, "The Difference of Knowledge and Primary Preventive for Typhoid Fever Between Parents in Rural and Urban Areas to Children Under-Five Years. *J Berk Epidemiol* 2016; 4(3): 12.
- Nester EW, Jr DA, Roberts CE. *Microbiology: a Human Perspective*. 7th ed. New York, NY, USA: Mc-Graw Hill 2012.
- P. Prof. dr. Syarifuddin Wahid Sp.PA (K) Sp.F, P.dr. Upik AMiskad Sp.PA (K)A.Wijaya *IMUNOLOGI Lebih Mudah Dipahami*. Brillian Internasional 2016.
- Guk KD, Kuprash DV. [Interleukin-11, an IL-6 like cytokine]. *Mol Biol (Mosk)* 2011; 45(1): 44-55. [PMID: 21485496]
- Liang J, Lei Z, Xu X, *et al*. Role of interleukin-6 in differentiating interleukin-11 induced fever and early bacterial infection. *Indian J Pediatr* 2014; 81(9): 871-5. [http://dx.doi.org/10.1007/s12098-014-1361-3] [PMID: 24677115]
- Lee J, Tian Y, Chan ST, Kim JY, Cho C, Ou JH. TNF- $\alpha$  Induced by Hepatitis C Virus via TLR7 and TLR8 in Hepatocytes Supports Interferon Signaling via an Autocrine Mechanism. *PLoS Pathog* 2015; 11(5):e1004937 [http://dx.doi.org/10.1371/journal.ppat.1004937] [PMID: 26023919]
- Zhao X-K, Che P, Cheng ML, *et al*. Tristetraprolin Down-Regulation Contributes to Persistent TNF-Alpha Expression Induced by Cigarette Smoke Extract through a Post-Transcriptional Mechanism. *PLoS One* 2016; 11(12): e0167451-. [http://dx.doi.org/10.1371/journal.pone.0167451] [PMID: 27911957]
- Doll DN, Rellick SL, Barr TL, Ren X, Simpkins JW. Rapid mitochondrial dysfunction mediates TNF-alpha-induced neurotoxicity. *J Neurochem* 2015; 132(4): 443-51. [http://dx.doi.org/10.1111/jnc.13008] [PMID: 25492727]
- Yesil S, Tanyildiz HG, Tekgunduz SA, *et al*. Vitamin D receptor polymorphisms in immune thrombocytopenic purpura. *Pediatr Int (Roma)* 2017; 59(6): 682-5. [http://dx.doi.org/10.1111/ped.13273] [PMID: 28258612]
- van Crevel R, Ottenhoff THM, van der Meer JWM. Innate immunity to Mycobacterium tuberculosis. *Clin Microbiol Rev* 2002; 15(2): 294-309. [http://dx.doi.org/10.1128/CMR.15.2.294-309.2002] [PMID: 11932234]
- Abul Abbas SP, Andrew H. Lichtman, *Cellular and Molecular Immunology*. 8th ed. Philadelphia: W.B. Saunders Company 2014.
- Hurley D, McCusker MP, Fanning S, Martins M. Salmonella-host interactions - modulation of the host innate immune system. *Front Immunol* 2014; 5: 481. [http://dx.doi.org/10.3389/fimmu.2014.00481] [PMID: 25339955]
- Chanana V, Ray P, Rishi DB, Rishi P. Reactive nitrogen intermediates and monokines induce caspase-3 mediated macrophage apoptosis by anaerobically stressed Salmonella typhi. *Clin Exp Immunol* 2007; 150(2): 368-74. [http://dx.doi.org/10.1111/j.1365-2249.2007.03503.x] [PMID: 17888027]
- Besung INK, Astawa NM, Suata K, Suwiti K. HUBUNGAN ANTARA AKTIVASI MAKROFAG DENGAN KADAR INTERLEUKIN-6 DAN ANTIBODI TERHADAP Salmonella typhi PADA MENCIT (Relationship between the Macrophage Activity with Interleukin-6 Levels and Titers of Antibodies against Salmonella typhi). *J Kedokt Hewan* 2016; 10(1): 1-4.
- Coppack SW. Pro-inflammatory cytokines and adipose tissue. *Proc Nutr Soc* 2001; 60(3): 349-56. [http://dx.doi.org/10.1079/PNS20011110] [PMID: 11681809]
- Skoog T, Dichtl W, Boquist S, *et al*. Plasma tumour necrosis factor-alpha and early carotid atherosclerosis in healthy middle-aged men. *Eur Heart J* 2002; 23(5): 376-83. [http://dx.doi.org/10.1053/ehj.2001.2805] [PMID: 11846495]
- Ma K, Zhang H, Baloch Z. Pathogenetic and Therapeutic Applications of Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ) in Major Depressive Disorder: A Systematic Review. *Int J Mol Sci* 2016; 17(5) [http://dx.doi.org/10.3390/ijms17050733] [PMID: 27187381]
- Hayes CE, Nashold FE, Spach KM, Pedersen LB. The immunological functions of the vitamin D endocrine system. *Cell Mol Biol* 2003; 49(2): 277-300. [PMID: 12887108]
- Liu PT, *et al*. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response *Science* (80- ). 2006; 311: pp. (5768)1770-3. [http://dx.doi.org/10.1126/science.1123933]
- Dusso AS, Brown AJ, Slatopolsky E. Vitamin D. *Am J Physiol Renal Physiol* 2005; 289(1): F8-F28. [http://dx.doi.org/10.1152/ajprenal.00336.2004] [PMID: 15951480]
- Dini C, Bianchi A. The potential role of vitamin D for prevention and treatment of tuberculosis and infectious diseases. *Ann Ist Super Sanita* 2012; 48(3): 319-27. [http://dx.doi.org/10.4415/ANN\_12\_03\_13] [PMID: 23007057]

# The Role of IL-6, TNF- $\alpha$ , and VDR in Inhibiting the Growth of Salmonella Typhi: in vivo Study

## ORIGINALITY REPORT

**25%**  
SIMILARITY INDEX

**23%**  
INTERNET SOURCES

**12%**  
PUBLICATIONS

**7%**  
STUDENT PAPERS

## PRIMARY SOURCES

<b>1</b>	<a href="http://pagepressjournals.org">pagepressjournals.org</a> Internet Source	<b>13%</b>
<b>2</b>	Juan Liang, Zhiyu Lei, Xiaojun Xu, Ning Zhao et al. "Role of Interleukin-6 in Differentiating Interleukin-11 Induced Fever and Early Bacterial Infection", The Indian Journal of Pediatrics, 2014 Publication	<b>1%</b>
<b>3</b>	<a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	<b>1%</b>
<b>4</b>	<a href="http://akjournals.com">akjournals.com</a> Internet Source	<b>1%</b>
<b>5</b>	Submitted to Tallahassee Community College Student Paper	<b>1%</b>
<b>6</b>	<a href="http://journal.umpalangkaraya.ac.id">journal.umpalangkaraya.ac.id</a> Internet Source	<b>1%</b>
<b>7</b>	Eifan Boyke Pattiasina, Pieter Kakisina, Ferymon Mahulette. "Decreasing SGPT level and macrophage activity through CD68	<b>1%</b>

expression in the Balb/c mice (*Mus musculus*)  
liver infected with *Salmonella typhi* after  
treating with atung seeds (*Parinarium*  
*glaberimmun Hassk*)", *BIOEDUPAT: Pattimura*  
*Journal of Biology and Learning*, 2021

Publication

---

8	<a href="http://m.scirp.org">m.scirp.org</a> Internet Source	1 %
9	<a href="http://biofarmaka.ipb.ac.id">biofarmaka.ipb.ac.id</a> Internet Source	1 %
10	<a href="http://opennursingjournal.com">opennursingjournal.com</a> Internet Source	1 %
11	<a href="http://repository.unhas.ac.id">repository.unhas.ac.id</a> Internet Source	1 %
12	M. Wielgos. "P07.01: Giant endometrial cyst of the uterine wall-a case report", <i>Ultrasound in Obstetrics and Gynecology</i> , 09/08/2005 Publication	1 %
13	Submitted to Universitas Indonesia Student Paper	1 %
14	<a href="http://www.jimmunol.org">www.jimmunol.org</a> Internet Source	1 %
15	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1 %

---

---

Exclude quotes      On

Exclude matches      < 1%

Exclude bibliography      On