



**Submit**





#### Article

# Artisanal and Small-Scale Gold Mining Activities and Mercury Exposure in Gorontalo Utara Regency, Indonesia

Yayu Indriati Arifin<sup>1,2,\*</sup>, Masayuki Sakakibara<sup>2,3</sup>, Sayaka Takakura<sup>4</sup>, Mohamad Jahja<sup>2,5</sup>, Fitryane Lihawa<sup>6</sup> and Koichiro Sera<sup>7</sup>

- <sup>1</sup> Department of Geology, State University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia
- <sup>2</sup> The Research Institute for Humanity and Nature (RIHN), 457-4 Motoyama, Kamigamo, Kita-ku, Kyoto, 603-8047, Japan
- <sup>3</sup> Graduate School of Sciences and Engineering, Ehime University, 2-5 Bunkyo-cho, Matsuyama, 790-8577, Japan
- <sup>4</sup> Ehime University, 3 Bunkyo-cho, Matsuyama, 790-8577, Japan
- <sup>5</sup> Department of Physics, State University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia
- <sup>6</sup> Department of Educational Geography, University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia
- <sup>7</sup> Cyclotron Research Center, Iwate Medical University, 348-58 Tomegamori, Takizawa, Iwate 020-0173, Japan

\* Correspondence: yayu\_arifin@ung.ac.id; Tel.: +81-089-927-9649; Fax: +81-089-927-9640

Received: date; Accepted: date; Published: date

**Abstract:** Small-scale gold mining activities and their mercury emissions in the Utara Regency of the Province, Sulawesi Island, Indonesia, have been studied. The mining occurs in many areas of this

**Commented [01]:** Instructions for Authors (IfA): <https://www.tandfonline.com/action/authorSubmission?show=instructions&journalCode=gtec20>

Please include a word count for your paper.

**Commented [L2]:** Word Its is deketed; s in exposure also deleted number of word ( with spaces) in title is 107

**Commented [03]:** IfA: Write in an objective, neutral manner in the third person, not in the 1st person 'we' or 'I'.

**Commented [L4]:** (We studied and) is deleted

**Commented [L5]:** Small s is replaced with Captial S

**Commented [L6]:** (, have been studied) is inserted

region, but especially in the three villages Hulawa, Ilangata and Ilangata Barat. Mining activities in the Hulawa village are the oldest (19th century) in the regency, whereas mining in the Ilangata village began just fifteen years ago. The activity level changes rapidly from time to time depending on the fluctuations of the gold market. Women and children are involved in the gold mining process. Female workers crush ore, process sediment wastes, and pan for gold in rivers. Children assist their mothers in these activities and play around the mining sites. Gold mining in the Utara Regency is estimated to produce about 290 kg of gold and emits approximately 860 kg of mercury into the environment annually. The impact of such large mercury release into the environment have caused environmental degradation and contamination of fish as food for humans. Collaborative works of stakeholders are urgently needed to find solutions to the problems related to small-scale gold mining such as marginalization and environmental pollution.

**Keywords:** artisanal and small-scale gold mining; mercury exposure; human health problems; Gorontalo Utara; Indonesia

## 1. Introduction

Indonesia is the world's second-largest source of mercury emissions from artisanal and small-scale gold mining (ASGM) after China (Veiga, Maxson, and Hylander 2006; Pacyna *et al.* 2010; Ismawati 2014). The top ten global ASGM mercury emitters in 2000 and 2005 are shown in Figure 1 (Telmer and Veiga 2009; Cordy *et al.* 2011; Pacyna *et al.* 2010; Veiga, Maxson, and Hylander 2006; Ismawati 2014). Annual mercury emissions from ASGM in Indonesia in 2000, 2005, and 2013 were approximately 125, 150, and 175 tons, respectively. In a five-year period (from 2000 to 2005), Indonesia's ASGM mercury emissions were increased by approximately 20 %.

According to UNEP (2008), there are 115 ASGM industries distributed evenly throughout Indonesia that collectively produce 60 tons of gold per annum. The number of ASGM sites in Indonesia is estimated to be ca. 900 involving about 250,000 miners (Purwana 2013). The largest gold mining site in Indonesia is located near Puncak Jaya in Papua. Tailings from this site are dumped in Timika in eastern Indonesia (Nakagawa 2008; Vogt 2012). Here ASGM activities can be found in several places, e.g.: Jayapura, Nabire, Paniai and Manokwari. ASGM activities also occur in Maluku Province, Gunung Botak and Gogorea on Buru Island, Maluku Province (Male *et al.* 2013; Idrus *et al.* 2014).

Several ASGM locations have also been reported in Sulawesi, including Poboya in Palu (Basir-Cyio *et al.* 2012). According to Iswamaty, Petrlik, and DiGangi (2013), ASGM miners' activities in Poboya use 100 tons of mercury annually (Nakazawa *et al.* 2016). Basyir-Cyio *et al.* (2012) estimate that about 34 tons of mercury are used daily and produce 22.4–45 kg gold daily. High concentrations above 1000 ng m<sup>-3</sup> the limit of air-quality guideline for annual mercury exposure by the World Health Organization of mercury vapor from ASGM were detected in Palu City, and almost all residents of the city are exposed (Nakazawa *et al.* 2016). Several ASGM has been reported in Mamuju of Sulawesi Barat Province, TanaToraja and Luwu of Sulawesi Selatan Province and the Bombana of Sulawesi Tenggara Province. The Bombana ASGM has been reported to have negative impact on social, economic and cultural status of the people around the ASGM (Rianse *et al.* 2015).

There are four other ASGM locations in Sulawesi Utara Province (Basir-Cyio *et al.* 2012), at Talawaan (Filho and Villas-bôas 2004; Bose-O'Reilly *et al.* 2010), Buyat (Limbong *et al.* 2005; Kambey, Farrell, and Bendell-Young 2001), Tobongon and Tanoyan in the Bolaang Mongondow Regency. Mercury emissions from Talawaan alone are 70 t per annum (Lasut *et al.* 2010), which is seven times the value reported earlier by Telmer and Veiga (2009).

Commented [L7]: Small h is replaced by Capital H

Commented [L8]: Small i is replaced with Capital I

Commented [L9]: Small b is replace with Capital B

Commented [L10]: Small h is replace with Capital H

Commented [L11]: A space is deleted

Commented [L12]: Furthermore, is deleted

Commented [L13]: Small t is replaced with Capital T

Commented [L14]: es is inserted

Commented [L15]: d is deleted

Commented [L16]: and human population is deleted

Commented [L17]: fish as food for is inserted

Commented [L18]: s and fish is deleted

Commented [L19]: to the is inserted

Commented [L20]: Y is inserted

Commented [L21]: Comma is inserted

Commented [L22]: Comma is inserted

Commented [L23]: and miners is deleted

Commented [L24]: involving about is inserted

Commented [L25]: and is deleted

Commented [L26]: a space is deleted

Commented [L27]: miners is inserted

Commented [L28]: people, respectively is deleted

Commented [L29]: comma is deleted

Commented [L30]: comma is inserted

Commented [L31]: Moreover is deleted

Commented [L32]: Small s is replace with Capital S

Commented [L33]: Small c is replaced with capital C

Commented [L34]: Yuyun Ismawaty, Petrlik and DiC (... [1]

Commented [L35]: Ehile is deleted

Commented [L36]: Comma is deleted

Commented [L37]: (2012) is inserted

Commented [L38]: s is deleted

Commented [L39]: they used is deleted

Commented [040]: Daily? Monthly?... daily is inserted

Commented [L41]: are used is inserted

Commented [L42]: d is deleted

Commented [L43]: a space is inserted

Commented [L44]: a space is inserted

Commented [L45]: Basyir-cyio et al 2012 is deleted

Commented [L46]: A space is inserted

Commented [L47]: A space is inserted

Commented [048]: What is the limit? Here the limit n (... [2]

Commented [L49]: The limit is mentioned clearly

Commented [L50]: Elsewhere is deleted

Commented [L51]: Status is inserted

Commented [L52]: The is inserted

Commented [L53]: The is inserted

Commented [L54]: Small c is replaced with C

Commented [L55]: of is replaced with the

At least there is one ASGM site in each regency of the Gorontalo Province: (1) Pohuwato Regency: Gunung Pani and Bulontio; (2) Boalemo Regency: Bilato; (3) Gorontalo Regency: Bumela (Lihawa and Mahmud 2019) (4) Bonebolango Regency: Tulabolo and Mopuya; and (5) Gorontalo Utara Regency: Hulawa and Ilangata villages.

Gold deposits in the Gorontalo Utara Regency have existed in Wubudu breccias within Boliyohuto diorite of the middle Miocene age (Carlile, Digdowirogo, and Darius 1990). Gold in hydrothermal breccias and quartz veinlets has been reported in Sumalata and Boliyohuto subdistricts (Carlile, Digdowirogo, and Darius 1990). Gold mining in the Hulawa village is considered to date back to Indonesia Dutch era (Henley 2005).

This study describes mercury pollution from ASGM sites in the Gorontalo Utara Regency where gold mining activity is intense, but information is scarce. The report describes how miners work and what the mercury emission situation is, as a basis for further discourse on health risks in and around these mining sites.

Commented [L56]: comma is deleted

## 2. Materials and Methods

### 2.1. Study Area

The Hulawa (meaning 'gold' in the Gorontalo language) mining area (Figure 2) is also known as the Buladu mining area in Sumalata. This area lies within the Buladu River watershed, between the Pasolo and Padengo villages, approximately 109 km northeast of Gorontalo City, the capital of Gorontalo Province. From Gorontalo City, the area can be reached by car in approximately four hours. Buladu has a typical equatorial climate with two seasons (rainy: November–April; dry: May–October), with the highest rainfall occurring in December (annual average 1,210 mm). The temperature is 27 °C on average. The Buladu River watershed drains from the mountain peaks with the highest point being 2,000 m, on the northern coast of the Sumalata district. The distance from the mountain peaks to the Sulawesi Sea is approximately 10 km. Several rivers run through this watershed, including the Buladu and Wubudu rivers.

The Ilangata and Ilangata Barat mining areas (Figure 2) comprise the Anggrek mining area. The area lies within the Anggrek River watershed between the Diata, Lunggulo, and Lantolo villages, which are approximately 68 km northwest of Gorontalo City. It is three hours by car from Gorontalo City. The Anggrek River watershed drains from the peaks of several hills with elevations of 300 m at most. The distance from the peaks to the sea is approximately 5 km. Number of rivers, including the Anggrek and Ilangata rivers, flow through the Anggrek watershed.

Residents of Hulawa, Ilangata and Ilangata Barat are exposed directly and indirectly to mercury from mercury vapor of ASGM process and sea fish consumption, while residents of other subdistricts (Kwandang, Monano and Tolinggula) are treated as control group since they might only be exposed to mercury through sea fish consumption.

### 2.2. Sample and data collection

The ASGM locations within Gorontalo Utara regency have been visited many times over the course of one month each year, from 2012 to 2014. Data on mercury use during ASGM and the activities of miners and their families were obtained via direct interviews with miners, mine owners, and local government officials. These interviews were conducted on-site at the mining locations in the Gorontalo Utara Regency (Figure 2). The simple questionnaire consisted of number of workers, number of gold trommels, amount of ore filled into trommels, and amount of mercury consumed and gold produced in a single mill process. This was done in Hulawa, Ilangata and Ilangata Barat ASGM sites (n=23): 6 ASGM operators, 2 panners, 3 villagers in Hulawa, 6 ASGM operators, 4 villagers and 2 gold shop workers in Ilangata and Ilangata barat, 2 officers in Kwandang). The interviewed were key-

Commented [L57]: ( is deleted

Commented [L58]: Were is inserted

actors, including mine site managers, head of villages, and elderly people who were working as ASGM miners 20 to 40 years ago.

Commented [L59]: and is inserted

Data obtained from different sources were compared with each other via triangulation procedure, for example, number of miners from the owner, from direct investigation and from local officers were confronted other to keep accountable. Any information obtained from participants was recorded and is used in triangulation process. For instance, we recorded the following information while communicating with villagers who have a motorcycle repair kiosk in Ilangata, his friend said that: "he started that business after he earned about 6000 US-\$ for a week from ASGM one year ago". Once we were in Hulawa village, a man who got training on the Minamata disease problem informed us that in nearby villages live three people who were suspected to have tremor disturbance related to their former activities as ASGM miners in Hulawa. The first person showed tremor during our conversation and he has a problem on oral communication already. The second one, also showing tremor with a lesser degree, said that: "We were working as ASGM miners, dealing with mercury without safety equipment during smelting; sometimes they put gold amalgam in their mouths to sell later on". They also consumed fish from contaminated rivers, the estuary, and the sea.

Commented [L60]: were confronted is inserted

Commented [061]: ??

Commented [L62]: other to keep accountable is deleted

Commented [L63]: We is deleted

Commented [L64]: s is deleted

Information on the economy and social life in ASGM were also obtained from interviews with miners, managers of ASGM businesses, and villagers. Compared to large-scale mining, ASGM activities have more positive economic impacts on local communities, since the capital is shared within local investors, miners, and villagers (Langston *et al.* 2015). Economy and social life information, in this case, will enrich our understanding on ASGM activities, its impacts on the environment and human health, and ongoing struggles against social marginalization in Indonesia's gold mining sector (Spiegel and Veiga 2010).

Commented [L65]: comma is inserted

Commented [L66]: s is inserted

### 2.3. Analytical Procedure

The health effects on ASGM miners and villagers were examined by collecting scalp hair strands from 115 donors from five subdistricts. The hair strand from the occipital region was cut close to the skull, collected, put in a clear plastic bag, and labeled accordingly. Mercury content in hair was determined using proton induced X-ray emission (PIXE) at Iwate Medical University, Japan, as reported elsewhere (Sera *et al.* 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera 2017). The procedure is known as a standard-free method for hair without powdering and acid treatment as reported in previous studies (Sera, Futatsugawa, and Matsuda 1999; Sera, Futatsugawa, and Murao 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera 2017). The widely accepted exposure limit values for mercury concentration in hair ( $\mu\text{g/g}$ ) as suggested by the German Human Biomonitoring Commission (HBM) as the following categories are adopted: concentration ( $\mu\text{g/g}$ ) below 1.0 are considered normal, 1.0 - 5.0 are an alert level, and above 5.0 are indicative of a serious risk of health effects.

Commented [L67]: s is deleted

Commented [068]: IfA: Abbreviations are to be introduced upon first mentioning of the fully spelled term (initial letters of each word in lower-case, not capital).

Commented [L69]: We follow the IfA all capitals replaced with lower-case letters

Commented [L70]: Comma is inserted

Commented [L71]: The method is deleted

Commented [L72]: A space is inserted

Commented [L73]: two commas are replaced with two ; and et al. is replaced with , Sakakibara and Sera

Commented [L74]: full sentence is deleted

Commented [075]: IfA: Write in an objective, neutral manner in the third person, not in the 1st person 'we' or 'I'.

Commented [L76]: The is replacing We adopt

Commented [L77]: are adopted is inserted

Commented [L78]: comma is inserted

### 2.4. Neurological Assessment

The subjects with elevated hair mercury concentration were examined clinically according to a questionnaire especially for subjective symptoms, and the use of mercury-containing soap (for female only), and a neurological test. Neurological examination was performed by a medical doctor on participants, using a UNIDO protocol which consists of ten symptoms: (1) signs of bluish discoloration of gums; (2) rigidity and ataxia (walking and standing); (3) alternating movements or dysidiadochokinesia; (4) irregular eye movements or nystagmus; (5) field of vision; (6) knee jerk reflex and biceps reflex; (7) Babinski reflex and labial reflex; (8) salivation and dysarthria; (9) sensory examination and (10) tremor: tongue, eyelids, finger to nose, pouring, posture holding, and the Romberg test. Score 1 was given for positively-observed symptoms, and 0 for negatively-observed symptoms. The sum of score for every participant is calculated and analyzed

Commented [L79]: We is deleted and small s is replaced with capital S following the IfA

Commented [L80]: D is deleted

Commented [L81]: Was given is inserted

Commented [L82]: s is deleted

### 2.5. Description of sites and processes in of ASGM

Ore is manually excavated from nearby hills or mountains (Figure 3a) using vertical holes in the ground (Figure 3b). Traditional tools such as broad hoes, bars and simple pulleys are used for the ore excavation (Figure 3c). The ore is manually crushed (Figure 3d) using hammers or other percussion tools (Figure 3e). Some processing plants use homemade mechanical crushers. The ore is then packed into sacks and manually transported to the processing plant (Figure 3f).

At the processing plant, crushed ore (30–40 kg) is fed into a trommel (known as “Tromol” in the local language (James 1994), a steel mill grinder where gold is extracted from ore (Male et al. 2013) (Figure 4a) with water (Figure 4b) and several decimeter-sized rocks (Figure 4c) and milled for 3–4 hours until the material is fine enough to release the gold. Subsequently, approximately 500 g of mercury are added to the trommel (Figure 4d), which then rotates for 30 minutes until the amalgamation has completed. This amalgamation process is the major gold production method for ASGM miners in Gorontalo Province.

After 30 minutes, the material is discharged (Figure 4e), and the tailings are manually separated using bare hands (Figure 4f). The tailings are washed and rinsed several times (Figures 4g – 4j) to separate the gold from the unamalgamated mercury. The mixture is filtered using a fine cloth to separate the mercury amalgam (Figure 4k). The gold amalgam is recovered from this process as shown in Figure 4l.

After milling the gold, the amalgam is recovered, and the waste sediment is kept (Figure 4m) for a second recovery process. Then, mercury-contaminated water is released into the environment (Figure 4n). The amalgams are then smelted to obtain gold bullion. Gold smelters with simple tools smelt the gold outdoors (Figure 4o) or inside houses and gold shops. Gold bullion is recovered from the smelting process (Figure 4p).

The secondary process begins when miners, usually women, and children, wash sediment wastes from the primary milling process in running water. The milling process is then repeated, as shown in Figure 4a through Figure 4l, with the difference being that the trommel is generally rotated by a wheel powered with river water (Figure 4n).

Tertiary processing involves bagging the waste from the secondary processing and sending it to another processing plant to undergo cyanidation. There is no cyanidation plant in the Gorontalo Utara Regency. However, there is a cyanidation plant in the other Regency of Gorontalo Province. Numerous miners pan in the Wubudu River estuary (Figure 4q) and (Figure 4r) where the mercury enters the environment (air, soil, and water) directly.

Smelting activities occur in gold shops in the Ilangata Barat village. The owners of these gold shops determine the gold quality (amount of gold in “dore”) via these processes. The amount of gold in a dore varies from 40 – 80% of the weight. A dore with 80 % quality can be purchased for approximately US \$30,-/gram. Mining wastes have solid and liquid forms that contaminate river waters and soils. Gold bullion is extracted after smelting the amalgam, which emits mercury vapor.

### **3. Results**

#### *3.1. Mercury Emissions*

Because ASGM activities are unregistered, no exact data were available on the number of workers; therefore, it is difficult to estimate the amount of gold produced, and mercury used, for ASGM activities throughout Indonesia and its local provinces or regencies. This difficulty in estimating amounts of gold produced and mercury used by ASGM throughout the world is described elsewhere (Seccatore et al. 2014). We used direct observation and interviews to get information and compared it with data reported by several researchers. Data on mercury imported to Indonesia will be used as the upper limit of mercury lost to ASGM. Socio-economic factors which should be

considered when dealing with ASGM (Sulaiman et al. 2007; Spiegel 2012; Langston et al. 2015; Spiegel and Veiga 2010), will be elaborated further to discuss any possible scenario or solutions on sustainable development in local ASGM site areas.

Data on the number of mine workers, population and mercury used during the milling process in each village are listed in Table 1. These figures were estimated based on each milling process requiring 0.5 kg mercury for one trommel mill to extract gold from ca. 30-40 kg of crushed ore. Assuming all trommel mills are operating during the milling process, a maximum of 275 kg of mercury is needed (these data and the amount of mercury emitted per gram of gold produced allows us to estimate the mercury emitted into the environment). Each primary process produced ca. 3 g of gold per trommel mill. The secondary processing of waste produces ca. 0.5 g of gold per trommel mill. We also assume the miners only work five days a week and sell the gold on Saturday. The ratio of mercury emissions to gold produced is ca. 2:1 (Darmutji 2003). The amount of gold produced by the process is ca. 1134 g. Thus, the mercury emissions are ca. 2.2 kg per milling process. The total annual mercury emission in the Gorontalo Utara Regency is approximately 572 kg. If we used 1-3 grams of mercury lost per gram of gold produced, the mercury emission ranges from 286 – 858 kg (Veiga et al. 2009).

### 3.2. Mercury Concentration in Hair Samples According to Geographic Location

The distributions and range of mercury levels in 115 hair samples collected from the five subdistricts are shown in Figure 5. Mercury concentrations in all hair donors are more than 1 µg/g, which indicates the concentrations are already within the alert level range according to HBM (Schulz et al. 2007). The number of subjects with high mercury levels over 10 µg/g was 10 (40 %), 7(30 %), and 4 (8.5 %) in Anggrek, Sumalata, and the control group (Kwandang, Monano and Tolinggula), respectively.

According to the Kolmogorov-Smirnov test, the distribution of mercury in the hair data from the Gorontalo Utara Regency was not normal; instead, it had a log normal distribution (Figure 6). The geometric mean is more suitable for log normal distribution data. Mean mercury level (+/- standard deviation) of Gorontalo Utara regency is 6.7 (2.0) µg/g and is higher than minimum limit for danger level (5.0 µg/g) according to HBM. Only residents of Tolinggula has average hair mercury level less than 5.0 µg/g. This may due to their residence away from the seaside; their common diet along with rice is chicken. Even though one male and two females have mercury higher than 6.0 µg/g, the Tolinggula group could be considered as a control group in Gorontalo Utara regency.

The data were plotted using a box-and-whisker plot to make a clear view of data distribution in each residence and among Gorontalo Utara regency data. Each Box is filled with the horizontal line representing the position of the median. The position of the median and mean relative to HBM was used to put the population into three categories according to HBM. Box plot of mercury distribution according to geographical location is shown in Figure 5, the results of Sumalata and Anggrek are positioned higher than others, which means that their hair average mercury is elevated in the regency. Other subdistricts (Tolinggula and Kwandang) have no known ASGM activities, but the population lives close to seashore (their common diet is fish from Sulawesi sea, same as the population of Anggrek and Sumalata).

### 3.3. Mercury concentration for male and female

The lognormal distribution of hair mercury levels in males and females is shown in Figure 6. Five females had mercury levels greater than 25 µg/g, and none of them worked as ASGM miners. Those five females may have been exposed to mercury from another source. The elevated hair mercury levels that were above average (7.1 µg/g) and even the highest (17.9 µg/g) mercury level were found among the ASGM miners. The average hair mercury levels for all, male and female inhabitants in the Monano subdistrict are 5.7, 5.0 and 6.2 µg/g, respectively. These levels show that

Commented [L83]: a space is inserted

Commented [L84]: .4 is deleted



there are no significant differences between the mean hair mercury of males and females in that subdistrict. While the average of all mercury hair content for females is 8.1 µg/g (more than 30 percent higher than males (6.0 µg/g), such conditions were also found for the subgroups of Kwandang and Sumalata. The condition in Anggrek is even higher (the average mercury for females is three times than the males). The large discrepancy of mercury levels between female and male inhabitants suggests that female inhabitants may be receiving mercury from another source (e.g., whitening cream, soap, etc).

The clinical symptoms were observed in most of the females with mercury higher than 50 mg/g (the minimum level at which Minamata disease symptoms are seen (Harada 1995)) are listed in Table 2, indicative of inorganic-mercury poisoning only. All the subjects have habitual use of whitening cream that are common in Gorontalo Utara regency. The mercury concentration on the whitening creams are shown in Table 3. It seems that elevated mercury concentration among female inhabitants of Hulawa and Ilangata may be caused by direct mercury contact from habitual use of whitening creams. Harada et al. (2001) found the similar cases in Kenya which were caused by female workers using elevated hair mercury levels while using soaps with high mercury concentrations.

#### 4. Discussion

##### 4.1. ASGM operation system and Mercury losses

Figure 7 shows the ASGM operation system which is used in Gorontalo Utara Regency; the system used in North Gorontalo is similar to those reported elsewhere (Veiga *et al.* 2009). Gold recovery systems are based on whole ore amalgamation using trommel, gravitation and panning methods were found in Wobudu river estuary and Anggrek river banks. There are two possible routes for mercury contamination from ASGM into the environment and humans. During the milling process, liquid mercury could contaminate miners and the environment. Working with mining wastes (tailings), and direct exposure to liquid mercury during panning, are also contamination routes. Mercury vapor could contaminate the smelters and environment during smelting.

For comparison, we also estimated the total annual mercury emissions in the Gorontalo Utara Regency from ASGM using data on Indonesian mercury emissions (Pacyna *et al.* 2010) and the number of known ASGM sites (Kocman *et al.* 2013). The rate for Indonesian mercury emissions from ASGM may be close to 200 t annually. The average emission from each mining site is approximately 200 kg/year based on the 900 known ASGM sites (Kocman *et al.* 2013). The three mining sites in the Gorontalo Utara Regency emit approximately 600 kg/year. Ismawaty (2014) estimated the ASGM production ranged from 65-130 t in 2012 (Ismawaty 2014). Thus, approximately 170 – 260 t of mercury was lost to the environment based on the multiplication factor from Darmutji (2003). Using the 1:1 to 3:1 ratio proposed by Ismawaty (2014), the amount of mercury lost was 65 – 390 t. These estimates agree with the total mercury import data for Indonesia in 2012, which was approximately 368 t.

##### 4.2. Health Issues

The involvement of women and children workers is unique to the Hulawa mining site. These women and children come from the Buladu area. They work with waste stored in wooden boxes, and sluice and pan in the Wobudu River estuary. Women and children typically work on mercury-bearing mining wastes during the secondary processing stage (Figures. 8a and 8b). Their bare hands and feet are directly exposed to the mining waste, which allows mercury to enter their body via dermal absorption.

Health problems in miners and inhabitants were documented by medical doctors using a standard questionnaire and neurological test tools (Figures. 9a and 9b). Ten neurological disorder indicators/symptoms were observed. Bluish gums and tremors are major symptoms observed in the miners and inhabitants of the Gorontalo Utara Regency. Several explanations could be addressed based on the evidence; first, they have experienced prolonged contact with mercury, and second, the mercury



has entered their body via the food chain (Basir-Cyio et al. 2012; Nakazawa et al. 2016; Bose-O'Reilly et al. 2010). Both can happen simultaneously for the miners and inhabitants of these mining sites. Further investigations are needed to explain the evidence and determine the most probable mercury contamination routes in the Gorontalo Utara Regency.

Without pollution controls and environmental cleanup, further mercury contamination of the environment (bottom sediments, water, and fish) and human population around ASGM sites within the Gorontalo Utara Regency is inevitable (Arifin, Sakakibara, and Sera 2015). Veiga and Hinton (Seccatore *et al.* 2014) reported that gold bullion contains two to five percent mercury just after smelting, and such gold bullion is refined at gold shops in towns within the Gorontalo Utara Regency with few emission controls. Mercury pollution from gold shops, like those observed in Colombian cities, is of great concern (Cordy *et al.* 2013).

#### 4.3. Socio-economic aspects of ASGM system in Gorontalo Utara Regency

The intensification of ASGM activities at the historical gold mining site (hulawa village) and expansion to the new gold mining site (ilangata and ilangata barat villages) has been seen as advantage of ASGM over large scale gold mining (LSGM)(Adler Miserendino *et al.* 2013). The ability of ASGM activities to cut down production cost through their reliance on flexible informal labor. The ASGM activities are driven by local investors which are intimately entangled with socio-political structures and land tenure systems (Verbrugge and Geenen 2019).

The hierarchy of people working in ASGM system in Gorontalo Utara regency can be explained using the network chart introduced by Langston et al. (2015), with slight modification mainly related to the number of ASGM miners. In general, a small number of miners (up to 3 to 4 people) work at the excavating site or processing site; these miners may be family members or friends. Hence supervisor, landowner position, and even investor itself can be the same person.

Local miners are formerly farmers or fisherman and live nearby in their houses, while migrants usually live inside the processing site camps, which are very simple structures and lack sanitation facilities. The basic needs (foods, drinks, cigarettes) were available in kiosk/gold shops which may belong to cooperation or investor/landowner/supervisor.

Positive economic impacts of ASGM activities to miners and inhabitants were not easily seen in Gorontalo Utara regency since there is no significant difference in economic activities in ASGM sites and non ASGM sites. In a stakeholder meeting, ASGM miners, local government, and investor seem to have opposing viewpoints on issues such as sustainable development, environmental protection, and negative impacts of ASGM on health.

Local governments and inhabitants would like to embrace long-term economic solution such as farming, fishery, and tourism, which could eventually replace ASGM. There is ongoing research on finding the ultimate solutions which fit to Gorontalo Utara Regency cases. The multidisciplinary research approach is being taken to help young local students who were living in the ASGM area to find solutions to their local problems. A transdisciplinary approach could facilitate the stakeholders in ASGM area together with the researcher to develop the ideas on sustainable development (Sippl and Selin 2012).

## 5. Conclusions

ASGM activities in the Gorontalo Utara Regency are dangerous to the environment, miners, and the local population. The amount of mercury used per mining cycle is approximately 275 kg, and approximately 2.2 kg of mercury is lost per cycle. Annually, ASGM in the Gorontalo Utara Regency produces approximately 286 kg of gold and emits approximately 572 kg of mercury. This estimate is probably conservative given the limited mining data available. The impact of such a large mercury

Commented [L85]: With is replaced by to

Commented [L86]: Comma is inserted

Commented [L87]: S is inserted

Commented [L88]: Comma is inserted

Commented [L89]: Are is replaced with is

Commented [L90]: es is deleted

loss into the environment and human population has been to degrade the environmental conditions and contaminating humans and fish as demonstrated by Arifin et al. (2015).

Gorontalo Utara Regency is an essential agricultural (rice) and horticultural (dairy) production area in the Gorontalo Province and Indonesia. The ASGM driven economy did not bring sustainable development in the vicinity of ASGM sites and Gorontalo Utara regency as a whole. Solutions to the problem are being developed using ongoing multi and trans-disciplinary research approach through the collaboration of Indonesian and Japanese researchers.

**Author Contributions:** All authors contributed to the work presented in the manuscript.

**Acknowledgments:** The authors wish to thank the regent of Gorontalo Utara and its officers, and the owners of mining plants for allowing us to conduct interviews and take photographs. This research was supported by The Research Institute for Humanity and Nature (RIHN: a constituent member of NIHU), Project No.14200102.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Adler Miserendino, Rebecca, Bridget A. Bergquist, Sara E. Adler, Jean Remy Davée Guimarães, Peter S J Lees, Wilmer Niquen, P. Colon Velasquez-López, and Marcello M. Veiga. 2013. "Challenges to Measuring, Monitoring, and Addressing the Cumulative Impacts of Artisanal and Small-Scale Gold Mining in Ecuador." *Resources Policy* 38 (4). <https://doi.org/10.1016/j.resourpol.2013.03.007>.
- Arifin, Yayu, Masayuki Sakakibara, and Koichiro Sera. 2015. "Impacts of Artisanal and Small-Scale Gold Mining (ASGM) on Environment and Human Health of Gorontalo Utara Regency, Gorontalo Province, Indonesia." *Geosciences* 5 (2): 160–76. <https://doi.org/10.3390/geosciences5020160>.
- Basir-Cyio, Muhammad, Uswah Hasanah, Isrun Nur, and Yuka Serikawa. 2012. "Gold Mining Activities and Its Impacts on Land Degradation in Central Sulawesi Indonesia." *Journal of Ecotechnology Research* 16 (3–4): 79–83. <https://doi.org/https://doi.org/10.11190/jer.16.79>.
- Basri, Masayuki Sakakibara, and Koichiro Sera. 2017. "Current Mercury Exposure from Artisanal and Small-Scale Gold Mining in Bombana, Southeast Sulawesi, Indonesia—Future Significant Health Risks." *Toxics* 5 (1): 7. <https://doi.org/10.3390/toxics5010007>.
- Bose-O'Reilly, Stephan, Gustav Drasch, Christian Beinhoff, Saulo Rodrigues-Filho, Gabriele Roider, Beate Lettmeier, Alexandra Maydl, Stefan Maydl, and Uwe Siebert. 2010. "Health Assessment of Artisanal Gold Miners in Indonesia." *Science of the Total Environment* 408 (4): 713–25. <https://doi.org/10.1016/j.scitotenv.2009.10.070>.
- Carlile, J C, S Digidowirogo, and K Darius. 1990. "Geological Setting, Characteristics and Regional Exploration for Gold in the Volcanic Arcs of North Sulawesi, Indonesia." *Journal of Geochemical Exploration* 35 (1): 105–40. [https://doi.org/http://dx.doi.org/10.1016/0375-6742\(90\)90037-B](https://doi.org/http://dx.doi.org/10.1016/0375-6742(90)90037-B).
- Cordy, Paul, Marcello Veiga, Ben Crawford, Oseas Garcia, Victor Gonzalez, Daniel Moraga, Monika Roeser, and Dennis Wip. 2013. "Characterization, Mapping, and Mitigation of Mercury Vapour Emissions from Artisanal Mining Gold Shops." *Environmental Research* 125: 82–91. <https://doi.org/10.1016/j.envres.2012.10.015>.
- Cordy, Paul, Marcello M. Veiga, Ibrahim Salih, Sari Al-Saadi, Stephanie Console, Oseas Garcia, Luis Alberto Mesa, Patricio C. Velásquez-López, and Monika Roeser. 2011. "Mercury Contamination from Artisanal Gold Mining in Antioquia, Colombia: The World's Highest per Capita Mercury Pollution." *Science of The Total Environment* 410–411: 154–60. <https://doi.org/10.1016/j.scitotenv.2011.09.006>.

- Darmutji, Selinawati T. 2003. "Report, UNIDO Project No EG/GLO/01/G34, Overview of Project Site in Indonesia Talawaan and Galangan Areas."
- Filho, Saulo Rodrigues, and Roberto Villas-bôas. 2004. "Biogeochemical Assessment of Mercury Pollution in the São Chico Gold Mining Site - Brazilian Amazon Rio De Janeiro Novembro / 2004." In *Environmental Geochemistry in Tropical Counties*, 235.
- Harada, Masazumi. 1995. "Minamata Disease: Methylmercury Poisoning in Japan Caused by Environmental Pollution." *Critical Reviews in Toxicology* 25 (1): 1–24. <https://doi.org/10.3109/10408449509089885>.
- Harada, Masazumi, Shigeharu Nakachi, Koa Tasaka, Sakae Sakashita, Kazue Muta, Kohichi Yanagida, Rikuo Doi, Takako Kizaki, and Hideki Ohno. 2001. "Wide Use of Skin-Lightening Soap May Cause Mercury Poisoning in Kenya." *Science of The Total Environment* 269 (1–3): 183–87. [https://doi.org/http://dx.doi.org/10.1016/S0048-9697\(00\)00812-3](https://doi.org/http://dx.doi.org/10.1016/S0048-9697(00)00812-3).
- Henley, David E F. 2005. *Fertility, Food and Fever: Population, Economy and Environment in North and Central Sulawesi, 1600-1930*. Leiden: Brill.
- Idrus, Arifudin, Sukmandaru Prihatmoko, Hill. Gendoet Hartono, Fadlin Idrus, Ernowo Ernowo, Franklin Franklin, Moetamar Moetamar, and Iwan Setiawan. 2014. "Some Key Features and Possible Origin of the Metamorphic Rock-Hosted Gold Mineralization in Buru Island, Indonesia." *Indonesian Journal on Geoscience* 1 (1): 9–19. <http://ijog.bgl.esdm.go.id>.
- Ismawati, Yuyun. 2014. "Gold, Mercury and the next Minamata." *Indonesian Journal of Leadership, Policy and World Affairs*.
- Ismawati, Yuyun, Jundrich Petrlik, and Joe DiGangi. 2013. "Mercury Hotspots in Indonesia ASGM Sites : Poboya and Sekotong in Indonesia."
- James, Laurence P. 1994. "The Mercury 'Tromol' Mill: An Innovative Gold Recovery Technique, and a Possible Environmental Concern." *Journal of Geochemical Exploration* 50 (1): 493–500. [https://doi.org/https://doi.org/10.1016/0375-6742\(94\)90038-8](https://doi.org/https://doi.org/10.1016/0375-6742(94)90038-8).
- Kambey, Joice L., A. P. Farrell, and L. I. Bendell-Young. 2001. "Influence of Illegal Gold Mining on Mercury Levels in Fish of North Sulawesi's Minahasa Peninsula, (Indonesia)." *Environmental Pollution* 114 (3): 299–302. [https://doi.org/10.1016/S0269-7491\(01\)00163-4](https://doi.org/10.1016/S0269-7491(01)00163-4).
- Kocman, David, Milena Horvat, Nicola Pirrone, and Sergio Cinnirella. 2013. "Contribution of Contaminated Sites to the Global Mercury Budget." *Environmental Research* 125 (i): 160–70. <https://doi.org/10.1016/j.envres.2012.12.011>.
- Langston, James Douglas, Muhammad I Lubis, Jeffrey A Sayer, Chris Margules, Agni Klintuni Boedhihartono, and Paul H G M Dirks. 2015. "Comparative Development Benefits from Small and Large Scale Mines in North Sulawesi, Indonesia." *The Extractive Industries and Society* 2 (3): 434–44. <https://doi.org/http://dx.doi.org/10.1016/j.exis.2015.02.007>.
- Lasut, Markus T., Yoshiaki Yasuda, Evan N. Edinger, and Jane M. Pangemanan. 2010. "Distribution and Accumulation of Mercury Derived from Gold Mining in Marine Environment and Its Impact on Residents of Buyat Bay, North Sulawesi, Indonesia." *Water, Air, and Soil Pollution* 208 (1–4): 153–64. <https://doi.org/10.1007/s11270-009-0155-0>.
- Lihawa, Fitriyane, and Marike Mahmud. 2019. "The Content of Mercury in Sediments around Artisanal Small-Scale Gold Mining (ASGM) Bumela District, Gorontalo Regency, Gorontalo Province, Indonesia." *IOP Conference Series: Earth and Environmental Science* 314 (August): 012016. <https://doi.org/10.1088/1755-1315/314/1/012016>.
- Limbong, D., J. Kumampung, D. Ayhuan, T. Arai, and N. Miyazaki. 2005. "Mercury Pollution Related to Artisanal Gold Mining in North Sulawesi Island, Indonesia." *Bulletin of Environmental Contamination and Toxicology* 75 (August): 989–96.

<https://doi.org/10.1007/s00128-005-0847-0>.

- Male, Yusthinus Thobias, Amanda Jean Reichelt-Brushett, Matt Pocock, and Albert Nanlohy. 2013. "Recent Mercury Contamination from Artisanal Gold Mining on Buru Island, Indonesia - Potential Future Risks to Environmental Health and Food Safety." *Marine Pollution Bulletin* 77 (1–2): 428–33. <https://doi.org/10.1016/j.marpolbul.2013.09.011>.
- Nakagawa, Junji. 2008. *Freeport's Grasberg/Ertsberg Mine in West Papua, Indonesia. Opportunities and Challenges for Foreign Investment in the APEC Region: Case Studies*.
- Nakazawa, Koyomi, Osamu Nagafuchi, Tomonori Kawakami, Takanobu Inoue, Kuriko Yokota, Yuka Serikawa, Basir Cyio, and Rosana Elvince. 2016. "Human Health Risk Assessment of Mercury Vapor around Artisanal Small-Scale Gold Mining Area, Palu City, Central Sulawesi, Indonesia." *Ecotoxicology and Environmental Safety* 124 (February): 155–62. <https://doi.org/10.1016/j.ecoenv.2015.09.042>.
- Pacyna, E. G., J. M. Pacyna, K. Sundseth, J. Munthe, K. Kindbom, S. Wilson, F. Steenhuisen, and P. Maxson. 2010. "Global Emission of Mercury to the Atmosphere from Anthropogenic Sources in 2005 and Projections to 2020." *Atmospheric Environment* 44 (20): 2487–99. <https://doi.org/10.1016/j.atmosenv.2009.06.009>.
- Purwana, Rachmadhi. 2013. "Safety and Health in Mining in Indonesia." In *Occupational Safety and Health in Mining Anthology on the Situation in 16 Mining Countries*, edited by Kaj Elgstrand and Eva Vingård, Arbete och Hälsa, 47(2):42–49. Gothenburg: Vetenskaplig Skriptserie.
- Rianse, Usman, Weka Gusmiarty Abdullah, Abdi La Abdi, Ilma Sarimustaqiyma Rianse, Zulfikar La Zulfikar, Wa Kuasa Baka, La Ode Midi, and Weka Widayati. 2015. "The Impact of the Gold Mining on the Social, Economic, and Cultural in the Bombana District Southeast Sulawesi Province." *International Journal of Sustainable Tropical Agricultural Sciences* 1 (1): 53–65.
- Schulz, C, J Angerer, U Ewers, and M Kolossa-Gehring. 2007. "The German Human Biomonitoring Commission." *International Journal of Hygiene and Environmental Health* 210 (3–4): 373–82. <https://doi.org/10.1016/j.ijheh.2007.01.035>.
- Seccatore, Jacopo, Marcello Veiga, Chiara Origliasso, Tatiane Marin, and Giorgio De Tomi. 2014. "An Estimation of the Artisanal Small-Scale Production of Gold in the World." *Science of The Total Environment* 496: 3–8. <https://doi.org/10.1016/j.scitotenv.2014.05.003>.
- Sera, K., S. Futatsugawa, and S. Muraio. 2002. "Quantitative Analysis of Untreated Hair Samples for Monitoring Human Exposure to Heavy Metals." *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms* 189: 174–79. [https://doi.org/10.1016/S0168-583X\(01\)01034-5](https://doi.org/10.1016/S0168-583X(01)01034-5).
- Sera, K, S Futatsugawa, and K Matsuda. 1999. "Quantitative Analysis of Untreated Bio-Samples." *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 150 (1): 226–33. [https://doi.org/http://dx.doi.org/10.1016/S0168-583X\(98\)01071-4](https://doi.org/http://dx.doi.org/10.1016/S0168-583X(98)01071-4).
- Sippl, Kristin, and Henrik Selin. 2012. "Global Policy for Local Livelihoods: Phasing out Mercury in Artisanal and Small-Scale Gold Mining." *Environment: Science and Policy for Sustainable Development* 54 (3): 18–29. <https://doi.org/https://doi.org/10.1080/00139157.2012.673452>.
- Spiegel, Samuel J. 2012. "Governance Institutions, Resource Rights Regimes, and the Informal Mining Sector: Regulatory Complexities in Indonesia." *World Development* 40 (1): 189–205. <https://doi.org/https://doi.org/10.1016/j.worlddev.2011.05.015>.
- Spiegel, Samuel J, and Marcello M Veiga. 2010. "International Guidelines on Mercury Management in Small-Scale Gold Mining." *Journal of Cleaner Production* 18 (4): 375–85. <https://doi.org/https://doi.org/10.1016/j.jclepro.2009.10.020>.

- Sulaiman, Rini, R Baker, Budi Susilorini, Kevin H Telmer, and Samuel Spiegel. 2007. "Removal of Barriers to Introduction of Cleaner Artisanal Gold Mining and Extraction Technologies: Indonesia Country Project Summary. Report to the United Nations Industrial Development Organization." <http://iwlearn.net/iw-projects/1223/reports/indonesia/indonesia-country-report>.
- Telmer, Kevin H, and Marcello M Veiga. 2009. "World Emissions of Mercury from Artisanal and Small Scale Gold Mining." In *Mercury Fate and Transport in the Global Atmosphere Emissions, Measurements and Models*, edited by N. Pirrone and R. Mason, 131–72. New York: Springer. <https://doi.org/10.1007/978-0-387-93958-2>.
- UNEP (United Nations Environment Programme). 2008. "The Global Atmospheric Mercury Assessment: Sources, Emissions and Transport." UNEP-Chemicals, Geneva. <https://www.unenvironment.org/resources/report/technical-background-report-global-atmospheric-mercury-assessment-2008-0>.
- Veiga, Marcello M., Peter a. Maxson, and Lars D. Hylander. 2006. "Origin and Consumption of Mercury in Small-Scale Gold Mining." *Journal of Cleaner Production* 14 (3–4): 436–47. <https://doi.org/10.1016/j.jclepro.2004.08.010>.
- Veiga, Marcello M, Denise Nunes, Bern Klein, Janis A Shandro, P Colon Velasquez, and Rodolfo N Sousa. 2009. "Mill Leaching: A Viable Substitute for Mercury Amalgamation in the Artisanal Gold Mining Sector?" *Journal of Cleaner Production* 17 (15): 1373–81. <https://doi.org/https://doi.org/10.1016/j.jclepro.2009.03.012>.
- Verbrugge, Boris, and Sara Geenen. 2019. "The Gold Commodity Frontier: A Fresh Perspective on Change and Diversity in the Global Gold Mining Economy." *Extractive Industries and Society* 6 (2): 413–23. <https://doi.org/10.1016/j.exis.2018.10.014>.
- Vogt, Craig. 2012. "International Assessment of Marine and Riverine Disposal of Mine Tailings." In *Proceedings of the 34th Meeting of the London Convention and the 7th Meeting of the London Protocol, London, UK. Vol. 1*.


Commented [L91]: O is replaced with lower- case o


Commented [L92]: dots is inserted after abbreviated first names if authors

Commented [L93]: 83 is replaced with 16(3-4)

Commented [L94]: lower-case c is substituted with capital one

## Review





#### Article

## Artisanal and Small-Scale Gold Mining Activities and Its Mercury Exposures in Gorontalo Utara Regency, Indonesia

Yayu Indriati Arifin<sup>1,\*</sup>, Masayuki Sakakibara<sup>2</sup>, Sayaka Takakura<sup>3</sup>, Mohamad Jahja<sup>4</sup>, Fitriyane Lihawa<sup>5</sup> and Koichiro Sera<sup>6</sup>

<sup>1</sup> Department of Geology, State University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia

<sup>2</sup> Graduate School of Sciences and Engineering, Ehime University, 2-5 Bunkyo-cho, Matsuyama, 790-8577, Japan

<sup>3</sup> Ehime University, 3 Bunkyo-cho, Matsuyama, 790-8577, Japan

<sup>4</sup> Department of Physics, State University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia

<sup>5</sup> Department of Educational Geography, University of Gorontalo, Jl. Jend. Sudirman No. 6, Kota Gorontalo, 96128, Indonesia

<sup>6</sup> Cyclotron Research Center, Iwate Medical University, 348-58 Tomegamori, Takizawa, Iwate 020-0173, Japan

\* Correspondence: yayu\_arifin@ung.ac.id; Tel.: +81-089-927-9649; Fax: +81-089-927-9640

Received: date; Accepted: date; Published: date

**Abstract:** We studied and Small-scale gold mining activities and their mercury emissions in the Utara Regency of the Province, Sulawesi Island, Indonesia, have been studied. The mining occurs in many areas of this region, but especially in the three villages Hulawa, Langata and Langata Barat. Mining activities in the Hulawa village are the oldest (19th century) in the regency, whereas mining in the Langata -village began just fifteen years ago. Furthermore, the activity level changes

**Commented [095]:** Instructions for Authors (IfA): <https://www.tandfonline.com/action/authorSubmission?show=instructions&journalCode=gtec20>

Please include a word count for your paper.

**Commented [096]:** IfA: Write in an objective, neutral manner in the third person, not in the 1st person 'we' or 'I'.



rapidly from time to time depending on the fluctuations of the gold market. Women and children are involved in the gold mining processes. Female workers crush ore, process sediment wastes, and pan for gold in rivers. Children assist their mothers in these activities and play around the mining sites. Gold mining in the Utara Regency is estimated to produce about 290 kg of gold and emits approximately 860 kg of mercury into the environment annually. The impact of such large mercury released into the environment ~~and human population~~ have caused environmental degradation and contamination of ~~fish as food for humans and fish~~. Collaborative works of stakeholders are urgently needed to find solutions ~~to the~~ problems related to small-scale gold mining such as marginalization and environmental pollution.

**Keywords:** artisanal and small-scale gold mining; mercury exposure; human health problems; Gorontalo Utara; Indonesia

## 1. Introduction

Indonesia is the world's second-largest source of mercury emissions from artisanal and small-scale gold mining (ASGM) after China (Veiga, Maxson, and Hylander 2006; Pacyna *et al.* 2010; Ismawati 2014). The top ten global ASGM mercury emitters in 2000 and 2005 are shown in Figure 1 (Telmer and Veiga 2009; Cordy *et al.* 2011; Pacyna *et al.* 2010; Veiga, Maxson, and Hylander 2006; ~~Y~~ Ismawati 2014). Annual mercury emissions from ASGM in Indonesia in 2000, 2005, and 2013 were approximately 125, 150, and 175 tons, respectively. In a five-year period (from 2000 to 2005), Indonesia's ASGM mercury emissions were increased by approximately 20 %.

According to UNEP (2008), there are 115 ASGM industries distributed evenly throughout Indonesia that collectively produce 60 tons of gold per annum. The number of ASGM sites ~~and miners~~ in Indonesia is estimated to be ca. 900 ~~involving about~~ 250,000 ~~miners~~ people, respectively (Purwana 2013). The largest gold mining site in Indonesia is located near Puncak Jaya in Papua. Tailings from this site are dumped in Timika, in eastern Indonesia (Nakagawa 2008; Vogt 2012). Here ASGM activities can be found in several places, e.g.: Jayapura, Nabire, Paniai and Manokwari. ASGM activities also occur in Maluku Province, Gunung Botak and Gogoreaon Buru Island, Maluku Province (Male *et al.* 2013; Idrus *et al.* 2014).

~~Moreover,~~ Several ASGM locations have also been reported in Sulawesi, including Poboya in Palu (Basir-Ceyio *et al.* 2012). According to Iswamaty, Petrlik, and DiGangi (~~Yuyun Ismawati, Petrlik, and DiGangi~~ 2013), ASGM miners' activities in Poboya use 100 tons of mercury annually (Nakazawa *et al.* 2016). ~~While~~ Basyir-Cyio, *et al.* (2012) estimates that ~~they used~~ about 34 tons of mercury ~~are used~~ and produced 22.4 - 45 kg gold daily (~~Basir-cyio et al. 2012~~). High concentrations ~~(above the limit)~~ of mercury vapor from ASGM were detected ~~elsewhere~~ in Palu City, and almost all residents of the city are exposed (Nakazawa *et al.* 2016). Several ASGM has been reported in Mamuju of Sulawesi Barat Province, TanaToraja and Luwu of Sulawesi Selatan Province and the Bombana of Sulawesi Tenggara Province. The Bombana ASGM has been reported to have negative impact on social, economic and cultural ~~status of the~~ people around ~~the~~ ASGM (Rianse *et al.* 2015).

There are four other ASGM locations in Sulawesi Utara Province (Basir-Ceyio *et al.* 2012), at Talawaan (Filho and Villas-bôas 2004; Bose-O'Reilly *et al.* 2010), Buyat (Limbong *et al.* 2005; Kambey, Farrell, and Bendell-Young 2001), Tobongon and Tanoyan in the Bolaang Mongondow Regency. Mercury emissions from Talawaan alone are 70 t per annum (Lasut *et al.* 2010), which is seven times ~~the~~ value reported earlier by Telmer and Veiga (2009).

At least there is one ASGM site in each regency of the Gorontalo Province: (1) Pohuwato Regency: Gunung Pani and Bulontio; (2) Boalemo Regency: Bilato; (3) Gorontalo Regency: Bumela

Commented [097]: Daily? Monthly?...

Commented [098]: What is the limit? Here the limit must be defined/mentioned concretely.

(Lihawa and Mahmud 2019) (4) Bonebolango Regency: Tulabolo and Mopuya; and (5) Gorontalo Utara Regency: Hulawa and Ilangata villages.

Gold deposits in the Gorontalo Utara Regency have existed in Wubudu breccias within Boliyohuto diorite of the middle Miocene age (Carlile, Digdowirogo, and Darius 1990). Gold in hydrothermal breccias and quartz veinlets has been reported in Sumalata and Boliyohuto subdistricts (Carlile, Digdowirogo, and Darius 1990). Gold mining in the Hulawa village is considered to date back to Indonesia Dutch era (Henley 2005).

This study describes mercury pollution from ASGM sites in the Gorontalo Utara Regency, where gold mining activity is intense, but information is scarce. The report describes how miners work and what the mercury emission situation is, as a basis for further discourse on health risks in and around these mining sites.

## 2. Materials and Methods

### 2.1. Study Area

The Hulawa (meaning 'gold' in the Gorontalo language) mining area (Figure 2) is also known as the Buladu mining area in Sumalata. This area lies within the Buladu River watershed, between the Pasolo and Padengo villages, approximately 109 km northeast of Gorontalo City, the capital of Gorontalo Province. From Gorontalo City, the area can be reached by car in approximately four hours. Buladu has a typical equatorial climate with two seasons (rainy: November–April; dry: May–October), with the highest rainfall occurring in December (annual average 1,210 mm). The temperature is 27 °C on average. The Buladu River watershed drains from the mountain peaks with the highest point being 2,000 m, on the northern coast of the Sumalata district. The distance from the mountain peaks to the Sulawesi Sea is approximately 10 km. Several rivers run through this watershed, including the Buladu and Wubudu rivers.

The Ilangata and Ilangata Barat mining areas (Figure 2) comprise the Anggrek mining area. The area lies within the Anggrek River watershed between the Diata, Lunggulo, and Lantolo villages, which are approximately 68 km northwest of Gorontalo City. It is three hours by car from Gorontalo City. The Anggrek River watershed drains from the peaks of several hills with elevations of 300 m at most. The distance from the peaks to the sea is approximately 5 km. Number of rivers, including the Anggrek and Ilangata rivers, flow through the Anggrek watershed.

Residents of Hulawa, Ilangata and Ilangata Barat are exposed directly and indirectly to mercury from mercury vapor of ASGM process and sea fish consumption, while residents of other subdistricts (Kwandang, Monano and Tolinggula) are treated as control group since they might only be exposed to mercury through sea fish consumption.

### 2.2. Sample and data collection

The ASGM locations within Gorontalo Utara regency have been visited many times over the course of one month each year, from 2012 to 2014. Data on mercury use during ASGM and the activities of miners and their families were obtained via direct interviews with miners, mine owners, and local government officials. These interviews were conducted on-site at the mining locations in the Gorontalo Utara Regency (Figure 2). The simple questionnaire consisted of number of workers, number of gold trommels, amount of ore filled into trommels, and amount of mercury consumed and gold produced in a single mill process. This was done in Hulawa, Ilangata and Ilangata Barat ASGM sites (n=23): 6 ASGM operators, 2 panners, 3 villagers in Hulawa, 6 ASGM operators, 4 villagers and 2 gold shop workers in Ilangata and Ilangata barat, 2 officers in Kwandang). The interviewed [were](#) key-actors, including mine site managers, head of villages, [and](#) elderly people who were working as ASGM miners 20 to 40 years ago.

Data obtained from different sources were compared with each other via triangulation procedure, for example, number of miners from the owner, from direct investigation and from local officers other to keep accountable. Any information ~~we~~ obtained from participants was recorded and is used in triangulation process. For instance, we recorded the following information while communicating with villagers who have a motorcycle repair kiosk in Ilangata, his friend said that: "he started that business after he earned about 6000 US-\$ for a week from ASGM one year ago". Once we were in Hulawa village, a man who got training on the Minamata diseases problem informed us that in nearby villages live three people who were suspected to have tremor disturbance related to their former activities as ASGM miners in Hulawa. The first person showed tremor during our conversation and he has a problem on oral communication already. The second one, also showing tremor with a lesser degree, said that: "We were working as ASGM miners, dealing with mercury without safety equipment during smelting; sometimes they put gold amalgam in their mouths to sell later on". They also consumed fish from contaminated rivers, the estuary, and the sea.

Commented [099]: ??

Information on the economy and social life in ASGM were also obtained from interviews with miners, managers of ASGM businesses, and villagers. Compared to large-scale mining, ASGM activities have more positive economic impacts on local communities, since the capital is shared within local investors, miners, and villagers (Langston *et al.* 2015). Economy and social life information, in this case, will enrich our understanding on ASGM activities, its impacts on the environment and human health, and ongoing struggles against social marginalization in Indonesia's gold mining sector (Spiegel and Veiga 2010).

### 2.3. Analytical Procedure

The health effects on ASGM miners and villagers were examined by collecting scalp hair strands from 115 donors from five subdistricts. The hair strand from the occipital region was cut close to the skull, collected, put in a clear plastic bag, and labeled accordingly. Mercury content in hairs was determined using ~~p~~Proton ~~i~~nduced X-ray ~~e~~mission (PIXE) at Iwate Medical University, Japan. ~~The method was reported elsewhere (Sera et al. 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera et al. 2017). Level of mercury is associated with a total of mercury weight divided by total weight of the sample.~~ The procedure is known as a standard-free method for hairs without powdering and acid treatment as reported in previous studies (Sera, Futatsugawa, and Matsuda 1999; Sera, Futatsugawa, and Murao 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera 2017). ~~We adopt~~ The widely accepted exposure limit values for mercury concentration in hair ( $\mu\text{g/g}$ ) as suggested by the German Human Biomonitoring Commission (HBM) as the following categories ~~are adopted~~: concentration ( $\mu\text{g/g}$ ) below 1.0 are considered normal, 1.0 - 5.0 are an alert level, and above 5.0 are indicative of a serious risk of health effects.

Commented [0100]: IfA: Abbreviations are to be introduced upon first mentioning of the fully spelled term (initial letters of each word in lower-case, not capital).

Commented [0101]: IfA: Write in an objective, neutral manner in the third person, not in the 1st person 'we' or 'I'.

### 2.4. Neurological Assessment

The subjects with elevated hair mercury concentration were examined clinically according to a questionnaire especially for subjective symptoms, and the use of mercury-containing soap (for female only), and a neurological test. Neurological examination was performed by a medical doctor on participants, using a UNIDO protocol which consists of ten symptoms: (1) signs of bluish discoloration of gums; (2) rigidity and ataxia (walking and standing); (3) alternating movements or dysdiadochokinesia; (4) irregular eye movements or nystagmus; (5) field of vision; (6) knee jerk reflex and biceps reflex; (7) Babinski reflex and labial reflex; (8) salivation and dysarthria; (9) sensory examination and (10) tremor: tongue, eyelids, finger to nose, pouring, posture holding, and the Romberg test. ~~We scored 1 was given~~ for positively-observed symptoms, and 0 for negatively-observed symptoms. The sum of score for every participant is calculated and analyzed

### 2.5. Description of sites and processes in of ASGM

Ore is manually excavated from nearby hills or mountains (Figure 3a) using vertical holes in the ground (Figure 3b). Traditional tools such as broad hoes, bars and simple pulleys are used for the ore excavation (Figure 3c). The ore is manually crushed (Figure 3d) using hammers or other percussion tools (Figure 3e). Some processing plants use homemade mechanical crushers. The ore is then packed into sacks and manually transported to the processing plant (Figure 3f).

At the processing plant, crushed ore (30–40 kg) is fed into a trommel (known as “Tromol” in the local language (James 1994), a steel mill grinder where gold is extracted from ore (Male et al. 2013) (Figure 4a) with water (Figure 4b) and several decimeter-sized rocks (Figure 4c) and milled for 3–4 hours until the material is fine enough to release the gold. Subsequently, approximately 500 g of mercury are added to the trommel (Figure 4d), which then rotates for 30 minutes until the amalgamation has completed. This amalgamation process is the major gold production method for ASGM miners in Gorontalo Province.

After 30 minutes, the material is discharged (Figure 4e), and the tailings are manually separated using bare hands (Figure 4f). The tailings are washed and rinsed several times (Figures 4g – 4j) to separate the gold from the unamalgamated mercury. The mixture is filtered using a fine cloth to separate the mercury amalgam (Figure 4k). The gold amalgam is recovered from this process as shown in Figure 4l.

After milling the gold, the amalgam is recovered, and the waste sediment is kept (Figure 4m) for a second recovery process. Then, mercury-contaminated water is released into the environment (Figure 4n). The amalgams are then smelted to obtain gold bullion. Gold smelters with simple tools smelt the gold outdoors (Figure 4o) or inside houses and gold shops. Gold bullion is recovered from the smelting process (Figure 4p).

The secondary process begins when miners, usually women, and children, wash sediment wastes from the primary milling process in running water. The milling process is then repeated, as shown in Figure 4a through Figure 4l, with the difference being that the trommel is generally rotated by a wheel powered with river water (Figure 4n).

Tertiary processing involves bagging the waste from the secondary processing and sending it to another processing plant to undergo cyanidation. There is no cyanidation plant in the Gorontalo Utara Regency. However, there is a cyanidation plant in the other Regency of Gorontalo Province. Numerous miners pan in the Wubudu River estuary (Figure 4q) and (Figure 4r) where the mercury enters the environment (air, soil, and water) directly.

Smelting activities occur in gold shops in the Ilangata Barat village. The owners of these gold shops determine the gold quality (amount of gold in “dore”) via these processes. The amount of gold in a dore varies from 40 – 80% of the weight. A dore with 80 % quality can be purchased for approximately US \$30,-/gram. Mining wastes have solid and liquid forms that contaminate river waters and soils. Gold bullion is extracted after smelting the amalgam, which emits mercury vapor.

### 3. Results

#### 3.1. Mercury Emissions

Because ASGM activities are unregistered, no exact data were available on the number of workers; therefore, it is difficult to estimate the amount of gold produced, and mercury used, for ASGM activities throughout Indonesia and its local provinces or regencies. This difficulty in estimating amounts of gold produced and mercury used by ASGM throughout the world is described elsewhere (Seccatore et al. 2014). We used direct observation and interviews to get information and compared it with data reported by several researchers. Data on mercury imported to Indonesia will be used as the upper limit of mercury lost to ASGM. Socio-economic factors which should be

considered when dealing with ASGM (Sulaiman et al. 2007; Spiegel 2012; Langston et al. 2015; Spiegel and Veiga 2010), will be elaborated further to discuss any possible scenario or solutions on sustainable development in local ASGM site areas.

Data on the number of mine workers, population and mercury used during the milling process in each village are listed in Table 1. These figures were estimated based on each milling process requiring 0.5 kg mercury for one trommel mill to extract gold from ca. 30-40 kg of crushed ore. Assuming all trommel mills are operating during the milling process, a maximum of 275 kg of mercury is needed (these data and the amount of mercury emitted per gram of gold produced allows us to estimate the mercury emitted into the environment). Each primary process produced ca. 3 g of gold per trommel mill. The secondary processing of waste produces ca. 0.5 g of gold per trommel mill. We also assume the miners only work five days a week and sell the gold on Saturday. The ratio of mercury emissions to gold produced is ca. 2:1 (Darmutji 2003). The amount of gold produced by the process is ca. 1134 g. Thus, the mercury emissions are ca. 2.2 kg per milling process. The total annual mercury emission in the Gorontalo Utara Regency is approximately 572 kg. If we used 1-3 grams of mercury lost per gram of gold produced, the mercury emission ranges from 286 – 858 kg (Veiga et al. 2009).

### 3.2. Mercury Concentration in Hair Samples According to Geographic Location

The distributions and range of mercury levels in 115 hair samples collected from the five subdistricts are shown in Figure 5. Mercury concentrations in all hair donors are more than 1 µg/g, which indicates the concentrations are already within the alert level range according to HBM (Schulz et al. 2007). The number of subjects with high mercury levels over 10 µg/g was 10 (40 %), 7(30-4 %), and 4 (8.5 %) in Anggrek, Sumalata, and the control group (Kwandang, Monano and Tolinggula), respectively.

According to the Kolmogorov-Smirnov test, the distribution of mercury in the hair data from the Gorontalo Utara Regency was not normal; instead, it had a log normal distribution (Figure 6). The geometric mean is more suitable for log normal distribution data. Mean mercury level (+/- standard deviation) of Gorontalo Utara regency is 6.7 (2.0) µg/g and is higher than minimum limit for danger level (5.0 µg/g) according to HBM. Only residents of Tolinggula has average hair mercury level less than 5.0 µg/g. This may due to their residence away from the seaside; their common diet along with rice is chicken. Even though one male and two females have mercury higher than 6.0 µg/g, the Tolinggula group could be considered as a control group in Gorontalo Utara regency.

The data were plotted using a box-and-whisker plot to make a clear view of data distribution in each residence and among Gorontalo Utara regency data. Each Box is filled with the horizontal line representing the position of the median. The position of the median and mean relative to HBM was used to put the population into three categories according to HBM. Box plot of mercury distribution according to geographical location is shown in Figure 5, the results of Sumalata and Anggrek are positioned higher than others, which means that their hair average mercury is elevated in the regency. Other subdistricts (Tolinggula and Kwandang) have no known ASGM activities, but the population lives close to seashore (their common diet is fish from Sulawesi sea, same as the population of Anggrek and Sumalata).

### 3.3. Mercury concentration for male and female

The lognormal distribution of hair mercury levels in males and females is shown in Figure 6. Five females had mercury levels greater than 25 µg/g, and none of them worked as ASGM miners. Those five females may have been exposed to mercury from another source. The elevated hair mercury levels that were above average (7.1 µg/g) and even the highest (17.9 µg/g) mercury level were found among the ASGM miners. The average hair mercury levels for all, male and female inhabitants in the Monano subdistrict are 5.7, 5.0 and 6.2 µg/g, respectively. These levels show that

there are no significant differences between the mean hair mercury of males and females in that subdistrict. While the average of all mercury hair content for females is 8.1 µg/g (more than 30 percent higher than males (6.0 µg/g), such conditions were also found for the subgroups of Kwandang and Sumalata. The condition in Anggrek is even higher (the average mercury for females is three times than the males). The large discrepancy of mercury levels between female and male inhabitants suggests that female inhabitants may be receiving mercury from another source (e.g., whitening cream, soap, etc).

The clinical symptoms were observed in most of the females with mercury higher than 50 mg/g (the minimum level at which Minamata disease symptoms are seen (Harada 1995)) are listed in Table 2, indicative of inorganic-mercury poisoning only. All the subjects have habitual use of whitening cream that are common in Gorontalo Utara regency. The mercury concentration on the whitening creams are shown in Table 3. It seems that elevated mercury concentration among female inhabitants of Hulawa and Ilangata may be caused by direct mercury contact from habitual use of whitening creams. Harada et al. (2001) found the similar cases in Kenya which were caused by female were elevated hair mercury levels were using soaps with high mercury concentrations.

#### 4. Discussion

##### 4.1. ASGM operation system and Mercury losses

Figure 7 shows the ASGM operation system which is used in Gorontalo Utara Regency; the system used in North Gorontalo is similar to those reported elsewhere (Veiga *et al.* 2009). Gold recovery systems are based on whole ore amalgamation using trommel, gravitation and panning methods were found in Wobudu river estuary and Anggrek river banks. There are two possible routes for mercury contamination from ASGM into the environment and humans. During the milling process, liquid mercury could contaminate miners and the environment. Working with mining wastes (tailings), and direct exposure to liquid mercury during panning, are also contamination routes. Mercury vapor could contaminate the smelters and environment during smelting.

For comparison, we also estimated the total annual mercury emissions in the Gorontalo Utara Regency from ASGM using data on Indonesian mercury emissions (Pacyna *et al.* 2010) and the number of known ASGM sites (Kocman *et al.* 2013). The rate for Indonesian mercury emissions from ASGM may be close to 200 t annually. The average emission from each mining site is approximately 200 kg/year based on the 900 known ASGM sites (Kocman *et al.* 2013). The three mining sites in the Gorontalo Utara Regency emit approximately 600 kg/year. Ismawaty (2014) estimated the ASGM production ranged from 65-130 t in 2012 (Y Ismawati 2014). Thus, approximately 170 – 260 t of mercury was lost to the environment based on the multiplication factor from Darmutji (2003). Using the 1:1 to 3:1 ratio proposed by Ismawaty (2014), the amount of mercury lost was 65 – 390 t. These estimates agree with the total mercury import data for Indonesia in 2012, which was approximately 368 t.

##### 4.2. Health Issues

The involvement of women and children workers is unique to the Hulawa mining site. These women and children come from the Buladu area. They work with waste stored in wooden boxes, and sluice and pan in the Wobudu River estuary. Women and children typically work on mercury-bearing mining wastes during the secondary processing stage (Figures. 8a and 8b). Their bare hands and feet are directly exposed to the mining waste, which allows mercury to enter their body via dermal absorption.

Health problems in miners and inhabitants were documented by medical doctors using a standard questionnaire and neurological test tools (Figures. 9a and 9b). Ten neurological disorder indicators/symptoms were observed. Bluish gums and tremors are major symptoms observed in the miners and inhabitants of the Gorontalo Utara Regency. Several explanations could be addressed based on the evidence; first, they have experienced prolonged contact with mercury, and second, the mercury

has entered their body via the food chain (Basir-cyio et al. 2012; Nakazawa et al. 2016; Bose-O'Reilly et al. 2010). Both can happen simultaneously for the miners and inhabitants of these mining sites. Further investigations are needed to explain the evidence and determine the most probable mercury contamination routes in the Gorontalo Utara Regency.

Without pollution controls and environmental cleanup, further mercury contamination of the environment (bottom sediments, water, and fish) and human population around ASGM sites within the Gorontalo Utara Regency is inevitable (Arifin, Sakakibara, and Sera 2015). Veiga and Hinton (Seccatore *et al.* 2014) reported that gold bullion contains two to five percent mercury just after smelting, and such gold bullion is refined at gold shops in towns within the Gorontalo Utara Regency with few emission controls. Mercury pollution from gold shops, like those observed in Colombian cities, is of great concern (Cordy *et al.* 2013).

#### 4.3. Socio-economic aspects of ASGM system in Gorontalo Utara Regency

The intensification of ASGM activities at the historical gold mining site (hulawa village) and expansion to the new gold mining site (ilangata and ilangata barat villages) has been seen as advantage of ASGM over large scale gold mining (LSGM)(Adler Miserendino *et al.* 2013). The ability of ASGM activities to cut down production cost through their reliance on flexible informal labor. The ASGM activities are driven by local investors which are intimately entangled with socio-political structures and land tenure systems (Verbrugge and Geenen 2019).

The hierarchy of people working in ASGM system in Gorontalo Utara regency can be explained using the network chart introduced by Langston et al. (2015), with slight modification mainly related [to with](#) the number of ASGM miners. In general, a small number of miners (up to 3 to 4 people) work at the excavating site or processing site; these miners may be family members or friends. Hence supervisor, landowner position, [and](#) even investor itself can be the same person.

Local miners are formerly farmers or fisherman and live nearby in their houses, while migrants usually live inside the processing site camps, which are very simple structures and lack sanitation facilities. The basic needs (foods, drinks, cigarettes) were available in kiosk/gold shop [s](#) which may belong to cooperation or investor/landowner/supervisor.

Positive economic impacts of ASGM activities to miners and inhabitants were not easily seen in Gorontalo Utara regency since there is no significant difference in economic activities in ASGM sites and non ASGM sites. In a stakeholder meeting, ASGM miners, local government, and investor seem to have opposing viewpoints on issues such as sustainable development, environmental protection, [and](#) negative impacts of ASGM on health.

Local governments and inhabitants would like to embrace long-term economic solution such as farming, fishery, and tourism, which could eventually replace ASGM. There [is are](#) ongoing researches on finding the ultimate solutions which fit to Gorontalo Utara Regency cases. The multidisciplinary research approach is being taken to help young local students who were living in the ASGM area to find solutions to their local problems. A transdisciplinary approach could facilitate the stakeholders in ASGM area together with the researcher to develop the ideas on sustainable development (Sippl and Selin 2012).

## 5. Conclusions

ASGM activities in the Gorontalo Utara Regency are dangerous to the environment, miners, and the local population. The amount of mercury used per mining cycle is approximately 275 kg, and approximately 2.2 kg of mercury is lost per cycle. Annually, ASGM in the Gorontalo Utara Regency produces approximately 286 kg of gold and emits approximately 572 kg of mercury. This estimate is probably conservative given the limited mining data available. The impact of such a large mercury



loss into the environment and human population has been to degrade the environmental conditions and contaminating humans and fish as demonstrated by Arifin et al. (2015).

Gorontalo Utara Regency is an essential agricultural (rice) and horticultural (dairy) production area in the Gorontalo Province and Indonesia. The ASGM driven economy did not bring sustainable development in the vicinity of ASGM sites and Gorontalo Utara regency as a whole. Solutions to the problem are being developed using ongoing multi and trans-disciplinary research approach through the collaboration of Indonesian and Japanese researchers.

**Author Contributions:** All authors contributed to the work presented in the manuscript.

**Acknowledgments:** The authors wish to thank the regent of Gorontalo Utara and its officers, and the owners of mining plants for allowing us to conduct interviews and take photographs. Thanks also to Syamsu Qamar Badu, the Rector of the State University of Gorontalo, for his support of this research.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Adler Miserendino, Rebecca, Bridget A. Bergquist, Sara E. Adler, Jean Remy Davée Guimarães, Peter S J Lees, Wilmer Niquen, P. Colon Velasquez-López, and Marcello M. Veiga. 2013. "Challenges to Measuring, Monitoring, and Addressing the Cumulative Impacts of Artisanal and Small-Scale Gold Mining in Ecuador." *Resources Policy* 38 (4). <https://doi.org/10.1016/j.resourpol.2013.03.007>.
- Arifin, Yayu, Masayuki Sakakibara, and Koichiro Sera. 2015. "Impacts of Artisanal and Small-Scale Gold Mining (ASGM) on Environment and Human Health of Gorontalo Utara Regency, Gorontalo Province, Indonesia." *Geosciences* 5 (2): 160–76. <https://doi.org/10.3390/geosciences5020160>.
- Basir-Ceyio, Muhammad, Uswah Hasanah, Isrun Nur, and Yuka Serikawa. 2012. "Gold Mining Activities and Its Impacts on Land Degradation in Central Sulawesi Indonesia." *Journal of Ecotechnology Research* 16(3-4): 79–83. <https://doi.org/https://doi.org/10.11190/jer.16.79>.
- Basri, Masayuki Sakakibara, and Koichiro Sera. 2017. "Current Mercury Exposure from Artisanal and Small-Scale Gold Mining in Bombana, Southeast Sulawesi, Indonesia—Future Significant Health Risks." *Toxics* 5 (1): 7. <https://doi.org/10.3390/toxics5010007>.
- Bose-O'Reilly, Stephan, Gustav Drasch, Christian Beinhoff, Saulo Rodrigues-Filho, Gabriele Roeder, Beate Lettmeier, Alexandra Maydl, Stefan Maydl, and Uwe Siebert. 2010. "Health Assessment of Artisanal Gold Miners in Indonesia." *Science of the Total Environment* 408 (4): 713–25. <https://doi.org/10.1016/j.scitotenv.2009.10.070>.
- Carlile, J. C., S. Digdowirogo, and K. Darius. 1990. "Geological Setting, Characteristics and Regional Exploration for Gold in the Volcanic Arcs of North Sulawesi, Indonesia." *Journal of Geochemical Exploration* 35 (1): 105–40. [https://doi.org/http://dx.doi.org/10.1016/0375-6742\(90\)90037-B](https://doi.org/http://dx.doi.org/10.1016/0375-6742(90)90037-B).
- Cordy, Paul, Marcello Veiga, Ben Crawford, Oseas Garcia, Victor Gonzalez, Daniel Moraga, Monika Roeser, and Dennis Wip. 2013. "Characterization, Mapping, and Mitigation of Mercury Vapour Emissions from Artisanal Mining Gold Shops." *Environmental Research* 125: 82–91. <https://doi.org/10.1016/j.envres.2012.10.015>.
- Cordy, Paul, Marcello M. Veiga, Ibrahim Salih, Sari Al-Saadi, Stephanie Console, Oseas Garcia, Luis Alberto Mesa, Patricio C. Velásquez-López, and Monika Roeser. 2011. "Mercury Contamination from Artisanal Gold Mining in Antioquia, Colombia: The World's Highest per Capita Mercury Pollution." *Science of The Total Environment* 410–411: 154–60. <https://doi.org/10.1016/j.scitotenv.2011.09.006>.

- Darmutji, Selinawati T. 2003. "Report, UNIDO Project No EG/GLO/01/G34, Overview of Project Site in Indonesia Talawaan and Galangan Areas."
- Filho, Saulo Rodrigues, and Roberto Villas-bôas. 2004. "Biogeochemical Assessment of Mercury Pollution in the São Chico Gold Mining Site - Brazilian Amazon Rio De Janeiro Novembro / 2004." In *Environmental Geochemistry in Tropical Counties*, 0–3. [????](#).
- Harada, Masazumi. 1995. "Minamata Disease: Methylmercury Poisoning in Japan Caused by Environmental Pollution." *Critical Reviews in Toxicology* 25 (1): 1–24. <https://doi.org/10.3109/10408449509089885>.
- Harada, Masazumi, Shigeharu Nakachi, Koa Tasaka, Sakae Sakashita, Kazue Muta, Kohichi Yanagida, Rikuo Doi, Takako Kizaki, and Hideki Ohno. 2001. "Wide Use of Skin-Lightening Soap May Cause Mercury Poisoning in Kenya." *Science of The Total Environment* 269 (1–3): 183–87. [https://doi.org/http://dx.doi.org/10.1016/S0048-9697\(00\)00812-3](https://doi.org/http://dx.doi.org/10.1016/S0048-9697(00)00812-3).
- Henley, David E. F. 2005. *Fertility, Food and Fever: Population, Economy and Environment in North and Central Sulawesi, 1600-1930*. Leiden: Brill.
- Idrus, Arifudin, Sukmandaru Prihatmoko, Hill. Gendoet Hartono, Fadlin Idrus, [Ernowo](#) Ernowo, [Franklin](#) Franklin, [Moetamar](#) Moetamar, and Iwan Setiawan. 2014. "Some Key Features and Possible Origin of the Metamorphic Rock-Hosted Gold Mineralization in Buru Island, Indonesia." *Indonesian Journal on Geoscience* 1 (1): 9–19. <http://ijog.bgl.esdm.go.id>.
- Ismawati, Y. 2014. "Gold, Mercury and the next Minamata." *Indonesian Journal of Leadership, Policy and World Affairs*.
- Ismawati, Yuyun, Jundrich Petrlik, and Joe DiGangi. 2013. "Mercury Hotspots in Indonesia ASGM Sites : Poboya and Sekotong in Indonesia."
- [Ismawati, Yuyun. 2014. "Gold, Mercury and the next Minamata."](#)
- James, Laurence P. 1994. "The Mercury 'Tromol' Mill: An Innovative Gold Recovery Technique, and a Possible Environmental Concern." *Journal of Geochemical Exploration* 50 (1): 493–500. [https://doi.org/https://doi.org/10.1016/0375-6742\(94\)90038-8](https://doi.org/https://doi.org/10.1016/0375-6742(94)90038-8).
- Kambey, Joice L., A. P. Farrell, and L. I. Bendell-Young. 2001. "Influence of Illegal Gold Mining on Mercury Levels in Fish of North Sulawesi's Minahasa Peninsula, (Indonesia)." *Environmental Pollution* 114 (3): 299–302. [https://doi.org/10.1016/S0269-7491\(01\)00163-4](https://doi.org/10.1016/S0269-7491(01)00163-4).
- Kocman, David, Milena Horvat, Nicola Pirrone, and Sergio Cinnirella. 2013. "Contribution of Contaminated Sites to the Global Mercury Budget." *Environmental Research* 125 (i): 160–70. <https://doi.org/10.1016/j.envres.2012.12.011>.
- Langston, James Douglas, Muhammad I Lubis, Jeffrey A Sayer, Chris Margules, Agni Klintuni Boedihartono, and Paul H G M Dirks. 2015. "Comparative Development Benefits from Small and Large Scale Mines in North Sulawesi, Indonesia." *The Extractive Industries and Society* 2 (3): 434–44. <https://doi.org/http://dx.doi.org/10.1016/j.exis.2015.02.007>.
- Lasut, Markus T., Yoshiaki Yasuda, Evan N. Edinger, and Jane M. Pangemanan. 2010. "Distribution and Accumulation of Mercury Derived from Gold Mining in Marine Environment and Its Impact on Residents of Buyat Bay, North Sulawesi, Indonesia." *Water, Air, and Soil Pollution* 208 (1–4): 153–64. <https://doi.org/10.1007/s11270-009-0155-0>.
- Lihawa, Fitriyane, and Marike Mahmud. 2019. "The Content of Mercury in Sediments around Artisanal Small-Scale Gold Mining ({ASGM}) Bumela District, Gorontalo Regency, Gorontalo Province, Indonesia." In *{IOP} Conference Series: Earth and Environmental Science*, 314:12016. *{IOP} Publishing*. <https://doi.org/10.1088/1755-1315/314/1/012016>.
- Limbong, D., J. Kumampung, D. Ayhuan, T. Arai, and N. Miyazaki. 2005. "Mercury Pollution

Related to Artisanal Gold Mining in North Sulawesi Island, Indonesia.” *Bulletin of Environmental Contamination and Toxicology* 75 (August): 989–96.  
<https://doi.org/10.1007/s00128-005-0847-0>.

Male, Yusthinus Thobias, Amanda Jean Reichelt-Brushett, Matt Pocock, and Albert Nanlohy. 2013. “Recent Mercury Contamination from Artisanal Gold Mining on Buru Island, Indonesia - Potential Future Risks to Environmental Health and Food Safety.” *Marine Pollution Bulletin* 77 (1–2): 428–33. <https://doi.org/10.1016/j.marpolbul.2013.09.011>.

Nakagawa, Junji. 2008. *Freeport’s Grasberg/Ertsberg Mine in West Papua, Indonesia. Opportunities and Challenges for Foreign Investment in the APEC Region: Case Studies*.

Nakazawa, Koyomi, Osamu Nagafuchi, Tomonori Kawakami, Takanobu Inoue, Kuriko Yokota, Yuka Serikawa, Basir Cyio, and Rosana Elvince. 2016. “Human Health Risk Assessment of Mercury Vapor around Artisanal Small-Scale Gold Mining Area, Palu City, Central Sulawesi, Indonesia.” *Ecotoxicology and Environmental Safety* 124 (February): 155–62.  
<https://doi.org/10.1016/j.ecoenv.2015.09.042>.

Pacyna, E. G., J. M. Pacyna, K. Sundseth, J. Munthe, K. Kindbom, S. Wilson, F. Steenhuisen, and P. Maxson. 2010. “Global Emission of Mercury to the Atmosphere from Anthropogenic Sources in 2005 and Projections to 2020.” *Atmospheric Environment* 44 (20): 2487–99.  
<https://doi.org/10.1016/j.atmosenv.2009.06.009>.

Purwana, Rachmadhi. 2013. “Safety and Health in Mining in Indonesia.” In *Occupational Safety and Health in Mining Anthology on the Situation in 16 Mining Countries*, edited by Ed Kaj Elgstrand and Eva Vingård, *Arbete och Hälsa* 47(2): 42-49+86. University of Gothenburg.

Rianse, Usman, Weka Gusmiarty Abdullah, Abdi La Abdi, Ilma Sarimustaqiyma Rianse, Zulfikar La Zulfikar, Wa Kuasa Baka, La Ode Midi, [and Weka Widayati](#), ~~and others~~. 2015. “The Impact of the Gold Mining on the Social, Economic, and Cultural in the Bombana District Southeast Sulawesi Province.” *International Journal of Sustainable Tropical Agricultural Sciences* 1-(1): [53-65](#).

Schulz, C., J. Angerer, U. Ewers, and M. Kolossa-Gehring. 2007. “The German Human Biomonitoring Commission.” *International Journal of Hygiene and Environmental Health* 210 (3–4): 373–82. <https://doi.org/10.1016/j.ijheh.2007.01.035>.

Seccatore, Jacopo, Marcello Veiga, Chiara Origliasso, Tatiane Marin, and Giorgio De Tomi. 2014. “An Estimation of the Artisanal Small-Scale Production of Gold in the World.” *Science of The Total Environment* 496: 3–8. <https://doi.org/10.1016/j.scitotenv.2014.05.003>.

Sera, K., S. Futatsugawa, and S. Muraio. 2002. “Quantitative Analysis of Untreated Hair Samples for Monitoring Human Exposure to Heavy Metals.” *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms* 189: 174–79.  
[https://doi.org/10.1016/S0168-583X\(01\)01034-5](https://doi.org/10.1016/S0168-583X(01)01034-5).


Sera, K., S. Futatsugawa, and K. Matsuda. 1999. “Quantitative Analysis of Untreated Bio-Samples.” *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 150 (1): 226–33. [https://doi.org/http://dx.doi.org/10.1016/S0168-583X\(98\)01071-4](https://doi.org/http://dx.doi.org/10.1016/S0168-583X(98)01071-4).

Sippl, Kristin, and Henrik Selin. 2012. “Global Policy for Local Livelihoods: Phasing out Mercury in Artisanal and Small-Scale Gold Mining.” *Environment: Science and Policy for Sustainable Development* 54 (3): 18–29. <https://doi.org/https://doi.org/10.1080/00139157.2012.673452>.

Spiegel, Samuel J. 2012. “Governance Institutions, Resource Rights Regimes, and the Informal Mining Sector: Regulatory Complexities in Indonesia.” *World Development* 40 (1): 189–205.  
<https://doi.org/https://doi.org/10.1016/j.worlddev.2011.05.015>.

- Spiegel, Samuel J., and Marcello M. Veiga. 2010. "International Guidelines on Mercury Management in Small-Scale Gold Mining." *Journal of Cleaner Production* 18 (4): 375–85. <https://doi.org/https://doi.org/10.1016/j.jclepro.2009.10.020>.
- Sulaiman, Rini, R Baker, Budi Susilorini, Kevin H Telmer, and Samuel Spiegel. 2007. "Indonesia Country Report." [???? Bibliography?](#)
- Telmer, Kevin H., and Marcello M. Veiga. 2009. "World Emissions of Mercury from Artisanal and Small Scale Gold Mining." In *Mercury Fate and Transport in the Global Atmosphere Emissions, Measurements and Models*, edited by N. Pirrone and R. Mason, 131–72. New York: Springer. <https://doi.org/10.1007/978-0-387-93958-2>.
- UNEP ([United Nations Environment Programme](#)). 2008. "The Global Atmospheric Mercury Assessment: Sources, Emissions and Transport." UNEP-Chemicals, Geneva. <https://www.unenvironment.org/resources/report/technical-background-report-global-atmospheric-mercury-assessment-2008-0>.
- Veiga, Marcello M., Peter A. Maxson, and Lars D. Hylander. 2006. "Origin and Consumption of Mercury in Small-Scale Gold Mining." *Journal of Cleaner Production* 14(3-4): 436–47. <https://doi.org/10.1016/j.jclepro.2004.08.010>.
- Veiga, Marcello M., Denise Nunes, Bern Klein, Janis A. Shandro, P. Colon Velasquez, and Rodolfo N. Sousa. 2009. "Mill Leaching: A Viable Substitute for Mercury Amalgamation in the Artisanal Gold Mining Sector?" *Journal of Cleaner Production* 17-(15): 1373–81. <https://doi.org/https://doi.org/10.1016/j.jclepro.2009.03.012>.
- Verbrugge, Boris, and Sara Geenen. 2019. "The Gold Commodity Frontier: A Fresh Perspective on Change and Diversity in the Global Gold Mining Economy." *Extractive Industries and Society* 6 (2): 413–23. <https://doi.org/10.1016/j.exis.2018.10.014>.
- Vogt, Craig. 2012. "International Assessment of Marine and Riverine Disposal of Mine Tailings." In *Proceedings of the 34th Meeting of the London Convention and the 7th Meeting of the London Protocol*, London, UK. Vol. 1.

**Published**





Yayu Arifin <yayu\_arifin@ung.ac.id>

## Toxicological & Environmental Chemistry - Decision on Manuscript ID GTEC-2019-0447.R1

**Toxicological & Environmental Chemistry** <onbehalfof@manuscriptcentral.com>

6 September 2020 05:10

Balas Ke: hartmut.frank@uni-bayreuth.de

Kepada: yayu\_arifin@ung.ac.id

Cc: toxenchem@uni-bayreuth.de

05-Sep-2020

Dear Dr Arifin:

Your revised manuscript entitled "Artisanal and Small-Scale Gold Mining Activities and Its Mercury Exposures in Gorontalo Utara Regency, Indonesia" which you submitted to Toxicological & Environmental Chemistry, has been checked for compliance with the Instructions for Authors, the reference style guide, and general grammar/style - see some examples in the attached annotated file (not comprehensive, just a few examples).

There are a considerable number of incompatibilities and flaws, but subject to further, more careful revisions (especially of your references), your paper could be suitable for publication. Please consider these suggestions, and I look forward to receiving your revision.

When you revise your manuscript please highlight the changes you make in the manuscript by using the track changes mode in MS Word or by using bold or coloured text. We also need a "clean" copy, i.e. one without correction marks.

At this stage it is particularly important that you meticulously check the correctness of your manuscript in respect to spelling, punctuation, grammar and style, compatibility with the Information for Authors

(<http://www.tandfonline.com/action/authorSubmission?journalCode=gtec20&page=instructions#.VrCdTrLRHfA>) and especially the reference style guide

([http://www.tandf.co.uk/journals/authors/style/reference/tf\\_ChicagoAD.pdf](http://www.tandf.co.uk/journals/authors/style/reference/tf_ChicagoAD.pdf)). Non-compliance with these general rules will result in delays of the publication of your article, as the manuscript will be returned for further corrections until it is largely flawless; it is exclusively the authors' duty to supply a manuscript which is free of errors or incompatibilities with the style of T&EC.

To submit your revision, log into <https://mc.manuscriptcentral.com/gtec> and enter your Author Centre, where you will find your manuscript title listed under "Manuscripts with Decisions." Under "Actions," click on "Create a Revision." Your manuscript number has been appended to denote a revision. Please enter your responses to the reviewers' comments in the space provided. You can use this space to

document any changes you made to the original manuscript. Please be as specific as possible in your response.

Alternatively, once you have revised your paper, it can be resubmitted to Toxicological & Environmental Chemistry by way of the following link:

\*\*\* PLEASE NOTE: This is a two-step process. After clicking on the link, you will be directed to a webpage to confirm. \*\*\*

[https://mc.manuscriptcentral.com/gtec?URL\\_MASK=97a47694ed804fedada4e489fc6f9748](https://mc.manuscriptcentral.com/gtec?URL_MASK=97a47694ed804fedada4e489fc6f9748)

IMPORTANT: Your original files are available to you when you upload your revised manuscript. Please delete any redundant files before completing the submission.

Thank you for submitting your manuscript to Toxicological & Environmental Chemistry, and I look forward to receiving your revision.

<https://mail.google.com/mail/u/0/?ik=617d61bd3f&view=pt&search=all&permmsgid=msg-f%3A1677029354406058657&simpl=msg-f%3A1677029354406058657> 1/2 12/9/22, 8:42 AM Email Universitas Negeri Gorontalo - Toxicological & Environmental Chemistry - Decision on Manuscript ID GTEC-2019-0447.R1

Sincerely,  
Dr Frank  
Editor in Chief,  
Toxicological &  
Environmental  
Chemistry  
[hartmut.frank@un  
i-bayreuth.de](mailto:hartmut.frank@uni-bayreuth.de)







**2020Yayu-getc-main-document.docx**

108K <https://mail.google.com/mail/u/0/?ik=617d61bd3f&view=pt&search=all&permmsgid=msg-f%3A1677029354406058657&simpl=msg-f%3A1677029354406058657> 2/2



## Artisanal and small-scale gold mining activities and mercury exposure in Gorontalo Utara Regency, Indonesia

Yayu Indriati Arifin<sup>a,b</sup> , Masayuki Sakakibara<sup>b,c</sup> , Sayaka Takakura<sup>d</sup>, Mohamad Jahja<sup>b,e</sup> , Fitriyane Lihawa<sup>f</sup> and Koichiro Sera<sup>g</sup> 

<sup>a</sup> Department of Geology, State University of Gorontalo, Kota Gorontalo, Indonesia; <sup>b</sup> The Research Institute for Humanity and Nature (RIHN), Kita-ku, Kyoto, Japan; <sup>c</sup> Graduate School of Sciences and Engineering, Ehime University, Matsuyama, Japan; <sup>d</sup> Tsuneshi Shipbuilding Co., Ltd. Fukuyama, Japan; <sup>e</sup> Department of Physics, State University of Gorontalo, Kota Gorontalo, Indonesia; <sup>f</sup> Department of Educational Geography, University of Gorontalo, Kota Gorontalo, Indonesia; <sup>g</sup> Cyclotron Research Center, Iwate Medical University, Takizawa, Japan

### ABSTRACT

Small-scale gold mining activities and their mercury emissions in the Utara Regency of the Province, Sulawesi Island, Indonesia, have been studied. The mining occurs in many areas of this region, but especially in the three villages Hulawa, Ilangata and Ilangata Barat. Mining activities in the Hulawa village are the oldest (19th century) in the regency, whereas mining in the Ilangata village began just 15 years ago. The activity level changes rapidly from time to time depending on the fluctuations of the gold market. Women and children are involved in the gold mining process. Female workers crush ore, process sediment wastes, and pan for gold in rivers. Children assist their mothers in these activities and play around the mining sites. Gold mining in the Utara Regency is estimated to produce about 290 kg of gold and emits approximately 860 kg of mercury into the environment annually. The impact of such large mercury release into the environment have caused environmental degradation and contamination of fish as food for humans. Collaborative works of stakeholders are urgently needed to find solutions to the problems related to small-scale gold mining such as marginalization and environmental pollution.

### ARTICLE HISTORY

Received 17 October 2019

Accepted 10 October 2020

### KEYWORDS

Artisanal and small-scale gold mining; mercury exposure; human health problems; Gorontalo Utara; Indonesia

### 1. Introduction

*Indonesia is the world's second-largest source of mercury emissions from artisanal and small-scale gold mining (ASGM) after China (Veiga,*

**CONTACT** Yayu Indriati Arifin [yayu\\_arifin@ung.ac.id](mailto:yayu_arifin@ung.ac.id) Department of Geological Engineering, State University of Gorontalo, Gorontalo 96128, Indonesia

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial/NoDerivatives License

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.




Figure 1. The top ten of mercury emitter countries from ASGM in the World.

Maxson, and Hylander 2006; Pacyna et al. 2010; Ismawati 2014). The top 10 global ASGM mercury emitters in 2000 and 2005 are shown in Figure 1 (Veiga, Maxson, and Hylander 2006; Telmer and Veiga 2009; Pacyna et al. 2010; Cordy et al. 2011; Ismawati 2014). Annual mercury emissions from ASGM in Indonesia in 2000, 2005, and 2013 were approximately 125, 150, and 175 tons, respectively. In a five-year period (from 2000 to 2005), Indonesia's ASGM mercury emissions were increased by approximately 20%.

According to UNEP (2008), there are 115 ASGM industries distributed evenly throughout Indonesia that collectively produce 60 tons of gold per annum. The number of ASGM sites in Indonesia is estimated to be ca. 900 involving about 250,000 miners (Purwana 2013). The largest gold mining site in Indonesia is located near Puncak Jaya in Papua. Tailings from this site are dumped in Timika in eastern Indonesia (Nakagawa 2008; Vogt 2012). Here ASGM activities can be found in several places, for example, Jayapura, Nabire, Paniai and Manokwari. ASGM activities also occur in Maluku Province, Gunung Botak and Gogorea on Buru Island, Maluku Province (Male et al. 2013; Idrus et al. 2014).

Several ASGM locations have also been reported in Sulawesi, including Poboya in Palu (Basir-Cyio et al. 2012). According to Ismawati, Petrlik, and DiGangi (2013), ASGM miners' activities in Poboya use 100 tons of mercury annually (Nakazawa et al. 2016). Basir-Cyio et al. (2012) estimate that about 34 tons of mercury are used daily and produce 22.4–45 kg gold daily. High concentrations above 1000 ng m<sup>-3</sup> the limit of air-quality guideline for annual mercury exposure by the World Health Organization) of mercury vapor from ASGM were detected in Palu City, and almost all residents of the city are exposed (Nakazawa et al. 2016). Several ASGM has been reported in Mamuju of Sulawesi Barat Province, Tana Toraja and Luwu of Sulawesi Selatan Province and the Bombana of Sulawesi Tenggara Province. The Bombana ASGM has been reported to have negative impact on social, economic and cultural status of the people around the ASGM (Rianse et al. 2015).

There are four other ASGM locations in Sulawesi Utara Province (Basir-Cyio et al. 2012), at Talawaan (Filho and Villas-Boas 2004; Bose O'Reilly et al. 2010), Buyat (Kambey, Farrell, and Bendell-Young 2001; Limbong et al. 2005), Tobongon and Tanoyan in the Bolaang Mongondow Regency. Mercury emissions from Talawaan alone are 70 t per annum (Lasut et al. 2010), which is seven times the value reported earlier by Telmer and Veiga (2009).

*At least there is one ASGM site in each regency of the Gorontalo Province: (1) Pohuwato Regency: Gunung Pani and Bulontio; (2) Boalemo Regency: Bilato; (3) Gorontalo Regency: Bumela (Lihawa and Mahmud 2019) (4) Bonebolango Regency: Tulabolo and Mopuya; and (5) Gorontalo Utara Regency: Hulawa and Ilangata villages.*


*Gold deposits in the Gorontalo Utara Regency have existed in Wubudubreccias within Boliyohuto diorite of the middle Miocene age (Carlile, Digdowirogo, and Darius 1990). Gold in hydrothermal breccias and quartz veinlets has been reported in Sumalata and Boliyohuto subdistricts (Carlile, Digdowirogo, and Darius 1990). Gold mining in the Hulawa village is considered to date back to Indonesia Dutch era (Henley 2005).*

*This study describes mercury pollution from ASGM sites in the Gorontalo Utara Regency where gold mining activity is intense, but information is scarce. The report describes how miners work and what the mercury emission situation is, as a basis for further discourse on health risks in and around these mining sites.*

## **2. Materials and methods**

### **2.1. Study area**

*The Hulawa (meaning ‘gold’ in the Gorontalo language) mining area (Figure 2) is also known as the Buladu mining area in Sumalata. This area lies within the Buladu River watershed, between the Pasolo and Padengo villages, approximately 109 km northeast of Gorontalo City, the capital of Gorontalo Province. From Gorontalo City, the area can be reached by car in approximately four hours. Buladu has a typical equatorial climate with two seasons (rainy: November–April; dry: May–October), with the highest rainfall occurring in December (annual average 1,210 mm). The temperature is 27 C on average. The Buladu*



**Figure 2.** Map of the Gorontalo Utara Regency showing the location of Sumalata sub district, where the Hulawa mining area is, and the Anggrek sub district where the Ilangata and Ilangata Barat mining areas are located.

River watershed drains from the mountain peaks with the highest point being 2,000 m, on the northern coast of the Sumalata district. The distance from the mountain peaks to the Sulawesi Sea is approximately 10 km. Several rivers run through this watershed, including the Buladu and Wubudu rivers.

The Ilangata and Ilangata Barat mining areas (Figure 2) comprise the Anggrek mining area. The area lies within the Anggrek River watershed between the Diata, Lunggulo, and Lantolo villages, which are approximately 68 km northwest of Gorontalo City. It is three hours by car from Gorontalo City. The Anggrek River watershed drains from the peaks of several hills with elevations of 300 m at most. The distance from the peaks to the sea is approximately 5 km. Number of rivers, including the Anggrek and Ilangata rivers, flow through the Anggrek watershed.

Residents of Hulawa, Ilangata and Ilangata Barat are exposed directly and indirectly to mercury from mercury vapor of ASGM process and sea fish consumption, while residents of other subdistricts (Kwandang, Monano and Tolinggula) are treated as control group since they might only be exposed to mercury through sea fish consumption.

## 2.2. Sample and data collection

The ASGM locations within Gorontalo Utara regency have been visited many times over the course of one month each year, from 2012 to 2014. Data on mercury use during ASGM and the activities of miners and their families were obtained via direct interviews with miners, mine owners, and local government officials. These interviews were conducted onsite at the mining locations in the Gorontalo Utara Regency (Figure 2). The simple questionnaire consisted of number of workers, number of gold trommels, amount of ore filled into trommels, and amount of mercury consumed and gold produced in a single mill process. This was done in Hulawa, Ilangata and Ilangata Barat ASGM sites (n = 23): 6 ASGM operators, 2 panners, 3 villagers in Hulawa, 6 ASGM operators, 4 villagers and 2 gold shop workers in Ilangata and Ilangata Barat, 2 officers in Kwandang). The interviewed were key-actors, including mine site managers, head of villages, and elderly people who were working as ASGM miners 20 to 40 years ago.

Data obtained from different sources were compared with each other via triangulation procedure, for example, number of miners from the owner, from direct investigation and from local officers were confronted other to keep accountable. Any information obtained from participants was recorded and is used in triangulation process. For instance, we recorded the following information while communicating with villagers who have a motorcycle repair kiosk in Ilangata, his friend said that: "he started that business after he earned about 6000 US-\$ for a week from ASGM one year ago". Once we were in Hulawa village, a man who got training on the Minamata disease problem informed us that in nearby villages live three people who were suspected to have tremor disturbance related to their former activities as ASGM miners in Hulawa. The first person showed tremor during our conversation and he has a problem on oral communication already. The second one, also showing tremor with a lesser degree, said that: 'We were working as ASGM miners, dealing with mercury without safety equipment during smelting; sometimes they put gold amalgam in their mouths to sell later on'. They also consumed fish from contaminated rivers, the estuary, and the sea.

Information on the economy and social life in ASGM were also obtained from interviews with miners, managers of ASGM businesses, and villagers. Compared to large-scale mining, ASGM activities have more positive economic impacts on local communities, since the capital is shared within local investors, miners, and villagers (Langston et al. 2015). Economy and social life information, in this case, will enrich our understanding on ASGM activities, its impacts on the environment and human health, and ongoing struggles against social marginalization in Indonesia's gold mining sector (Spiegel and Veiga 2010).

## 2.3. Analytical procedure

The health effects on ASGM miners and villagers were examined by collecting scalp hair strands from 115 donors from five subdistricts. The hair strand from the occipital region was cut close to the skull, collected, put in a clear plastic bag, and labeled accordingly. Mercury

content in hair was determined using proton induced X-ray emission (PIXE) at Iwate Medical University, Japan, as reported elsewhere (Sera, Futatsugawa, and Murao 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera 2017). The procedure is known as a standard-free method for hair without powdering and acid treatment as reported in previous studies (Sera, Futatsugawa, and Matsuda 1999; Sera, Futatsugawa, and Murao 2002; Arifin, Sakakibara, and Sera 2015; Basri, Sakakibara, and Sera 2017). The widely accepted exposure limit values for mercury concentration in hair (lg/g) as suggested by the German Human Biomonitoring Commission (HBM) as the following categories are adopted: concentration (mg/g) below 1.0 are considered normal, 1.0–5.0 are an alert level, and above 5.0 are indicative of a serious risk of health effects.

#### 2.4. Neurological assessment

The participants with elevated hair mercury concentration were assessed clinically according to a questionnaire specifically for subjective symptoms and a neurological examination, and the use of mercury-containing soap (for female only). Neurological examination was performed by a medical doctor on participants, using a UNIDO protocol which consists of 10 symptoms: (1) rigidity and ataxia (walking and standing); (2) tremor: tongue, eyelids, finger to nose, pouring, posture holding, and the Romberg test; (3) signs of bluish discoloration of gums; (4) irregular eye movements or nystagmus; (5) salivation and dysarthria; (6) field of vision; (7) alternating movements or dysdiadochokinesia; (8) knee jerk reflex and biceps reflex; (9) sensory examination and (10) Babinski reflex and labial reflex. Score 1 was given for positively-observed symptoms, and 0 for negatively-observed symptoms. The sum of score for every participant is calculated and analyzed.

#### 2.5. Description of sites and processes in of ASGM

Ore is manually excavated from nearby hills or mountains (Figure 3(a)) using vertical holes in the ground (Figure 3(b)). Traditional tools such as broad hoes, bars and simple pulleys are used for the ore excavation (Figure 3(c)). The ore is manually crushed (Figure 3(d)) using hammers or other percussion tools (Figure 3(e)). Some processing plants use homemade mechanical crushers. The ore is then packed into sacks and manually transported to the processing plant (Figure 3(f)).

At the processing plant, crushed ore (30–40 kg) is fed into a trommel (known as ‘Tromol’ in the local language (James 1994), a steel mill grinder where gold is extracted from ore (Male et al. 2013) (Figure 4(a)) with water (Figure 4(b)) and several decimeter-sized rocks (Figure 4(c)) and milled for 3–4 hours until the material is fine enough to release the gold. Subsequently, approximately 500 g of mercury are added to the trommel (Figure 4(d)), which then rotates for 30 minutes until the amalgamation has completed. This amalgamation process is the major gold production method for ASGM miners in Gorontalo Province.

After 30 minutes, the material is discharged (Figure 4(e)), and the tailings are manually separated using bare hands (Figure 4(f)). The tailings are washed and rinsed several times (Figure 4(g–j)) to separate the gold from the unamalgamated mercury. The mixture is filtered using a fine cloth to separate the mercury amalgam (Figure 4(k)). The gold amalgam is recovered from this process as shown in Figure 4(l).

After milling the gold, the amalgam is recovered, and the waste sediment is kept (Figure 4(m)) for a second recovery process. Then, mercury-contaminated water is released into the environment (Figure 4(n)). The amalgams are then smelted to obtain gold bullion. Gold smelters with simple tools smelt the gold outdoors (Figure 4(o)) or inside houses



**Figure 3.** ASGM in the Gorontalo Utara Regency: (a) An artisanal mine shaft on a slope. (b) A hole where the miners are working. (c) A sack of ore is being pulled from the base of the hole. (d) The ore is manually crushed using simple tools. (e) The percussion tools used to crush the ore. (f) Raw crushed ore is manually transported to the processing plant.


and gold shops. Gold bullion is recovered from the smelting process

(Figure 4(p)).

The secondary process begins when miners, usually women, and children, wash sediment wastes from the primary milling process in running water. The milling process is then repeated, as shown in Figure 4(a) through Figure 4(l), with the difference being that the trommel is generally rotated by a wheel powered with river water (Figure 4(n)).

Tertiary processing involves bagging the waste from the secondary processing and sending it to another processing plant to undergo cyanidation. There is no cyanidation plant in the Gorontalo Utara Regency.





**Figure 4.** Artisanal and Small-scale mining (ASGM) activities in three locations in the Gorontalo Utara Regency. (a) Crushed ore is inserted into the ball mill (b) Water is poured into the trommel (c) The softball size rocks used to mill the ore inside the trommel (d) Where mercury is poured into the trommel mill (e) Discharging the trommel. (f) Separating

**Figure 4. (Continued)**

the gold amalgam from tailing (g–j) Extracting the gold-containing mercury (k) Separating amalgam from mercury (l) The amalgam is recovered (m) Tailing waste is kept into wooden boxes for secondary processing (n) Wastewater flows into the rivers stream directly (o) Smelting the amalgam (p) Gold bullion (q) A woman panning the sediment for gold r: Extracting gold from sediment using mercury.

**Table 1.** Number of mining workers, population, trommel mills and mercury needed per milling process at several locations in the Gorontalo Utara Regency.

Subdistrict	Village	Number of Workers	Population	Number of Trommel Mills	Mercury/Milling Process (kg)
Ilangata Barat	Diata	10	382	35	23
	Lunggulo	6	247	43	23
Ilangata	Lantolo	8	248	60	71
Hulawa	Pasolo	29	558	164	105
	Padengo	14	0	76	54
Total	–	67	1435	378	275

However, there is a cyanidation plant in the other Regency of Gorontalo Province. Numerous miners pan in the Wubudu River estuary (Figure 4(q)) and (Figure 4(r)) where the mercury enters the environment (air, soil, and water) directly.

Smelting activities occur in gold shops in the Ilangata Barat village. The owners of these gold shops determine the gold quality (amount of gold in ‘dore’) via these processes. The amount of gold in a dore varies from 40–80% of the weight. A dore with 80% quality can be purchased for approximately US \$30,-/gram. Mining wastes have solid and liquid forms that contaminate river waters and soils. Gold bullion is extracted after smelting the amalgam, which emits mercury vapor.

### 3. Results

#### 3.1. Mercury emissions

Because ASGM activities are unregistered, no exact data were available on the number of workers; therefore, it is difficult to estimate the amount of gold produced, and mercury used, for ASGM activities throughout Indonesia and its local provinces or regencies. This difficulty in estimating amounts of gold produced and mercury used by ASGM throughout the world is described elsewhere (Seccatore et al. 2014). We used direct observation and interviews to get information and compared it with data reported by several researchers. Data on mercury imported to Indonesia will be used as the upper limit of mercury lost to ASGM. Socio-economic factors which should be considered when dealing with ASGM (Sulaiman et al. 2007; Spiegel and Veiga 2010; Spiegel 2012; Langston et al. 2015), will be elaborated further to discuss any possible scenario or solutions on sustainable development in local ASGM site areas.

Data on the number of mine workers, population and mercury used during the milling process in each village are listed in Table 1. These figures were estimated based on each milling process requiring 0.5 kg mercury for one trommel mill to extract gold from ca. 30–40 kg of crushed ore. Assuming all trommel mills are operating during the milling process, a maximum of 275 kg of mercury is needed (these data and the amount of mercury emitted per gram of gold produced allows us to estimate the mercury emitted into the environment). Each primary process produced ca. 3 g of gold per trommel mill. The secondary processing of waste produces ca. 0.5 g of gold per trommel mill. We also assume the miners only work five days a week and sell the gold on Saturday. The ratio of mercury emissions to gold produced is ca. 2:1 (Darmutji 2003). The amount of gold produced by the process is ca. 1134 g. Thus, the mercury emissions are ca. 2.2 kg per milling process. The total annual mercury emission

in the Gorontalo Utara Regency is approximately 572 kg. If we used 1-3 grams of mercury lost per gram of gold produced, the mercury emission ranges from 286858 kg (Veiga et al. 2009).

### 3.2. Mercury concentration in hair samples according to geographic location

The distributions and range of mercury levels in 115 hair samples collected from the five subdistricts are shown in Figure 5. Mercury concentrations in all hair donors are more than 1 lg/g, which indicates the




Figure 5. Mercury in hair data among the population of Gorontalo Utara regency as box plot, two threshold lines (HBM I and HBM II) according to German Human Biomonitoring (HBM) Commission are shown, number of sample of each group is presented in the bracket.




Figure 6. The hair mercury levels among females and males are lognormally distributed. Solid and open bar indicate female and male, respectively. Geometric mean and Standard Deviation (SD) are shown for female, male and total.

concentrations are already within the alert level range according to HBM (Schulz et al. 2007). The number of subjects with high mercury levels over 10 lg/g was 10 (40%), 7(30%), and 4 (8.5%) in Anggrek, Sumalata, and the control group (Kwandang, Monano and Tolinggula), respectively.

According to the Kolmogorov-Smirnov test, the distribution of mercury in the hair data from the Gorontalo Utara Regency was not normal; instead, it had a log normal distribution

(Figure 6). The geometric mean is more suitable for log normal distribution data. Mean mercury level ( $\mu$ / standard deviation) of Gorontalo Utara regency is 6.7 (2.0) lg/g and is higher than minimum limit for danger level (5.0 lg/g) according to HBM. Only residents of Tolinggula has average hair mercury level less than 5.0 lg/g. This may due to their residence away from the seaside; their common diet along with rice is chicken. Even though one male and two females have mercury higher than 6.0 lg/g, the Tolinggula group could be considered as a control group in Gorontalo Utara regency.

The data were plotted using a box-and-whisker plot to make a clear view of data distribution in each residence and among Gorontalo Utara regency data. Each Box is filled with the horizontal line representing the position of the median. The position of the median and mean relative to HBM was used to put the population into three categories according to HBM. Box plot of mercury distribution according to geographical location is shown in Figure 5, the results of Sumalata and Anggrek are positioned higher than others, which means that their hair average mercury is elevated in the regency. Other subdistricts (Tolinggula and Kwandang) have no known ASGM activities, but the population lives close to seashore (their common diet is fish from Sulawesi sea, same as the population of Anggrek and Sumalata).

### 3.3. Mercury concentration for male and female

The hair mercury levels among females and males are lognormally distributed is shown in Figure 6. Three females had mercury levels greater than 50 lg/g, and none of them worked as ASGM miners. Those three females may have been exposed to mercury from another source. The elevated hair mercury levels that were above average (6.8 lg/g) and even the highest (17.9 lg/g) mercury level were found among the ASGM miners. The average hair mercury levels for all, male and female inhabitants in the Kwandang subdistrict are 6.5, 6.1 and 6.9 lg/g, respectively. These levels show that there are no significant differences between the mean hair mercury of males and females in that subdistrict. While the average of all mercury hair content for females is 7.4 lg/g (only more than 11 percent higher than males (6.1 lg/g), such conditions were also found for the subgroups of Monano and Sumalata. The condition in Anggrek and Sumalata are even higher (the average mercury for females is 50 percent higher than the males). The large discrepancy of mercury levels between female and male inhabitants suggests that female inhabitants may receiving mercury from another source (e.g. whitening cream, soap, etc).

The clinical symptoms were observed in most of the females with mercury higher than 50 mg/g (the minimum level at which Minamata disease symptoms are seen (Harada 1995)) are listed in Table 2, indicative of inorganic-mercury poisoning only. All the subjects have habitual use of whitening cream that are common in Gorontalo Utara regency. The mercury concentration on the whitening creams are shown in Table 3. It seems that elevated mercury concentration among female inhabitants of


Table 2. Females with hair mercury level more than 50 mg/g.

Subdistricts	Age (years)	Mercury level (mg/g)	Duration of use	Clinical symptoms	Occupation	Notes on ASGM activities
llangata	45	144.8	10 years	Headache, tremor	Housewife	not active

	13	61.3	2 years	Headache	Student	not active
Hulawa	36	69.8	8 years	Headache, tremor	Housewife and owner of local ASGM industry	Burning Amalgam at home

**Table 3.** Mercury concentration in common whitening cream used among female in the ASGM.

Whitening cream	Mercury level (mg/g)
A	11.9
B	7.2
C	4.1
D	3.3



**Figure 7.** Flowchart showing artisanal mining operations to extract gold using mercury and mercury contamination pathways to humans and the environment.

Hulawa and Ilangata may caused by direct mercury contact from habitual use of whitening creams. Harada et al. (2001) found the similar cases in Kenya which were caused by female were elevated hair mercury levels were using soaps with high mercury concentrations.

## 4. Discussion

### 4.1. ASGM operation system and mercury losses

Figure 7 shows the ASGM operation system which is used in Gorontalo Utara Regency; the system used in North Gorontalo is similar to those reported elsewhere (Veiga et al. 2009). Gold recovery systems is based on whole ore amalgamation using trommel, gravitation and panning methods were found in Wobudu river estuary and Anggrek river banks. There are two possible routes for mercury contamination from ASGM into the environment and humans. During the milling process, liquid mercury could contaminate miners and the environment. Working with mining wastes (tailings), and direct exposure to liquid mercury during panning, are also contamination routes. Mercury vapor could contaminate the smelters and environment during smelting.

For comparison, we also estimated the total annual mercury emissions in the Gorontalo Utara Regency from ASGM using data on Indonesian mercury emissions (Pacyna et al. 2010) and the number of known ASGM sites (Kocman et al. 2013). The rate for Indonesian mercury emissions from ASGM may be close to 200 t annually. The average emission from each mining site is approximately 200kg/year based on the 900 known ASGM sites (Kocman et al.

2013). The three mining sites in the Gorontalo Utara Regency emit approximately 600kg/year. Ismawati (2014) estimated the ASGM production ranged from 65–130 t in 2012 (Ismawati 2014). Thus, approximately 170260 t of mercury was lost to the environment based on the multiplication factor from Darmutji (2003). Using the 1:1 to 3:1 ratio proposed by Ismawati (2014), the amount of mercury lost was 65390 t. These estimates agree with the total mercury import data for Indonesia in 2012, which was approximately 368 t.

#### 4.2. Health issues

The involvement of women and children workers is unique to the Hulawa mining site. These women and children come from the Buladu area. They work with waste stored in wooden boxes, and sluice and pan in the Wubudu River estuary. Women and children typically work on mercurybearing mining wastes during the secondary processing stage (Figure 8(a,b)). Their bare hands and feet are directly exposed to the mining waste, which allows mercury to enter their body via dermal absorption.

Health problems in miners and inhabitants were documented by medical doctors using a standard questionnaire and neurological test tools (Figure 9(a,b)). Ten neurological disorder indicators/symptoms were observed. Bluish gums and tremors are major symptoms observed in the




Figure 8. The wastewater discharge goes directly into the river. (a) A small child working with her mother handling tailings (solid waste) at the water-powered mill. (b) A woman extracting gold from sediment using mercury.



Figure 9. A medical doctor observing the inhabitants' symptoms during a neurological examination. (a) Observation of tremor symptoms. (b) Observation of bluish gum symptoms.



miners and inhabitants of the Gorontalo Utara Regency. Several explanations could be addressed based on the evidence; first, they have experienced prolonged contact with mercury, and second, the mercury has entered their body via the food chain (Bose-O'Reilly et al. 2010; BasirCyio et al. 2012; Nakazawa et al. 2016). Both can happen simultaneously for the miners and inhabitants of these mining sites. Further investigations are needed to explain the evidence and determine the most probable mercury contamination routes in the Gorontalo Utara Regency.

Without pollution controls and environmental cleanup, further mercury contamination of the environment (bottom sediments, water, and fish) and human population around ASGM sites within the Gorontalo Utara Regency is inevitable (Arifin, Sakakibara, and Sera 2015). Veiga and Hinton (Seccatore et al. 2014) reported that gold bullion contains two to five percent mercury just after smelting, and such gold bullion is refined at gold shops in towns within the Gorontalo Utara Regency with few emission controls. Mercury pollution from gold shops, like those observed in Colombian cities, is of great concern (Cordy et al. 2013).

#### 4.3. Socio-economic aspects of ASGM system in gorontalo utara regency

The intensification of ASGM activities at the historical gold mining site (hulawavillage) and expansion to the new gold mining site (Ilangata and Ilangata Barat villages) has been seen as advantage of ASGM over large scale gold mining (LSGM) (Adler Miserendino et al. 2013). The ability of ASGM activities to cut down production cost through their reliance on flexible informal labor. The ASGM activities are driven by local investors which are intimately entangled with socio-political structures and land tenure systems (Verbrugge and Geenen 2019).

The hierarchy of people working in ASGM system in Gorontalo Utara regency can be explained using the network chart introduced by Langston et al. (2015), with slight modification mainly related to the number of ASGM miners. In general, a small number of miners (up to 3 to 4 people) work at the excavating site or processing site; these miners may be family members or friends. Hence supervisor, landowner position, and even investor itself can be the same person.

Local miners are formerly farmers or fisherman and live nearby in their houses, while migrants usually live inside the processing site camps, which are very simple structures and lack sanitation facilities. The basic needs (foods, drinks, cigarettes) were available in kiosk/gold shops which may belong to cooperation or investor/landowner/supervisor.

Positive economic impacts of ASGM activities to miners and inhabitants were not easily seen in Gorontalo Utara regency since there is no significant difference in economic activities in ASGM sites and non ASGM sites. In a stakeholder meeting, ASGM miners, local government, and investor seem to have opposing viewpoints on issues such as sustainable development, environmental protection, and negative impacts of ASGM on health.

Local governments and inhabitants would like to embrace long-term economic solution such as farming, fishery, and tourism, which could eventually replace ASGM. There is ongoing research on finding the ultimate solutions which fit to Gorontalo Utara Regency cases. The multidisciplinary research approach is being taken to help young local students who were

living in the ASGM area to find solutions to their local problems. A transdisciplinary approach could facilitate the stakeholders in ASGM area together with the researcher to develop the ideas on sustainable development (Sippl and Selin 2012).

## 5. Conclusions

ASGM activities in the Gorontalo Utara Regency are dangerous to the environment, miners, and the local population. The amount of mercury used per mining cycle is approximately 275 kg, and approximately 2.2 kg of mercury is lost per cycle. Annually, ASGM in the Gorontalo Utara Regency produces approximately 286 kg of gold and emits approximately 572 kg of mercury. This estimate is probably conservative given the limited mining data available. The impact of such a large mercury loss into the environment and human population has been to degrade the environmental conditions and contaminating humans and fish as demonstrated by Arifin, Sakakibara, and Sera (2015).

Gorontalo Utara Regency is an essential agricultural (rice) and horticultural (dairy) production area in the Gorontalo Province and Indonesia. The ASGM driven economy did not bring sustainable development in the vicinity of ASGM sites and Gorontalo Utara regency as a whole. Solutions to the problem are being developed using ongoing multi and trans-disciplinary research approach through the collaboration of Indonesian and Japanese researchers.

## Acknowledgments

The authors thank the regent of Gorontalo Utara and its officers, and the owners of mining plants for allowing us to conduct interviews and take photographs.

## Author contributions

All authors contributed to the work presented in the manuscript.

## Disclosure statement

The authors declare no conflict of interest.

## Funding

This research was supported by The Research Institute for Humanity and Nature (RIHN: a constituent member of NIHU), Project No.14200102.

## ORCID

Yayu Indriati Arifin  <http://orcid.org/0000-0002-4892-6270>  
Masayuki Sakakibara  <http://orcid.org/0000-0002-8188-6979> Mohamad Jahja  
<http://orcid.org/0000-0001-9601-7269>

## References



- Adler Miserendino, Rebecca, Bridget A. Bergquist, Sara E. Adler, Jean Remy Davee Guimar~aes, Peter S. J. Lees, Wilmer Niquen, P. Colon Velasquez-Lopez, and Marcello M. Veiga. 2013. "Challenges to Measuring, Monitoring, and Addressing the Cumulative Impacts of Artisanal and Small-Scale Gold Mining in Ecuador." *Resources Policy* 38 (4): 713–722. doi:[10.1016/j.resourpol.2013.03.007](https://doi.org/10.1016/j.resourpol.2013.03.007).
- Arifin, Yayu, Masayuki Sakakibara, and Koichiro Sera. 2015. "Impacts of Artisanal and Small-Scale Gold Mining (ASGM) on Environment and Human Health of Gorontalo Utara Regency, Gorontalo Province." *Geosciences* 5 (2): 160–176. doi:[10.3390/geosciences5020160](https://doi.org/10.3390/geosciences5020160).
- Basir-Cyio, Muhammad, Uswah Hasanah, Isrun Nur, and Yuka Serikawa. 2012. "Gold Mining Activities and Its Impacts on Land Degradation in Central Sulawesi Indonesia." *Journal of Ecotechnology Research* 16 (3–4): 79–83. doi:[10.11190/jer.16.79](https://doi.org/10.11190/jer.16.79).
- Basri, Masayuki, Sakakibara, and Koichiro Sera. 2017. "Current Mercury Exposure from Artisanal and Small-Scale Gold Mining in Bombana, Southeast Sulawesi, Indonesia—Future Significant Health Risks." *Toxics* 5 (1): 7. doi:[10.3390/toxics5010007](https://doi.org/10.3390/toxics5010007).
- Bose-O'Reilly, Stephan, Gustav Drasch, Christian Beinhoff, Saulo Rodrigues-Filho, Gabriele Roeder, Beate Lettmeier, Alexandra Maydl, Stefan Maydl, and Uwe Siebert. 2010. "Health Assessment of Artisanal Gold Miners in Indonesia." *The Science of the Total Environment* 408 (4): 713–725. doi:[10.1016/j.scitotenv.2009.10.070](https://doi.org/10.1016/j.scitotenv.2009.10.070).
- Carlile, J. C., S. Digidowirogo, and K. Darius. 1990. "Geological Setting, Characteristics and Regional Exploration for Gold in the Volcanic Arcs of North Sulawesi." *Journal of Geochemical Exploration* 35 (1-3): 105–140. doi:[10.1016/0375-6742\(90\)90037-B](https://doi.org/10.1016/0375-6742(90)90037-B).
- Cordy, Paul, Marcello M. Veiga, Ibrahim Salih, Sari Al-Saadi, Stephanie Console, Oseas Garcia, Luis Alberto Mesa, Patricio C. Velasquez-Lopez, and Monika Roeser. 2011. "Mercury Contamination from Artisanal Gold Mining in Antioquia, Colombia: The World's Highest per Capita Mercury Pollution." *Science of the Total Environment* 410–411: 154–160. doi:[10.1016/j.scitotenv.2011.09.006](https://doi.org/10.1016/j.scitotenv.2011.09.006).
- Cordy, Paul, Marcello Veiga, Ben Crawford, Oseas Garcia, Victor Gonzalez, Daniel Moraga, Monika Roeser, and Dennis Wip. 2013. "Characterization, Mapping, and Mitigation of Mercury Vapour Emissions from Artisanal Mining Gold Shops." *Environmental Research* 125: 82–91. doi:[10.1016/j.envres.2012.10.015](https://doi.org/10.1016/j.envres.2012.10.015).
- Darmutji, Selinawati T. 2003. "Report, UNIDO Project No EG/GLO/01/G34, Overview of Project Site in Indonesia Talawaan and Galangan Areas."
- Filho, Saulo Rodrigues, and Roberto Villas-Boas. 2004. "Biogeochemical Assessment of Mercury Pollution in the S~ao Chico Gold Mining Site - Brazilian Amazon Rio De Janeiro Novembro/2004." In *Environmental Geochemistry in Tropical Counties*, 235.
- Harada, Masazumi. 1995. "Minamata Disease: Methylmercury Poisoning in Japan Caused by Environmental Pollution." *Critical Reviews in Toxicology* 25 (1): 1–24. doi: [10.3109/10408449509089885](https://doi.org/10.3109/10408449509089885).
- Harada, Masazumi, Shigeharu Nakachi, Koa Tasaka, Sakae Sakashita, Kazue Muta, Kohichi Yanagida, Rikuo Doi, Takako Kizaki, and Hideki Ohno. 2001. "Wide Use of Skin-Lightening Soap May Cause Mercury Poisoning in Kenya." *The Science of the Total Environment* 269 (1-3): 183–187. doi:[10.1016/S0048-9697\(00\)00812-3](https://doi.org/10.1016/S0048-9697(00)00812-3).
- Henley, David E. F. 2005. *Fertility, Food and Fever: Population, Economy and Environment in North and Central Sulawesi, 1600-1930*. Leiden: Brill.

- Idrus, Arifudin, Sukmandaru Prihatmoko, Hill Gendoet Hartono, Fadlin Idrus, Ernowo Ernowo, Franklin Franklin, Moetamar Moetamar, and Iwan Setiawan. 2014. "Some Key Features and Possible Origin of the Metamorphic Rock-Hosted Gold Mineralization in Buru Island." *Indonesian Journal on Geoscience* 1 (1): 9–19. doi:10.17014/ijog.v1i1.172.
- Ismawati, Yuyun. 2014. "Gold, Mercury and the Next Minamata." *Indonesian Journal of Leadership, Policy and World Affairs*, April-June 2014. [http://www.sr-indonesia.com/in\\_the\\_journal/view/gold-mercury-and-the-next-minamata?pg=all](http://www.sr-indonesia.com/in_the_journal/view/gold-mercury-and-the-next-minamata?pg=all).
- Ismawati, Yuyun, Jundrich Petrlik, and Joe DiGangi. 2013. "Mercury Hotspots in Indonesia ASGM Sites : Poboya and Sekotong in Indonesia." Accessed 30 October 2020. <http://www.ipen.org/hgmonitoring/pdfs/indonesia-report-en.pdf>.
- James, Laurence P. 1994. "The Mercury 'Tromol' Mill: An Innovative Gold Recovery Technique, and a Possible Environmental Concern." *Journal of Geochemical Exploration* 50 (1-3): 493–500. doi:10.1016/0375-6742(94)90038-8.
- Kambey, Joice L., A. P. Farrell, and L. I. Bendell-Young. 2001. "Influence of Illegal Gold Mining on Mercury Levels in Fish of North Sulawesi's Minahasa Peninsula, (Indonesia)." *Environmental Pollution (Barking, Essex : 1987)* 114 (3): 299–302. doi:10.1016/S02697491(01)00163-4.
- Kocman, David, Milena Horvat, Nicola Pirrone, and Sergio Cinnirella. 2013. "Contribution of Contaminated Sites to the Global Mercury Budget." *Environmental Research* 125 (i): 160–170. doi:10.1016/j.envres.2012.12.011.
- Langston, James Douglas, Muhammad I. Lubis, Jeffrey A. Sayer, Chris Margules, Agni Klintuni Boedihartono, and Paul H. G. M. Dirks. 2015. "Comparative Development Benefits from Small and Large Scale Mines in North Sulawesi." *The Extractive Industries and Society* 2 (3): 434–444. doi:10.1016/j.exis.2015.02.007.
- Lasut, Markus T., Yoshiaki Yasuda, Evan N. Edinger, and Jane M. Pangemanan. 2010. "Distribution and Accumulation of Mercury Derived from Gold Mining in Marine Environment and Its Impact on Residents of Buyat Bay, North Sulawesi, Indonesia." *Water, Air, and Soil Pollution* 208 (1-4): 153–164. doi:10.1007/s11270-009-0155-0.
- Lihawa, Fitryane, and Marike Mahmud. 2019. "The Content of Mercury in Sediments around Artisanal Small-Scale Gold Mining (ASGM) Bumela District, Gorontalo Regency, Gorontalo Province, Indonesia." *IOP Conference Series: Earth and Environmental Science* 314 (August): 012016. doi:10.1088/1755-1315/314/1/012016.
- Limbong, D., J. Kumampung, D. Ayhuan, T. Arai, and N. Miyazaki. 2005. "Mercury Pollution Related to Artisanal Gold Mining in North Sulawesi Island, Indonesia." *Bulletin of Environmental Contamination and Toxicology* 75 (5): 989–996. doi:10.1007/s00128-005-0847-0.
- Male, Yusthinus Thobias, Amanda Jean Reichelt-Brushett, Matt Pocock, and Albert Nanlohy. 2013. "Recent Mercury Contamination from Artisanal Gold Mining on Buru Island, Indonesia-potential future risks to environmental health and food safety." *Marine Pollution Bulletin* 77 (1-2): 428–433. doi:10.1016/j.marpolbul.2013.09.011.
- Nakagawa, Junji. 2008. *Freeport's Grasberg/Ertsberg Mine in West Papua, Indonesia. Opportunities and Challenges for Foreign Investment in the APEC Region: Case Studies.*
- Nakazawa, Koyomi, Osamu Nagafuchi, Tomonori Kawakami, Takanobu Inoue, Kuriko Yokota, Yuka Serikawa, Basir Cyio, and Rosana Elvince. 2016. "Human Health Risk Assessment of Mercury Vapor around Artisanal Small-Scale Gold Mining Area, Palu City, Central Sulawesi, Indonesia." *Ecotoxicology and Environmental Safety* 124 (February): 155–162. doi:10.1016/j.ecoenv.2015.09.042.
- Pacyna, E. G., J. M. Pacyna, K. Sundseth, J. Munthe, K. Kindbom, S. Wilson, F. Steenhuisen, and P. Maxson. 2010. "Global Emission of Mercury to the Atmosphere from Anthropogenic Sources in 2005 and Projections to 2020." *Atmospheric Environment* 44 (20): 2487–2499. doi:10.1016/j.atmosenv.2009.06.009.

- Purwana, Rachmadhi, Arbete och H  lsa 2013. "Safety and Health in Mining in Indonesia." In *Occupational Safety and Health in Mining Anthology on the Situation in 16 Mining Countries*, edited by Kaj Elgstrand and Eva Ving  rd, 47(2):42–49. Gothenburg: Vetenskaplig Skriptserie.
- Rianse, Usman, Weka Gusmiarty Abdullah, Abdi La Abdi, Ilma Sarimustaqyma Rianse, Zulfikar La Zulfikar, Wa Kuasa Baka, La Ode Midi, and Weka Widayati. 2015. "The Impact of the Gold Mining on the Social, Economic, and Cultural in the Bombana Distric Southeast Sulawesi Province." *International Journal of Sustainable Tropical Agricultural Sciences* 1 (1): 53–65.
- Schulz, C., J. Angerer, U. Ewers, and M. Kolossa-Gehring. 2007. "The German Human Biomonitoring Commission." *International Journal of Hygiene and Environmental Health* 210 (3-4): 373–382. doi:10.1016/j.ijheh.2007.01.035.
- Seccatore, Jacopo, Marcello Veiga, Chiara Origliasso, Tatiane Marin, and Giorgio De Tomi. 2014. "An Estimation of the Artisanal Small-Scale Production of Gold in the World." *The Science of the Total Environment* 496: 662–668. doi:10.1016/j.scitotenv.2014.05.003.
- Sera, K., S. Futatsugawa, and K. Matsuda. 1999. "Quantitative Analysis of Untreated BioSamples." *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 150 (1-4): 226–233. doi:10.1016/S0168-583X(98)01071-4.
- Sera, K., S. Futatsugawa, and S. Murao. 2002. "Quantitative Analysis of Untreated Hair Samples for Monitoring Human Exposure to Heavy Metals." *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 189 (1-4): 174–179. doi:10.1016/S0168-583X(01)01034-5.
- Sippl, Kristin, and Henrik Selin. 2012. "Global Policy for Local Livelihoods: Phasing out Mercury in Artisanal and Small-Scale Gold Mining." *Environment: Science and Policy for Sustainable Development* 54 (3): 18–29. doi:10.1080/00139157.2012.673452.
- Spiegel, Samuel J. 2012. "Governance Institutions, Resource Rights Regimes, and the Informal Mining Sector: Regulatory Complexities in Indonesia." *World Development* 40 (1): 189–205. doi:10.1016/j.worlddev.2011.05.015.
- Spiegel, Samuel J., and Marcello M. Veiga. 2010. "International Guidelines on Mercury Management in Small-Scale Gold Mining." *Journal of Cleaner Production* 18 (4): 375–385. doi:10.1016/j.jclepro.2009.10.020.
- Sulaiman, Rini, R. Baker, Budi Susilorini, Kevin H. Telme, and Samuel Spiegel. 2007. "Removal of Barriers to Introduction of Cleaner Artisanal Gold Mining and Extraction Technologies: Indonesia Country Project Summary. Report to the United Nations Industrial Development Organization." <http://iwlearn.net/iw-projects/1223/reports/indonesia/indonesia-country-report>.
- Telmer, Kevin H., and Marcello M. Veiga. 2009. "World Emissions of Mercury from Artisanal and Small Scale Gold Mining." In *Mercury Fate and Transport in the Global Atmosphere Emissions, Measurements and Models*, edited by N. Pirrone and R. Mason, 131–172. New York: Springer. doi:10.1007/978-0-387-93958-2.
- UNEP (United Nations Environment Programme) 2008. "The Global Atmospheric Mercury Assessment: Sources, Emissions and Transport." UNEP-Chemicals, Geneva. <https://www.unenvironment.org/resources/report/technical-background-report-globalatmospheric-mercury-assessment-2008-0>.
- Veiga, Marcello M., Denise Nunes, Bern Klein, Janis A. Shandro, P. Colon Velasquez, and Rodolfo N. Sousa. 2009. "Mill Leaching: A Viable Substitute for Mercury Amalgamation in the Artisanal Gold Mining Sector?" *Journal of Cleaner Production* 17 (15): 1373–1381. doi:10.1016/j.jclepro.2009.03.012.
- Veiga, Marcello M., Peter a Maxson, and Lars D. Hylander. 2006. "Origin and Consumption of Mercury in Small-Scale Gold Mining." *Journal of Cleaner Production* 14 (3-4): 436–447. doi:10.1016/j.jclepro.2004.08.010.
- Verbrugge, Boris, and Sara Geenen. 2019. "The Gold Commodity Frontier: A Fresh Perspective on Change and Diversity in the Global Gold Mining Economy." *The*

*Extractive Industries and Society* 6 (2): 413–423. doi:[10.1016/j.exis.2018.10.014](https://doi.org/10.1016/j.exis.2018.10.014).

Vogt, Craig. 2012. "International Assessment of Marine and Riverine Disposal of Mine Tailings." In *Proceedings of the 34th Meeting of the London Convention and the 7th*

*Meeting of the London Protocol*." London, UK. Vol. 1.

Yuyun Ismawaty, Petrlik and DiGangi is deleted

What is the limit? Here the limit must be defined/mentioned concretely.