

Effect Application of Sea Sand, Coconut and Banana Coir on the Growth and Yield of Rice Planted at Ustic Endoaquert Soil

by Nurdin Nurdin

Submission date: 16-Jul-2022 05:36PM (UTC+0700)

Submission ID: 1871182142

File name: Effect_Application_of_Sea_Sand_Coconut_and_BananaCoir.pdf (85.82K)

Word count: 5270

Character count: 26672

Effect Application of Sea Sand, Coconut and Banana Coir on the Growth and Yield of Rice Planted at Ustic Endoaquert Soil

Nurdin

Agrotechnology Department, Faculty of Agriculture, ⁵
Gorontalo State University
Jl. Jend. Sudirman No.6 Kota Gorontalo 96122, Indonesia.
E-mail: nurdin@ung.ac.id

Received 10 October 2013/ Received 18 December 2013

ABSTRACT

The ¹²arch was aimed to study effect application of sea sand (SS), coconut coir (CC) and banana coir (BC) on the growth and yield of rice (*Oryza sativa* L.) planted at Ustic Endoaquert soil. The pot experiment was carried out using a factorial design with 3 factors. The first factor was SS ¹⁷consisted of three levels i.e.: 0%, 25%, and 50%. The second and third factors were CC and BC, each consisted of three levels i.e.: 0, 10, and 20 Mg ha⁻¹. Application of SS and BC significantly increased leaf length where the highest increasing percentage was 16.47% which was achieved at 25% SS application. Their effect on leaf numbers and tiller numbers were relatively not similar pattern where leaf number only increased about 65.52% by BC application, while tiller numbers only increased about 10.77% by SS application. Furthermore, the application of CC and BC significantly increased panicle numbers to 29.53% and 29.05%, respectively compared to control. All ameliorants significantly increased panicle numbers, but the best was CC with the increasing up to 46.49% at 20 Mg ha⁻¹ CC compared to SS or BC application. However, only coconut coir significantly increased the rice grain numbers.

Keywords: Banana coir, coconut coir, rice plant, sea sand, vertisol

INTRODUCTION

Rate of population growth with a percentage of about 2% per year has resulted in increasing demand for rice consumption. Until 2012, national rice demand reached 34.055 million Mg and paddy production reached 68,956,293 Mg (BPS RI 2012) or equivalent to 37 million Mg of rice (Suswono 2012). Sudaryatno *et al.* (2010) predicted that the need of national rice in 2015 as many as 35,123,000 Mg and 37,021,000 Mg in 2020, or an average rate of rice consumption increased 0.92% per year. Viewing statistics in 2012, it was apparently a surplus as much as 3-4 million Mg of rice at the year. Although the current national rice needed was adequate and surplus, but considering Indonesia's populations that were about 247 million and the increasing rate of population growth, the rice production was needed to be maintained and improved, one through the increasing productivity of paddy soil.

Paddy rice field soil is an ecosystem which water source are dominantly from irrigation and as mainly national rice sources which coverage 2.1 million ha (Toha and Pirngadi 2004). In Paguyaman, Gorontalo Province paddy rice field areas are dominantly classified as Vertisols that developed from lacustrine deposition materials (Hikmatullah *et al.* 2002; Prasetyo 2007; Nurdin 2011). From chemically aspect, Vertisol is classified as nutrient rich soil that has high nutrient sources (Deckers *et al.* 2001). However, the physical properties are limiting factor for plant growth which are heavy clay texture, swelling and shrinking properties, lowest water infiltration, and slow drainage (Mukanda and Mapiki 2001). As a result, it frequently has stunted plant growth and low yields. It is necessary to repair these properties by giving ameliorant materials.

Sand is one type of the ameliorant materials that can be applied to high clay soils. Ravina and Magier (1984); Narka and Wiyanti (1999) showed that application of sand had significantly effect to decline COLE value and soil plasticity index, while soil permeability became large, but the water content availability was low. This was in line with statement of Kusnarta (2012) that sand could improve

aggregate and structure stability of Vertisol (*stability quotient*), way of declining clay function mechanism in swelling-shrinking processes. Besides, the application of sand about 20% of weight declined COLE values. The SS was used as planting media (Sari *et al.* 2006), but it had high NaCl contents that caused some crops grew abnormally (Kusnarta *et al.* 2014). Walter *et al.* (2000); Oliver and Smetten (2002); Al-Omran *et al.* (2004) has stated sandy texture soil was very influential on status and distribution of water, so it influenced root system and root depth. Further Rajiman (2009) has stated that the sandy texture soil influenced on soil nutrient and pH too.

However, paddy soil cultivation requires medium permeability with sufficient water content, so it needs another ameliorant to fix these properties, such as by using CC and BC. CC has been used as water storage on farms (Subiyanto *et al.* 2003). While, the banana coir is still relatively not used yet. Water absorption of banana coir was relatively high when it was dried because it had pores that were interconnected (Indrawati 2009).

Applications of the three ameliorant were allegedly able to mutually improve soil physical and chemical properties of Vertisol under rice cultivation in paddy rice field, so that its productivity could be improved. This research was aimed to study the effect application of sea sand, coconut coir and banana coir on the growth and yield of rice planted at Ustic Endoaquert soil taken from paddy rice field.

MATERIALS AND METHODS

Study Site

The experiment was conducted in the green house of Mathematics and Natural Science Faculty of Gorontalo State University starting in April-August 2012. Meanwhile, the growing media was Ustic Endoaquert Soil taken from Mootilango Sub district, Gorontalo District, Gorontalo Province. Soil description and classification are presented in Table 1.

Coconut and Banana Coir Preparation

Materials were obtained from dried coconut husk that surrounded coconut shell inside. Coconut husk was peeled and separated from coconut shell, then, a smooth outer skin was peeled again until remaining coconut coir (CC). Furthermore, CC was milled to 1 mm. Banana coir (BC) materials were obtained from dried banana bark. Banana bark was peeled and separated from trunk up to 10 sheets of rods into core of the BC. Furthermore, the banana

peel was grounded into powder with a size of 0.05 mm. The CC and BC obtained were weighed according to dosage of each treatment. Before application, water absorption capacities (WAC) of CC and BC were tested through immersion. Immersion was used to determine maximum WAC of the ameliorant material. Calculation of WAC followed the equation:

$$WAC (\%) = [\text{Pre Weight (w0)} / \text{Dry weight (w1)}] \times 100\%$$

Experimental Designs

A factorial design with 3 factors of ameliorant was used, where each factor consisted of 3 dosages of ameliorant and each had 3 replications, so 81 pot experiments were obtained. Sea sand (SS) factors consisted of 0% (S0), 25% (S1), and 50% (S2). CC factor consisted of 0 (C0), 10 (C1), and 20 Mg ha⁻¹ CC (C2). While, the BC factors consisted of 0 (B0), 10 (B1), and 20 (B2) Mg ha⁻¹ BC.

Initial properties of the soil at 0-20 cm depth are presented in Table 2. The soil had clay loamy textures, slow of soil permeability, and ready of swelling and shrinking. Furthermore, the soil chemical properties indicated that the soil organic matter, total N, available P, and K exchanged were low. Relatively neutral of soil pH, cation exchange capacity and base saturation were high. Thus, based on criterion of Center for Soil Research (1983) the soil fertility status was classified as moderate.

Planting Rice and Its Maintenance

Before planting, basic fertilizers were weighted and applied to the soils. Urea 125 kg ha⁻¹ were given twice at ages 0 days after planting (DAP) and 60 DAP, each as much as 62.5 kg ha⁻¹, respectively. Meanwhile, SP36 100 kg ha⁻¹ were given twice at ages 0 days after planting (DAP) and 60 DAP, each as much as 50.0 kg ha⁻¹, respectively, whereas KCl 50 kg ha⁻¹ were given twice at ages 0 days after planting (DAP) and 60 DAP, each as much as 25.0 kg ha⁻¹, respectively.

Ciherang rice seed variety was tested its quality by soaking in saline solution, then planted in trays which were covered by leaves and soil media containing organic material with a ratio of 1:1 until 10 days. A day before planting, the planting medium was watered so that toxic compounds to the rice seedling were moved down to the bottom of the pot. Rice seeds 10 days aged were transferred into plant growing media. Planting was done in planting hole as deep as 8 cm and followed by application of basic fertilizers. During plant growth and development, the maintenance performed was weed cleaning, watering was done every 7 DAP to end

the grain filling phase until flooding, water content was filed capacity. Additication of second basic fertilizers were at 60 DAP. Harvesting was done when plant age was less than 115 days after planting. Observation of plant growth components were plant height, leaf length, leaf number and tiller number. Meanwhile, the observations of yield components were panicle length, panicle number, and rice grain number.

Statistical Analysis

Analyses of Variance (ANOVA) for factorials design were done to study the growth and yield response of rice plants due to application of

ameliorant materials. If there was a significant effect, then it was continued by the least significant difference (LSD) test at 5% level.

RESULTS AND DISCUSSION

Growth Components of Rice Plant

The results of variance analysis for rice growth component planted at Ustic Endoaquert showed that CC and BC did not significantly effect to plant high (Table 3), while the SS significantly affected to plant high. The highest plant height was shown at application of 20% SS which significantly increased plant height compared other treatments. However,

Table 1. Soil description and classification of the Experiment Site.

Location	:	Sidomukti Village, Mootilango Subdistrict, Gorontalo District, Gorontalo Province
Soil classification		
Taxonomy (USDA 2010)	:	Ustic Endoaquert
PPT System	:	Eutric Cambisol
FAO-UNESCO System	:	Cambisol
Parent material	:	Lacustrine
Position physiographic	:	Foot slope, Depression
Topography	:	Flat-Ramps, slopes <2%
Elevation	:	58 m sl
Drainage	:	Poor
Ground water depth	:	Shallow
Vegetation	:	Rice (<i>Oryza sativa</i> L.)
Depth (cm)	Horizon	Descriptions
0-12	Apg1	Gray (10YR 5/1); clay loamy; massive structure; very sticky, plastic; smooth roots, a lot; clear flat.
12-31	Apg2	Gray (10YR 5/1); clays; angular blocky structure, smooth, Weak; very sticky, plastic; rusty brown (10YR 5/3), plain, smooth, clear, spots, sharp; rooting smooth, much; gradually average
31-53	Bwg1	Gray (10YR 5/1); clay; moderate, angular blocky structure, weak; very sticky, plastic; rooting smooth, slightly; frosted flat.
53-71/92	Bwg2	Gray (10YR 6/1); clay; angular blocky structure, rough, weak; very sticky, plastic; obviously choppy.
71/92-119	Bwssg	Dark gray (10YR 4/1); clay moderate, angular blocky structure, moderate; very sticky, plastic; slickenside; rusty brown (10YR 5/3), plain, smooth, clear, tube, clear; frosted flat.
119-150	BCg1	Dark gray (10YR 4/1); clay; angular blocky structure, harsh, strong; very sticky, plastic; rusty brown (10YR 5/3), plain, smooth, clear, tube, clear; clear flat.
150-200	BCg2	Dark gray (10YR 4/1); clay, very sticky, very friable; frosted flat.

Table 2. Initial soil properties of ustic endoaquert at 0-20 cm depth.

Soil Properties	Values	Criterion*
Texture:		
Sand (%)	27	Clay Loamy
Clay (%)	35	
Silt (%)	38	
Soil permeability (cm hour ⁻¹)	1.59	Slow
COLE value	0.98	Real swell-shrinking
Water content availability	8.47	10
Organic (%)	0.69	Very low
Total N (%)	0.06	Very low
C/N ratio	11.62	Moderate
Available-P, Bray 1 (mg P kg ⁻¹)	3.80	Very low
pH H ₂ O	6.48	slightly acid
NH ₄ OAc 1 N pH 7 Extraction:		
K (cmol+ kg ⁻¹)	0.24	Very low
Ca (cmol+ kg ⁻¹)	14.90	High
Mg (cmol+ kg ⁻¹)	6.05	High
Na (cmol+ kg ⁻¹)	0.50	Moderate
CEC (cmol+kg ⁻¹)	30.93	High
Base saturation (%)	70.08	High
Extra 24 KCl 1 M:		
Al ³⁺ (cmol+ kg ⁻¹)	0.00	
H ⁺ (cmol+ kg ⁻¹)	0.06	

*Center for Soil Research (1983).

the plant height still lower than normal growth as reported by Supriatno *et al.* (2007), that plant height of Ciherang varieties were ranged from 107-115 cm.

The highest increasing of plant height was indicated at application of 25% SS which increased plant height by 11.80%, while the lowest one was shown at application of 10 Mg ha⁻¹ CC which only increased by 0.83%. At growth phase, especially plant height it needed sufficient water availability. Application of 25% SS had not yet reduced water storage in soil, so it did not effect on water availability, but it could increase plant height. At application of 50% SS, the plant height decreased until 2.45%. Mishra *et al.* (2012) reported that water available significantly decreased with the increasing of sand in mixture. Water shortages did not apparent at the beginning of vegetative phase and it still could stimulate root development, but at next vegetative phase, plant would be stunted (Sholeh and Riajaya 2000).

Furthermore, Yakup (2008) reported that plant height was significantly affected by soil water availability. In the water-saturated state, despite resistance by soil particles, no water could be

absorbed by the soil except in certain plants such as paddy (Kasli and Efendi 2011).

Test results of WAC of both ameliorant showed that CC had a high WAC (71.77%), while BC was only 28.23%. Wuryaningsih *et al.* (2008) who conducted a study of growth *Anthurium andraeanum* in pot by using coconut bulk media reported that its physical properties were: a very high water content (1,314.41%), low-bulk density (0.09%), high total porosity (120.31%), and high pores water holding (116.6%). In additional, soil texture was classified as clay loamy (Table 2), so the soil puddling and flooding would become more easily dispersible and soil was relatively compact (Isnaini and Suwarno 2005).

Unlike the application of CC, the application of SS and BC had significant effect to leaf length (Table 3). The largest increasing of leaf length was shown at SS application as much as 25% (16.47%) and it was significantly different with control. It seems that application of 25% SS had not yet reduced water storages in soil but it might increase soil pores, so the leaf length was the highest. With the application of 50% SS, the leaf length decreased

Table 3. The rice plant growth components with the application of sea sand, coconut coir and banana coir planted at Ustic Endoaquent soil.

Treatments	Plant Height (cm)	Leaf length (cm)	Leaf number	Tiller number
Sea sand				
0%	27.74a	20.64a	10.29 ^{ns}	12.07a
25%	30.97b	24.04b	14.18	13.37b
50%	30.23ab	23.01ab	14.04	13.29ab
Coconut coir				
0 Mg ha ⁻¹	29.57 ^{ns}	21.99 ^{ns}	11.88 ^{ns}	12.37 ^{ns}
10 Mg ha ⁻¹	30.02	22.92	13.29	13.29
20 Mg ha ⁻¹	29.32	22.78	13.33	13.07
Banana coir				
0 Mg ha ⁻¹	30.95 ^{ns}	23.89a	9.89a	12.56 ^{ns}
10 Mg ha ⁻¹	29.29	21.43b	16.37b	13.19
20 Mg ha ⁻¹	28.65	22.37ab	12.26ab	13.00
Interaction	ns	ns	ns	ns
LSD _{0.05}	3.21	2.42	4.57	1.25

Note: Number that following by same latter in same column did not significantly different at LSD level of 0.05; ns=not significant effect at F level test 0.05.

until 11.48%. Mishra *et al.* (2012) reporting that the water available decreased significantly with increasing sand in mixture. The lowest leaf length was shown at the 0 Mg ha⁻¹ CC (0.62%) and it was not significantly different with 10 and 20 Mg ha⁻¹ CC.

Fazeli *et al.* (2007) stated that effect of a turn single of water availability would reduce growth. Furthermore, Kasli and Effendi (2011) stated that effects of water shortages on some physiological processes of plants were seen in a slow accumulation of dry matter, decreasing leaf expansion rate, and limiting stomata closure photosynthesis. In addition, short-term leaves and leaf area in plants were thought to be influenced by the intensity of the light received in green house where the research took place. The presence of trees around the green house allegedly indirectly affected the reception of light by the plant, even though the distance was about 6 feet from the green house. Garwulan *et al.* (2008) stated that light played an important role in the physiological processes of plants, especially photosynthesis, respiration, and transpiration. The plants required quite diverse intensity of light.

The applications of SS and CC did not have significant effect to leaf number, but BC had significant effect (Table 3). Application of 10 Mg ha⁻¹ BC had significantly increased leaf number by 65.52% and it was significantly different with control. This was presumably due to the ability of

BC to maintain soil moisture and water availability. Indrawati (2009) stated that the stem of BC was strong fiber and resistant to water. It also had pores that were interconnected, and when dry a material would be having absorption and high shelf.

The application of CC and BC did not have significant effect to tiller number, but application of SS had significant effect to it (Table 3). Application of 25% SS increased significantly tiller numbers by 10.77%. Indrawati (2009) stated that the flooding growing media as high as 5 cm was also associated with the formation of tiller numbers. This was in line with reports of Utomo and Rudi, (2000) in Kaseli and Effendi (2011) that the flooding up to approximately 3-5 cm above soil surface condition that considered as good for formation of tillers and at fall phase the flooding would inhibit seedling establishment. The results were reinforced by reports Astuti (2010) that the tiller numbers produced from crops by flooding system were more than by intermittent and kemalir systems. However, in contrast with results of Shi *et al.* (2002) that at the maximum tillering growth phase, the highest tiller numbers were in kemalir water management compared to intermittent and continuous floodings.

Sumardi *et al.* (2007) that water use efficiency for rice cultivation without flooding condition were as much as 19.581%, while with continuously flooding its efficiency was as much as 70.907% only. In this study also showed that the higher the plant would be followed by a greater tiller numbers. This

was in contrast with the result of Aldi *et al.* (2004); Hartati and Suwarta (2004); Rahayu and Harjoso (2010) who reported that tiller number showed contrary to values of plant height, where the higher tiller number was the lower plant high.

Although it had not significant effect to tiller numbers, but application of 10 Mg ha⁻¹ CC and BC increased tiller number by 7.44% and 5.04%, respectively compared to control. This was related a to high ability to absorb water of CC and BC, so that they would enough moisture and water availability. Ramesh *et al.* (2010) reported that combination of soil and coconut coir as much as 4% showed changes in the nature of Catton black soil compactness better that control. Contribution of ameliorant effect to plant growth was in order of SS > BC > CC.

Yield Components of Rice Plant

Analysis of variance for rice yield component resulted that application of CC and BC had significantly increased panicle numbers, but application of SS did not have significantly effect to panicle numbers. It seems that percentage of panicle numbers with CC and BC application were relative similar. However, the different between treatments were increasing of percentage figures. Application of 20 Mg ha⁻¹ BC significantly increased panicle number by 20.47% compared to 10 Mg ha⁻¹ BC and by 23.41 compared to control. This suggested that BC application was able to maintain water

availability due a to high ability to absorb water. The BC was quite strong and resistant to water and had pores that where interconnected and when dry, a material would be having absorption and high shelf (Indrawati 2009). Meanwhile, although it had not significantly effect to panicle numbers, but application of 50% SS increased tiller numbers by 19.36% compared to control.

Soil sand fraction was clay loamy textures, so that application of 50% SS to the soil could reduce weight to light fraction, particularly declined COLE index. Sand textures affected status and distribution of water, so affecting root system and root depth (Walter *et al.* 2000; Oliver and Smettem 2002). Thus, the plant root system was not disturbed due to lock of the possibility of soil cracking, so the process of water and nutrients absorption could be better.

Application of 20 Mg ha⁻¹ CC significantly increased panicle numbers by 29.54% compared to 10 Mg ha⁻¹ CC. This was presumably related to soil water availability. The WAC level of CC was high and would be able to absorb and retain water availability in the soil, so that more CC would be followed by the greater panicle numbers. This was consistent with statement of Dachban (2012) that rice storage water for 4 days at generative periods and following 2 weeks was a sensitive periods to water storages.

In addition, the SS, CC and BC significantly increased panicle length. Application of 10 Mg ha⁻¹

Table 4. The rice yield component at application of sea sand, coconut coir and banana coir planted at Ustic Endoaquert soil.

Treatments	Panicle number	Panicle length (cm)	Rice grain number
Sea sand			
0%	2.48ns	8.97b	8.64ns
25%	2.59	7.35a	7.98
50%	2.96	8.28ab	9.07
Coconut coir			
0 Mg ha ⁻¹	2.59ab	6.41a	7.14a
10 Mg ha ⁻¹	2.37a	8.80b	9.52b
20 Mg ha ⁻¹	3.07b	9.39b	9.04ab
Banana coir			
0 Mg ha ⁻¹	2.52ab	7.33a	7.49ns
10 Mg ha ⁻¹	2.41a	9.49b	9.48
20 Mg ha ⁻¹	3.11b	7.78a	8.72
Interaction	ns	ns	ns
LSD _{0.05}	0.63	1.24	2.05

Note: Number that following by same latter in same column did not significantly effect at LSD level of 0.05; ns=not significant effect at F level test 0.05.

BC significantly increased panicle length by 29.47% compared to control and by 21.8% compared to 20 Mg ha⁻¹ BC. This was presumably related to soil water availability of each treatment. The research conducted by Dachban (2012) showed that the panicle length at level of 3 mm, 2 mm, and 1 mm decreased long panicle length by 0.77%, 5.80%, and 45.10% respectively, due to reduction of plant yields because of water. In fact, Astuti (2010) reported that panicle length and rice grain numbers were not affected by irrigation, but each variety had panicle length and rice grain numbers per panicle significantly different due to the genetic. Meanwhile, 10 Mg ha⁻¹ CC significantly increased rice grain numbers by 33.33% compared to control but did not have significantly different with 20 Mg ha⁻¹ CC (Table 4). Apparently, long panicles would be relatively followed by a greater rice grain numbers. It was in line with Setiobudi *et al.* (2008) that the rice grain numbers was determined by genetic properties of plants especially panicle length, tassel branch, and a differentiation rice grains.

CONCLUSIONS

Application of sea sand and banana coir significantly increased leaf length where the highest increasing percentage of 16.47% was achieved at 25% SS application. Their effect on leaf numbers and tiller numbers were relatively not similar pattern where leaf number only increased about 65.52% by banana coir application, while tiller numbers only increased about 10.77% by sea sand application. Furthermore, the application of coconut coir and banana coir significantly increased panicle numbers up to 29.53% and 29.05%, respectively compared to control. All ameliorants significantly increased panicle numbers, but the best was coconut coir that could increase up to 46.49% at 20 Mg ha⁻¹ CC compared to sea sand and banana coir application. Only coconut coir significantly increased the rice grain numbers.

ACKNOWLEDGEMENTS

The research team would like to thank Directorate General of Higher Education (DGHE), the Ministry of Education and Culture of Republic Indonesia that had given us the opportunity to get a Competitive research Grant funding in fiscal year 2012 with contract number was 33/UN47.D2/PL/2012.

REFERENCES

- Aldi MA, Darjanto and ADH Totok. 2004. Effect of weed control methods on the results of four upland rice cultivars. *J Agrin* 8: 100-107.
- Al-Omran AM, AM Falatah, AS Sheta and AR Al-Harbi. 2004. Clay deposits for water management of sandy soils. *Arid land Res Manage* 1: 171-183.
- Anggarwulan E, Solichatun and W Mudyantini. 2008. Physiological characters of purse (*Xanthosoma sagittifolium* L. Schott) in several of light intensity (shading) and water availability. *J Biodiver* 9: 264-268.
- Astuti DN. 2010. Effect of irrigation systems on the growth and productivity of some varieties of rice (*Oryza sativa* L.). [BSc thesis] Bogor Agricultural University, Bogor (in Indonesian).
- BPS RI [Badan Pusat Statistik Republik Indonesia]. 2012. Statistics Indonesia in 2012. Central Bureau of Statistics, Jakarta.
- Center for Soil Research. 1983. Terms of Reference of Soil Capability Survey No 22119&3. Transmigration Supporting Agricultural Research Project Agency for Agricultural Research and Development Ministry of Agriculture, Bogor (in Indonesian).
- Cepy and W Wangiyana. 2011. Growth and yield of rice (*Oryza sativa* L.) in the media and Entisols-Vertisols at different water management techniques and types of fertilizers. *J Agro Crop* 4: 49-56.
- Dachban SMB. 2012. Effect of giving water level on the growth and yield several varieties of upland rice. *J Sci Higher Edu* 5: 6-15.
- Deckers J, O Spargaren and F Nachtergaele. 2001. Vertisols: Genesis properties and soil scape management for sustainable development. In: JK Syers, FWT Penning De Vries and P Nyamudeza (eds). The Sustainable Management of Vertisols. IBSSAM Proceeding No. 20, pp. 3-24.
- Fazeli F, M Ghorbanli and V Nikham. 2007. Effects of drought on biomass, protein content, lipid peroxidation and antioxidant enzymes in two sesame cultivars. *Biol Plantarum* 5: 98-103.
- Hartati and Suwarta. 2004. Yield and quality of two varieties of upland rice grown intercropped with biofertilizer treatments. *J Agronomika* 4: 7-9.
- Hikmatullah, BH Praseiyo and M Hendrisman. 2002. Vertisol from Gorontalo areas: physical properties of the chemical and mineralogical composition. *Soil Water J* 3. 21-32 (in Indonesian).
- Indrawati E. 2009. Sound absorption coefficient of acoustic materials of banana with different densities. [BSc Thesis] Islamic University Maliki, Malang.
- Isnaini S and P Suwarno. 2005. Rice cultivation with no tillage systems for ten seasons in Kerjaloman, Talangpadang, Lampung. *J Stigma* 8: 551-554 (in Indonesian).
- Kasli and AAR Effendi. 2011. High influence on plant growth puddle rice (*Oryza sativa* L.) in pots. *J Jerami* 4: 246-212.
- Kusnarta IGM. 2012. Kusnarta IGM. 2012. Study of soil properties affecting stability of furrows fields forever in rainfed paddy soil of Vertisol-Lombok. Summary [PhD Dissertation] Gadjah Mada University, Yogyakarta. 6 Pp.
- Mukanda N and A Mapiki. 2001. Vertisols management in Zambia. In: JK Syers, FWT Penning De Vries and

- P Nyamudeza (eds). The sustainable management of Vertisol. Proceedings IBSSAM no 20, pp. 129-127.
- Mishra PK, GRM Sankar, UK Mandal, SK Nalatwadmath, SL Patil and S Manmohan. 2012. Amelioration of Vertisol with sand for management of soil physical, chemical and hydraulic properties in south India. *Indian J Soil Conser* 40: 13-21.
- Narka IW and Wiyanti. 1999. Effect of sand and organic materials against each Vertisol soil physical properties. *J Agri Trop* 18: 11-15.
- Nugroho AW. 2013. Effect of Planting Media Composition on *Casuarina equisetifolia* var. *Incana* Growth in the Coastal Sand Dune. *Forest Rehabilitation* 1(1): 114.
- Nurdin. 2011. Development and rainfed paddy soils potency derived from lacustrine material in Paguyaman, Gorontalo. *J Trop Soil* 16: 267-278. doi: 10.5400/jts.2011.16.3.267
- Nursyamsi D. 2009. Effect of potassium and com varieties for organic acids from root exudates, uptake of N, P and K production plants and corn stover (*Zea mays* L.). *J Agron Indon* 37: 107-114.
- Oliver YM and KRJ Smethem. 2002. Predicting water balance in a sandy soil: model sensitivity to the variability of measured saturated and near saturated hydraulic properties. *Aust Soil Res* 43: 87-96.
- Prasetyo BH. 2007. Differences vertisol soil properties from a variety of parent materials. *J Agric Sci* 9: 20-31.
- Rajiman. 2009. The Effect of NPK Fertilizer on shallot field at coastal sandy land. *J Agric Sci* 5: 52-60.
- Rahayu AY and T Harjoso. 2010. Agronomist and physiological characters of upland rice grown in soil husky media on water conditions in the lower capacity field. *J Akta Agrosia* 13: 40-49.
- Ramesh HN, KVM Krishna and HV Mamatha. 2010. Compaction and strength behavior of lime-treated coir fiber Black Cotton soil. *J Geomech Eng* 2: 19-28.
- Ravina I and J Magier. 1984. Hydraulic conductivity and water retention of clay soil containing coarse fragments. *Soil Sci Soc Am J* 48: 736-740.
- Sari HC, S Darmanti and ED Hastuti. 2006. Plant growth of emprit ginger (*Zingiber officinale* var. *Rubrum*) on growing media sand with different salinity. *Bull Anat Physiol* XIV (2): 19-29.
- Setiobudi D, B Abdullah, H Sembiring and IP Wardana. 2008. Peningkatan hasil padi tipe baru melalui pengelolaan hara pupuk nitrogen. Prosiding Simposium V Tanaman Pangan-Inovasi Teknologi Tanaman Pangan. Pusat Penelitian dan Pengembangan Tanaman Pangan, Badan Penelitian dan Pengembangan Pertanian 2: 345-353 (in Indonesian).
- Shi Q, X Zeng, M Li, X Tan and F Xu. 2002. Effects of different water management practices on rice growth. In: BAM Bouman, H Hengsdijk, B Hardy, PS Bindraban, TP Tuong and JK Ladha (eds). Water-wise rice production. International Rice Research Institute, Los Banos, pp. 3-13.
- Sholeh M and PD Riajaya. 2000. Anticipating climate storage on the cultivation of tobacco. *Monogr Balittas* 6: 12-15.
- Subiyanto B, R Saragih and E Husin. 2003. Utilization of coconut coir dust as water and oil absorbent materials such as particle board panels. *J Trop Wood Sci Tech* 1: 26-34.
- Sudaryatno T, R Kustiari and HP Saliem. 2010. Estimates food needs in 2010-2050. In: Book Analysis of Land Resources towards Sustainable Food Security. Agency Research and Development of Agriculture, Jakarta, pp. 1-23.
- Sumardi, Kasli, M Kasim, A Syarif and N Akhir. 2007. Respon padi sawah pada teknik budidaya secara aerobik dan pemberian bahan organik. *J Akta Agrosia* 10: 65-71 (in Indonesian).
- Supriatno B, AA Darudjat, Satoto, SE Baehaki, N Widiarta, SD Indrasari, OS Lesmana and H Sembiring. 2007. Description rice varieties. Rice Research Institute, 80p.
- Suswono. 2012. Indonesia rice consumption too much. <http://www.republika.co.id/berita/nasional/umum/12/04/03/mlwjln-mentan-konsumsi-berasindonesia-terlalu-banyak>. Accessed on 18 December 2012.
- Toha HM and K Pirngadi. 2004. Effect of plant density and weed control on the results of some rice varieties seeded systems in rainfed land. *J Agrivigor* 3: 170-177.
- Utomo M and Nazaruddin. 2000. Paddy rice cultivation without tillage. Penebar Swadaya, Jakarta.
- Walter A, WK Silk and U Schur. 2000. Effect of soil pH on growth and cation deposition in the root tip of *Zea mays* L. *J Plant Growth Reg* 19: 65-76.
- Wuryaningsih S, T Sutater and B Tjia. 2008. Growth *Anthurium andraeanum* pot plant on coconut bulk media. *J Agric Res* 18: 31-38.
- Yakup. 2008. Effect of soil water availability and the time interval mulching on the growth and development of upland rice crop. *J Agria* 5: 5-8

Effect Application of Sea Sand, Coconut and Banana Coir on the Growth and Yield of Rice Planted at Ustic Endoaquert Soil

ORIGINALITY REPORT

7 %

SIMILARITY INDEX

4 %

INTERNET SOURCES

4 %

PUBLICATIONS

1 %

STUDENT PAPERS

PRIMARY SOURCES

1	World Soils Book Series, 2014. Publication	1 %
2	epdf.pub Internet Source	1 %
3	core.ac.uk Internet Source	<1 %
4	www.europeanproceedings.com Internet Source	<1 %
5	Akram La Kilo, La Ode Aman, Ismail Sabihi, Jafar La Kilo. "Studi Potensi Pirazolin Tersubstitusi 1-N dari Thiosemicarbazone sebagai Agen Antiamuba melalui Uji In Silico", Indo. J. Chem. Res., 2019 Publication	<1 %
6	etheses.bham.ac.uk Internet Source	<1 %
7	Submitted to Universitas Jenderal Soedirman Student Paper	<1 %

8	csdlkhoahoc.hueuni.edu.vn Internet Source	<1 %
9	www.tsbf.org Internet Source	<1 %
10	Funda Yirmibesoglu, Nilgun Ergun. "Property and Personal Crime in Istanbul", European Planning Studies, 2007 Publication	<1 %
11	Marlene Alves, Antonio Gonzalez, Gilberto Colodro, Helio Junior, Eva Vázquez. "Influence of Biosolids Rate on Chemical Properties of an Oxisol in São Paulo, Brazil", Communications in Soil Science and Plant Analysis, 2006 Publication	<1 %
12	protan.studentjournal.ub.ac.id Internet Source	<1 %
13	Suwignyo, B., P. Suryanto, E. T. S. Putra, T. Alam, and S. D. A. Prianto. "Potential of Corn as Forage on Alfisol and Vertisol Soil in Agrosilvopastural System with Kayuputih (Melaleuca leucadendron Linn)", Journal of Agricultural Science, 2015. Publication	<1 %
14	jurnal.polbangtanyoma.ac.id Internet Source	<1 %
15	www.mdpi.com Internet Source	<1 %

<1 %

16

Choudhury, B.U.. "Yield and water productivity of rice-wheat on raised beds at New Delhi, India", Field Crops Research, 20070201

Publication

<1 %

17

journal.ipb.ac.id

Internet Source

<1 %

18

manajemenkesuburantanah.blogspot.com

Internet Source

<1 %

19

ml.scribd.com

Internet Source

<1 %

20

pt.scribd.com

Internet Source

<1 %

21

Craven, D.. "Between and within-site comparisons of structural and physiological characteristics and foliar nutrient content of 14 tree species at a wet, fertile site and a dry, infertile site in Panama", Forest Ecology and Management, 20070130

Publication

<1 %

22

Mohammad Asadul Haque, Md Jahiruddin, Md. Islam, Md Fazlul Hoque et al. "Identifying suitable source and optimum rate of silicon in

<1 %

rice at Ganges Delta Coastal Zone Soils",
Research Square Platform LLC, 2022

Publication

23

Sushanta Kumar Naik, Dilip Kumar Das. "
Effect of split application of zinc on yield of
rice (L.) in an inceptisol ", Archives of
Agronomy and Soil Science, 2007

Publication

<1 %

24

C Ribeiro, M Madeira, M.C Araújo.
"Decomposition and nutrient release from
leaf litter of Eucalyptus globulus grown under
different water and nutrient regimes", Forest
Ecology and Management, 2002

Publication

<1 %

25

I A Sipahutar, A F Siregar, L Anggria, Husnain.
"Magnesium and silicon fertilizer application
to promote rice growth and production", IOP
Conference Series: Earth and Environmental
Science, 2021

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On