SUBMISSION



Submitted of manuscript journal

2 pesan

Nurdin <nurdin@ung.ac.id> Kepada: editor@sysrevpharm.or 11 November 2020 pukul 21.15

Best regards, Editor in Chief

I hereby submit a journal manuscript with the title: Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo. We hope that you will be accepted and have the opportunity to contribute to this journal. Regards

Nurdin

Study of Quality and Land Characteristics that Controlling of Composite Maize Yield in Gorontalo- Nurdin.docx
1438K

------ Pesan Yang Diteruskan ------From: Nurdin <nurdin@ung.ac.id> To: editor@sysrevpharm.or Cc: Bcc: Date: Wed, 11 Nov 2020 21:15:53 +0700 Subject: Submitted of manuscript journal ----- Message truncated -----

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Nurdin^{1*}, M. L Rayes², Soemarno², Sudarto²

 ¹Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145
 ³Soil Science Departmen, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145

*e-mail: nurdin@ung.ac.id

Abstracts. The challenge of composite maize developing in the future is the low productivity because the maize is grown on land that is not suitable for land quality. This study aims to determine the land quality and land characteristics that control the composite maize productivity in Gorontalo Province. A total of 33 land units were surveyed and their land observed to obtain data on morphology and soil characteristics, climate and terrain characteristics, as well as composite maize productivity data through ubinan plots and direct interviews with maize farmers. Partial least square of structural equation models (PLS-SEM) analysis has been used to determine the land quality and land characteristics that control the composite maize productivity through variable validity and reliability tests, as well as structural model tests. The results showed that the manifest variables were air temperature, rainfall, wet months, dry months, LGP, drainage, coarse materials, effective depth, pH H₂O, pH KCl, C-organic, total N, available P, available K, ESP, slopes, soil erosion, inundation height, inundation time, surface rock, and rock outcrops were valid and able to explain well the latent variables. Furthermore, the latent variables were temperature, water availability, oxygen availability, nutrient retention, nutrients availability, sodicity, erosion hazard, flood hazard, and land preparation used has good composite reliability and high reliability because of the composite reliability and alpha cronbach >0.6. Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of coarse materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

Keywords: Quality, characteristic, land, productivity, maize, composite.

INTRODUCTION

Low maize productivity is still a major problem in efforts to increase maize production in Indonesia. Until 2019, maize productivity had only reached 5.24 tons/ha [1], while the potential for maize productivity in Indonesia could reach 10-11 tons/ha [2]. Even though the government has rolled out various programs to increase maize production and productivity in order to achieve national maize self-sufficiency.

Gorontalo Province is one of the maize producing centers in Indonesia with an average maize productivity achievement until 2019 of 5.03 tons/ha [3] or still far below the average national maize productivity. All this time, maize farmers has been more dominant in planting maize with hybrid and composite varieties. There are no references to the productivity of

hybrid or composite maize in this area, so it is assumed that the maize productivity achievements are relatively the same at 5.24 tons/ ha. In fact, specifically the productivity of composite maize can reach 5-6 tons/ha [4] [5] or as much as 8.5 tons/ha [6]. Composite maize, besides its productivity potential is relatively similar with hybrid maize productivity, it also has advantages including being more adaptive in acid soils [5] and can be used as seeds for the next growing season, while hybrid maize can't be planted again. The use of composite maize can reduce the dependence of maize farmers on subsidized hybrid maize seed from the government, so that if the maize seed subsidy has stopped, the farmers can plant the composite maize again.

The challenge ahead in developing of composite maize was the low of composite maize productivity, so it is necessary to address the root of the problems. Maize planted on land with low productivity potential is one of the causes for the low of maize productivity [7]. Meanwhile, land characteristics and land quality have a close relationship with maize productivity [8] and each land quality has a significant effect on land suitability for certain uses [9], especially for maize. Research on land quality that controls the productivity of composite maize has been conducted in the Bogor area using stepwise regression analysis [10]. The use of structural equation modeling (SEM) analysis in determining the land characteristics and land quality that control plant productivity has not been widely published, except on older cocoa plants in Kolaka Timur Regency, Southeast Sulawesi Province that used SEM analysis [11]. Meanwhile, the use of SEM analysis specifically to determine the relationship between land quality and maize productivity has not been found.

The response of maize plants to the diversity of land characteristics and land quality will vary, so it is important to know of these that control of the maize productivity. The complex diversity of land characteristics and land quality in the field really requires a comprehensive analysis technique that is able to simplify the complexity in one analysis system. One analysis option is to use SEM analysis. SEM analysis is able to analyze how much influence each indicator (manifest) of soil physical and chemical properties (latent) has on production in one analysis unit [11]. The use of SEM is very helpful to determine the effect on indicators and to produce a model that is better than other multivariate analyzes [12] [13]. Partial Least Square (PLS) is a variant of SEM which has a higher level of flexibility because PLS is based on variants, so that the number of samples used does not need to be large, ranges from 30-100, and does not require normal multivariate assumptions compared to CB-SEM that requires a large data sample size (>100) and the data must be multivariate normal distribution [14] [15]. Therefore, a research on land quality that controls the composite maize productivity was carried out using SEM-PLS analysis based on the consideration complexity of land characteristics and land quality, as well as limited data in the land unit in the study area. The purpose of this study was to determine the land quality and land characteristics that control the composite maize productivity oin Gorontalo.

MATERIALS AND METHOD

This research location in the Sustainable Agriculture Area of Gorontalo Province (Figure 1) and the Soil Laboratory of the Soil Department, Agriculture Faculty of Brawijaya University. The timing of this research was started in December 2019 - May 2020. The tools used included the computer, SmatPLS version 2.0, SPSS, Microsoft Excel, and Microsoft Word. While the materials studied included the morphological and soil characteristics data, climate and terrain characteristics data that had been grouped into their respective land qualities, as well as composite maize productivity data from the study area.

Soil surveys and land observations were carried out to obtain morphological and soil characteristics data, climate and terrain characteristics data from the research area.

Meanwhile, composite maize productivity data was obtained from the results of ubinan directly on the land of maize farmers and from direct interviews with maize farmers on 33 land units. Furthermore, the diversity of sizes and data units (ratio data) of land characteristics were converted in the form of interval data which were represented as follows were 1 (very low), 2 (low), 3 (medium), 4 (high), and 5 (very high)). After the data was ready, the analysis process is continued using SEM-PLS (Figure 2).



Figure 2. Research Operational Framework

The latent variable in this study was the quality of the land consisting of: temperature (X1), water availability (X2), oxygen availability (X3), root media (X4), nutrient retention (X5), available nutrients (X6), sodicity (X7), erosion hazard (X8), flood hazard (X9), and land preparation (X10). While the manifest variable was the characteristic of the land which consists of temperature (X1.1), rainfall (X2.1), wet months (X2.2), dry months (X2.3), LGP (X2.4), drainage. (X3.1), texture (X4.1), coarse material (X4.2), effective soil depth (X4.3), pH H2O (X5.1), pH KCI (X5.2), C-organic (X5.3), CEC (X5.4), base saturation (X5.5), total N (X6.1), available P2O5 (X6.2), K available (X6.3), ESP (X7.1), slope (X8.1), soil erosion (X8.2), inundation height (X9.1), length of inundation (X9.2), surface rock (X10.1), and rock outcrop (X10.2). The use of SEM-PLS in this study consisted of testing the validity, reliability of the research variables, and testing the structural model. In summary, the test using SEM-PLS was described as follows:

- a. Research Variable Validity Testing. The basic evaluation carried out in the SEM-PLS analysis is to evaluate the measurement model (outer model) with the aim of knowing the validity and reliability of indicators in measuring research latent variables through convergent validity, discriminant validity, and composite reliability. Convergent validity testing on SEM-PLS is seen from the size of the outer loading of each indicator on its latent variable. A loading factor value above 0.70 is highly recommended, but a loading factor value of 0.50-0.60 can still be tolerated with a t-statistic value of more than 1.96 or a small p-value of 0.05. The loading factor of an indicator with the highest value is the strongest or most important measure in reflecting the latent variable in question. Discriminant validity is an evaluation of the outer model in SEM-PLS using cross loading values to test valid and reliable indicators in explaining or reflecting latent variables. If the correlation of the latent variables, then the latent variable is able to predict the indicator better than other latent variables and is said to be valid.
- b. Research Variable Reliability Testing. Composite reliability and alpha cronbach were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is> 0.60.

c. Structural Model Testing. Testing of the structural model (inner model) is carried out after the relationship model is built in accordance with the observed data and the suitability of the overall model (goodness of fit model). Testing of structural models and hypotheses is carried out by looking at the estimated value of the path coefficient and the critical point value (t-statistic) which is significant at $\alpha = 0.05$. Testing the relationship model and hypothesis between variables can be done by testing the direct correlation coefficient between variables. The results of testing the relationship between the X variables and the Y variable in this study are shown by the correlation coefficient and t-statistic, and also seen in the path diagram.

RESULTS AND DISCUSSION

a. Research Variable Validity

The loading factor value of the research variables, where the loading factor value on the indicators was mostly more than the critical limit of 0.7 with a confidence level of 95% (Table 1). The value of the loading factor which is below the tolerance value of 0.5 at the 95% confidence level where the t-statistic value of each indicator is smaller than the t-table (1,960) is found in the soil texture indicator of the latent variable root media (X₄) which is only 0.173 is also the CEC indicator (X_{5.4}) and the base saturation indicator (X_{5.5}) of the nutrient retention latent variable (X₅), which are only 0.399 and 0.482 respectively. This means that these indicators have not been able to properly form or explain their latent variables.

Effect of	Loading factors	Status		
Air temperature $(X_{1.1})$	->	Temperature (X ₁)	1.000	Valid
Rainfall (X _{2.1})	->		0.981	Valid
Wet months $(X_{2,2})$	->	Water evailability (V)	0.989	Valid
Dry months $(X_{2.3})$	->	water availability (X_2)	0.827	Valid
LGP $(X_{2.4})$	->		0.968	Valid
Drainage (X _{3.1})	->	Oxygen availability (X ₃)	1.000	Valid
Texture $(X_{4,1})$	->		0.173	Not valid
Coarse material $(X_{4,2})$	->	Rooting media (X ₄)	-0.921	Valid
Effective depth $(X_{4.3})$	->		0.912	Valid
pH H ₂ O (X _{5.1})	->		0.768	Valid
pH KCI (X _{5.2})	->		0.772	Valid
C-Organic (X _{5.3})	->	Nutrient retention (X_5)	0.710	Valid
$CEC(X_{5.4})$	->		0.399	Not valid
Base saturation $(X_{5.5})$	->		0.482	Not valid
N Total $(X_{6.1})$	->		0.799	Valid
Available P $(X_{6.2})$	->	Nutrient availability (X ₆)	0.521	Valid
Available $K(X_{6.3})$	->		0.886	Valid
$ESP(X_{7.1})$	->	Sodicity (X ₇)	1.000	Valid
Slope $(X_{8.1})$	->	Encoder barand (\mathbf{V})	0.974	Valid
Soil erosion $(X_{8.2})$	->	Erosion nazaru (X_8)	0.957	Valid
Inundation height (X _{9.1})	->	E_{1} and H_{2} and (\mathbf{V}_{1})	0.993	Valid
Inundation period (X _{9.2})	->	Flooding hazard (A9)	0.991	Valid
Surface rock $(X_{10.1})$	->	Land momentian $(\mathbf{V}_{\mathbf{v}})$	0.998	Valid
Rock outcrop (X _{10.2})	->	Land preparation (Λ_{10})	0.998	Valid
Productivity $(Y_{1.1})$	->	Local maize productivity (Y_1)	1.000	Valid

Table 1. Outer loading research variables

Indikator	Temperature (X ₁)	Water availability (X ₂)	Oxygen availability (X ₃)	Rooting media (X ₄)	Nutrient retention (X ₅)	Nutrient availability (X ₆)	Sodicity (X ₇)	Erosion hazard (X ₈)	Flooding hazard (X9)	Land preparation (X ₁₀)	Composite maize productivity (Y ₁)
Air temperature $(X_{1.1})$	1	0.952309	0.059098	-0.08736	-0.37805	-0.06653	0.38176	0.016269	-0.10297	0.19833	0.042282
Rainfall (X _{2.1})	0.968555	0.980906	0.114576	0.052348	-0.24379	0.058536	0.356547	-0.0379	-0.04621	0.056015	0.156751
Wet months $(X_{2.2})$	0.926635	0.989185	0.173659	-0.005903	-0.25644	0.062873	0.374745	-0.06373	-0.04367	0.060342	0.177251
Dry months $(X_{2.3})$	0.759123	0.82697	0.141078	-0.238735	-0.42612	-0.10563	0.47553	-0.11715	0.027746	0.215367	0.076041
LGP $(X_{2.4})$	0.900431	0.96821	0.13569	-0.003834	-0.28223	0.056251	0.459669	-0.12209	-0.04398	0.059938	0.193991
Drainage $(X_{3.1})$	0.059098	0.144225	1	0.129338	-0.24128	0.057861	0.084339	-0.50344	0.236555	-0.22277	0.400657
Texture (X _{4.1})	-0.02057	-0.01261	-0.16957	0.172551	0.242032	0.12283	0.217308	0.196875	-0.00074	-0.02261	0.09248
Coarse material (X _{4.2})	-0.00333	-0.1005	-0.13244	-0.921096	-0.38256	-0.6112	0.18822	0.322934	-0.26391	0.846957	-0.35202
Effective depth $(X_{4.3})$	-0.17758	-0.09256	0.165016	0.912088	0.3519	0.355112	-0.23141	-0.19005	0.095721	-0.76736	0.180089
pH H ₂ O (X _{5.1})	-0.40346	-0.38437	-0.3719	0.29356	0.767791	0.27088	-0.17175	0.151553	-0.02966	-0.08478	0.186569
pH KCl (X _{5.2})	-0.25953	-0.22811	-0.44804	0.342269	0.771872	0.272936	-0.02729	0.167533	0.098977	-0.18312	0.268161
C-Organic $(X_{5.3})$	-0.29516	-0.13852	0.096529	0.248076	0.710022	0.612498	0.073184	-0.4692	0.063874	-0.1793	0.384332
$CEC(X_{5,4})$	0.066756	0.115697	0.003345	0.084182	0.399393	0.421251	0.373179	-0.05735	0.15285	-0.01387	0.281455
Base saturation $(X_{5.5})$	-0.30026	-0.25724	-0.10527	0.412102	0.481624	0.361795	-0.60079	-0.0895	-0.13592	-0.48759	0.136266
N Total $(X_{6.1})$	0.002878	0.137879	0.07154	0.268606	0.545283	0.798694	0.030267	-0.37884	-0.10212	-0.2485	0.427705
Available P $(X_{6.2})$	-0.09821	-0.09791	-0.44547	0.211821	0.409315	0.520984	-0.28705	-0.057	0.033581	-0.26033	-0.02547
Available $K(X_{6.3})$	-0.09732	-0.01031	0.06693	0.614343	0.51245	0.885686	-0.3292	-0.29441	0.237691	-0.6422	0.49531
$ESP(X_{7.1})$	0.38176	0.405078	0.084339	-0.186069	-0.06947	-0.21259	1	-0.01035	0.201152	0.361936	-0.0249
Slope $(X_{8.1})$	-0.02207	-0.12714	-0.51717	-0.295103	-0.1643	-0.40295	-0.03466	0.973779	-0.34215	0.324431	-0.64795
Soil erosion $(X_{8.2})$	0.064136	-0.00224	-0.44709	-0.166166	-0.11161	-0.32907	0.021581	0.956588	-0.12926	0.257787	-0.48649
Inundation height $(X_{9.1})$	-0.08956	-0.02635	0.225421	0.194354	0.082178	0.127762	0.193925	-0.26735	0.992798	-0.13415	0.175472
Inundation period $(X_{9.2})$	-0.11594	-0.06329	0.244833	0.199427	0.048584	0.078386	0.205739	-0.2425	0.991369	-0.11616	0.135302
Surface rock $(X_{10.1})$	0.212772	0.074279	-0.23401	-0.854273	-0.28568	-0.55023	0.376036	0.319248	-0.13208	0.997623	-0.28655
Rock outcrop $(X_{10.2})$	0.183196	0.051703	-0.21067	-0.868319	-0.29655	-0.55537	0.34638	0.290608	-0.12053	0.997697	-0.28228
Productivity $(Y_{1.1})$	0.042282	0.177277	0.400657	0.304774	0.418519	0.534535	-0.0249	-0.59733	0.157534	-0.28507	1

Table 2. Cross loading of research variables

The standard of loading factor is greater equal to 0.50 [16] [17] [13]. However, in general, based on the indicated values, it can be concluded that the latent variables of land quality have been able to be well established or explained by each indicator and can be said to be convergent valid on these indicators. The cross loading value for the indicators of latent variables on average is above the cross loading value of the indicator is for other latent variables (Table 2). That is, the greatest cross loading value on the indicator is found in the latent variable too, except for the texture indicator (X_{4.1}) of the root media variable (X₄), the CEC indicator (X_{5.4}) and base saturation (X_{5.5}) of the nutrient retention variable (X₅) whose cross loading value is still smaller (<0.5) than the cross loading value of other latent variables. The standard of loading factor is ≥ 0.50 [16] [17] [13]. Thus, the indicators of each latent variable are mostly able to explain the latent variable itself better than the other variables, so that the research variables are said to be discriminant valid.

b. Research Variable Reliability

Composite reliability and Cronbach alpha were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is above 0.60 [18]. The composite reliability value on each research variable is more than the limit value (>0.6), except for the root media variable (Table 3). The composite reliability value and the Cronbach alpha value is greater than 0.6 so that the latent variable has good composite reliability and high reliability. A construct is said to be reliable if the Cronbach Alpha value must be >0.6 [19]. Thus, all indicators used in this study have met the criteria or are feasible to be used in the measurement of all latent variables because they have good validity and high reliability. The results of the evaluation of convergent validity and discriminant validity of indicators or variables as well as composite reliability and alpha Cronbach for indicators or variables can be concluded that the indicators as measures of latent variables are valid and reliable measures respectively.

Laten variables	Composite reliability	Alpha Cronbach
Temperature (X ₁)	1.000000	1.000000
Water availability (X ₂)	0.970030	0.965126
Oxygen availability (X ₃)	1.000000	1.000000
Rooting media (X ₄)	0.020314	-1.055192
Nutrient retention (X ₅)	0.770518	0.628062
Nutrient availability (X ₆)	0.788289	0.681393
Sodicity (X ₇)	1.000000	1.000000
Erosion hazard (X_8)	0.964615	0.927731
Flooding hazard (X ₉)	0.992053	0.984010
Land preparation (X ₁₀)	0.997657	0.995304

Table 3. Composite reliability and Cronbach's Alpha values of research variables

b. Structural Model Testing

The structural model (inner model) is evaluated by looking at the coefficient value of the relationship path parameter between latent variables. It seems that the soil quality of the root media, nutrient retention, and available nutrients showed a positive correlation and had a significant effect on the productivity of composite maize (Table 4). The quality of land preparation shows a negative correlation and has a significant effect on the productivity of composite maize. This indicates that the better rooting media, available nutrient and nutrient retention and a decrease in the level of land preparation as the productivity of composite maize increases. The results of this study are slightly different from the research report [8] regarding the quality of soil rooting media which has not affected the productivity of maize in the Bogor area, but the quality of soil nutrient retention and available nutrients has a significant effect on the productivity of maize relatively the same as the results of this study.

	Endogenous variables					
Exogenous variables	Composite maize produktivity (Y)					
	Path coeffisient	t-statistics ($t_{critics} = 2.00$)				
Temperature (X_1)	0.086	1.531				
Water availability (X_2)	0.457	-0.491				
Oxygen availability (X ₃)	0.099	0.371				
Rooting media (X ₄)	0.091*	2.250				
Nutrient retention (X ₅)	0.740*	2.291				
Nutrient availability (X_6)	0.283**	6.509				
Sodicity (X ₇)	-0.194	-0.036				
Erosion hazard (X_8)	-0.043	-1.043				
Flooding hazard (X ₉)	0.050	-0.050				
Land preparation (X_{10})	-0.386*	-2.339				

Table 4	Path	coefficient	and	sioni	ficance	testing
I dule T.	1 au	COCHICICIU	anu	SIGILI	neance	count

Significant on level test of 5%; ** Significant on level test of 1%

c. Land quality and characteristics that controlling of composite maize productivity

Based on the previous structural model testing, the most influential land quality and control of composite maize productivity based on the order of importance were nutrient retention, root media, land preparation, and available nutrients. This was also based on the results of multiple regression tests with the best equation (equation 1) of the land quality that affects composite maize production were:

 $Y = 5.892 + 0.430X_1 + 0.453X_2 + 0.248X_3 - 0.443X_4 \dots (1)$ r = 0.56

Where: X_1 = root medium, X_2 = nutrient retention, X_3 = nutrients availability, X_4 = land preparation. Furthermore, the land characteristics that most influenced the productivity of composite maize based on the order of importance were pH KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. This was also based on the results of multiple regression tests with the best equation (equation 2) as follows:

 $Y = 4.531 + 0.450X_1 - 0.351X_2 - 0.365X_3 + 0.321X_4 - 0.352X_5 + 0.351X_6 + 0.337X_1... (2)$ r = 0.63

Where: $X_1 = pH$ of KCl, $X_2 = coarse$ material, $X_3 = rock$ outcrop, $X_4 = effective$ depth, $X_5 = surface rock$, $X_6 = K$ available, $X_7 = soil$ texture.

The relationship of each land characteristic and its contribution to land quality in influencing the composite maize productivity was presented in Table 5 and Figure 3. The land characteristics consisting of texture, effective depth, pH of KCI, and available K has a positive relationship and significant effect on the composite maize productivity. This shows that the increasing of the land characteristics by 1%, the composite maize productivity will increase by 30% to 47%. In contrast, the content of coarse material, surface rock, and rock outcrops has a negative relationship and significant effect on the composite maize productivity. This indicates that the decreasing content of coarse material, surface rock, and rock outcrops was 1% each in line with the increase in the composite maize productivity by 42% to 44%. The correlation of each of these land characteristics was quite strong in influencing the composite maize productivity. Coarse material is rock fragments measuring 2 mm in diameter or more which affect soil moisture, infiltration, erosion, and land use [20].

Coarse material <15% is very suitable for maize, while > 55% is not suitable [21] [22] [23]. The most suitable soil texture for maize is a fine or loamy texture [24]. Meanwhile, the deeper effective depth affects root growth and development, so that plants can grow and develop well [25]. Surface rocks and rock outcrops are limiting factors in the suitability of maize land in Saentis Village [26].

	1 1 2	
Coefficient of correlation	Contribution on land quality (%)	Coefficient of correlation
Temperature $(X_{1.1})$	0.127	0.20
Rainfall $(X_{2.1})$	0.279	17.2
Wet months $(X_{2.2})$	0.209	13.7
Dry months $(X_{2.3})$	-0.124	-13.2
$LGP(X_{2.4})$	0.166	12.2
Drainage $(X_{3.1})$	0.084	14.1
Texture $(X_{4.1})$	0.298*	18.4
Coarse material (X _{4.2})	-0.438**	-89.4
Effective depth $(X_{4.3})$	0.431**	76.1
pH H ₂ O (X _{5.1})	0.254	32.0
pH KCl (X _{5.2})	0.471**	43.2
C-Organic $(X_{5.3})$	0.264	41.5
$CEC(X_{5.4})$	0.123	24.7
Base saturation $(X_{5.5})$	0.216	47.3
N Total $(X_{6.1})$	0.158	46.7
Available P (X _{6.2})	0.012	33.2
Available $K(X_{6.3})$	0.368*	77.5
$\text{ESP}(X_{7.1})$	-0.024	-17.1
Slope $(X_{8.1})$	-0.266	-44.4
Soil erosion $(X_{8.2})$	-0.158	-28.3
Inundation height $(X_{9.1})$	0.014	23.1
Inundation period (X _{9.2})	0.010	20.1
Surface rock $(X_{10.1})$	-0.418**	-83.7
Rock outcrop $(X_{10,2})$	-0.436**	-85.0

 Table 5. Coefficient of correlation and contribution level on land quality of the land characteristics and composite maize productivity

*Significant on level test of 5%; ** Significant on level test of 1%.

CONCLUSION

Land quality that controls the productivity of composite maize based on the order of importance is nutrient retention, root media, land preparation, and available nutrients. Meanwhile, land characteristics that control the productivity of composite maize based on the order of importance are pH of KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH of KCI, and available K had a positive and significant effect on the productivity of composite maize, while the content of coarse material, surface rock, and rock outcrops had a negative relationship and had a significant effect on the productivity of composite maize.

ACKNOWLEDGMENT

Thank you to the Brawijaya University, which provided PNBP research funding for the 2019 budget year. Thanks also went to Ecan Adam, SE, MM and Rival Rahman, SP, MSi for helping with SEM-PLS analysis and mapping of land resources.

REFERENCES

- [1] BPS RI. 2019. Maize production by province (tons), 1993-2018. Central Bureau of Statistics of the Republic of Indonesia, Jakarta.
- [2] Yasin, H. G. M., W. Langgo, dan Faesal. 2014. White seed maize as an alternative staple food. J. Food Crops Science and Technology **9**(2): 108 117.
- [3] BPS Kabupaten Gorontalo. 2019. Gorontalo regency in 2019 figures. Gorontalo Regencyt Statistics Agency, Limboto.
- [4] Zakaria, A. K. 2011. Anticipatory policy and farmers consolidating strategy toward national maize self-sufficiency. J. Agricultural Policy Analysis **9**(3): 261-274.
- [5] Bahtiar and B. Kumontoi. 2015. Challenges of composite maize seed production in Bolaang Mongondow regency, North Sulawesi. Proceedings of the National Cereals Seminar 2015: 596 - 604.
- [6] Murni, A. M and R. W. Arief. 2008. Maize cultivation technology. Center for Agricultural Studies and Development, Bogor.
- [7] Swastika, D. K. S. 2002. Maize self sufficiency in Indonesia: The past 30 years and future prospect. J. Indonesian Agricultural Research and Development **21**(3): 75-83.
- [8] Subardja, D and Sudarsono. 2005. Effect of land quality on maize productivity in volcanic soils and sedimentary rocks in the Bogor area. J. Soils and Climate **23**: 38-47.
- [9] FAO. 1976. A Framework for land evaluation. Food and Agriculture Organization Soil Bull. No. 32, Rome.
- [10] Subardja, D. 2005. Land suitability criteria for land use types based on maize and peanuts in the Bogor area. Dissertation. IPB Postgraduate School, Bogor.
- [11] Syaf, H. 2014. Evaluation of the relationship between land quality, growth and production of aged cocoa in East Kolaka Regency, Southeast Sulawesi Province. J. Bioeducation 3(1): 267 – 276.
- [12] Elisanti A. D, W. Purnomo and S. Melaniani. 2013. Implementation of partial least square health status of children under five in Indonesia. J. Biometrics and Population 2(2): 99 107.
- [13] Ulum, M., I. M. Tirta, and D. Anggraeni. 2014. Analysis of structural equation modeling (SEM) for small samples using the partial least square (PLS) approach. Proceedings of the National Mathematics Seminar, University of Jember, 19 November 2014: 1 – 15.
- [14] Jaya, I. G. N. M and I. M. Sumertajaya. 2008. Structural equation modeling with partial least square. Proceedings of the National Seminar on Mathematics and Mathematics Education 2008: 119 – 132.
- [15] Hair, J. F., Ringle, C. M. and M. Sarstedt. 2013. Partial least squares structural equation modeling: Rigorous applications, better results and higher acceptance. J. Long Range Planning 46: 1 - 12.
- [16] Igbaria, M., N. Zinatelli, P. Cragg, and A. L. M. Cavaye. 1997. Personal computing acceptance factors in small firms: a structural equation model. J. MIS Quarterly 21(3): 279-305.
- [17] Mattjik, A. A and I. M. Sumertajaya. 2011. Investigate multiple variables using SAS. IPB Press, Bogor.
- [18] Sujarweni, V. W. 2014. SPSS for research. Pustaka Baru, Yogyakarta.
- [19] Abdilah, W., and J. Hartono. 2015. Partial least square (PLS): alternative structural equation modeling (SEM) in business research. Andi Offset, Yogyakarta.
- [20] Soil Research Institute. 2004. Technical guidelines for soil observation. Research and Development Center for Soil and Agro-climate, Bogor.

- [21] Djaenudin D, Marwan H, Subagjo H, and A. Hidayat. 2011. Technical guidelines for land evaluation for agricultural commodities. Center for R&D for agricultural land resources, Agricultural R&D Agency, Bogor. 36p.
- [22] Ritung S, K Nugroho, A Mulyani, and E Suryani. 2012. Technical guidelines for land evaluation for agricultural commodities. Center for Research and Development of Land Resources, Agricultural Research and Development Agency, Bogor. 166 p.
- [23] Wahyunto, Hikmatullah, E. Suryani, C. Tafakresnanto, S. Ritung, A. Mulyani, Sukarman, K. Nugroho, Y. Sulaeman, Y. Apriyana, Suciantini, A. Pramudia, Suparto, RE Subandiono, T. Sutriadi, D. Nursyamsi. 2016. Technical guidelines for land suitability assessment guidelines for semi-detailed strategic agricultural commodities at a scale of 1: 50,000. Center for Research and Development of Agricultural Land Resources, Agricultural Research and Development Agency, Bogor. 37 p.
- [24] Sudjana A, A. Rifin and M. Sudjadi. 1991. Maize. Agricultural research and development agency, Bogor.
- [25] Wirosoedarmo R, A. T. Sutanhaji, E. Kurniati and R. Wijayanti. 2011. Evaluation of land suitability for maize using spatial analysis methods. J. Agritech **31**(1): 71-78.
- [26] Elfayetti and Herdi. 2015. Evaluation of land suitability for maize crops in Saentis Village, Percut Sei Tuan. J Social Sciences Education 7(1): 33-40.



Figure 1. Map of the Research Location



Figure 3. Path coefficient diagram of land quality to productivity level of composite maize

INSTRUKSI REVISI

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Nurdin^{1*}, M. L Rayes², Soemarno², Sudarto²

¹Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 ³Soil Science Departmen, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 *E-mail: <u>nurdin@ung.ac.id</u>

ABSTRACT

The challenge of composite maize developing in the future is the low productivity because the maize is grown on land that is not suitable for land quality. This study aims to determine the land quality and land characteristics that control the composite maize productivity in Gorontalo Province. A total of 33 land units were surveyed and their land observed to obtain data on morphology and soil characteristics, climate and terrain characteristics, as well as composite maize productivity data through ubinan plots and direct interviews with maize farmers. Partial least square of structural equation models (PLS-SEM) analysis has been used to determine the land quality and land characteristics that control the composite maize productivity through variable validity and reliability tests, as well as structural model tests. The results showed that the manifest variables were air temperature, rainfall, wet months, dry months, LGP, drainage, coarse materials, effective depth, pH ${
m H_2O}$, pH KCl, C-organic, total N, available P, available K, ESP, slopes, soil erosion, inundation height, inundation time, surface rock, and rock outcrops were valid and able to explain well the latent variables. Furthermore, the latent variables were temperature, water availability, oxygen availability, nutrient retention, nutrients availability, sodicity, erosion hazard, flood hazard, and land preparation used has good composite reliability and high reliability because of the composite reliability and alpha cronbach >0.6. Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of course materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

INTRODUCTION

Low maize productivity is still a major problem in efforts to increase maize production in Indonesia. Until 2018, maize productivity had only reached 5.24 tons/ha [1], while the potential for maize productivity in Indonesia could reach 10-11 tons/ha [2]. Even though the government has rolled out various programs to increase maize production and productivity in order to achieve national maize selfsufficiency.

Gorontalo Province is one of the maizes producing centers in Indonesia with an average maize productivity achievement until 2019 of 5.03 tons/ha [3] or still far below the average national maize productivity. All this time, maize farmers has been more dominant in planting maize with hybrid and composite varieties. There are no references to the productivity of hybrid or composite maize in this area, so it is assumed that the maize productivity achievements are relatively the same at 5.24 tons/ ha. In fact, specifically the productivity of composite maize can reach 5-6 tons/ha [4] [5] or as much as 8.5 tons/ha [6]. Composite maize, besides its productivity potential is relatively similar to that of hybrid maize, it also has advantages, including being more adaptive in acid soils [5] and can be used as seeds for the next growing season, while hybrid maize can't be planted again. The use of composite maize can reduce the dependence of maize farmers on hybrid maize seed assistance from the government, so that if the maize seed subsidy is stopped, the farmers can plant the composite maize again.

Keywords: Quality, characteristic, land, productivity, maize, composite.

Correspondence:

Nurdin

Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 E-mail: nurdin@ung.ac.id

The challenge ahead in developing composite maize is the low productivity of composite maize, so it is necessary to address the root of the problem. Maize planted on land with low productivity potential is one of the causes for the low productivity of maize [7]. Meanwhile, land characteristics and quality have a close relationship with maize productivity [8] and each land quality has a significant effect on land suitability for certain uses [9], especially for maize. Research on land quality that controls the productivity of composite maize has been conducted in the Bogor area using stepwise regression analysis [10]. The use of structural equation modeling (SEM) analysis in determining the characteristics and quality of land that control plant productivity has not been widely published, except for [11] who used SEM analysis on older cocoa plants in Kolaka Timur Regency, Southeast Sulawesi Province. Meanwhile, the use of SEM analysis specifically to determine the relationship between land quality and maize productivity has not been found. The response of maize plants to the diversity of

characteristics and quality of land will vary, so it is important to know the quality and characteristics of the land that control the productivity of maize. The diversity of characteristics and complex quality of land in the field really requires a comprehensive analysis technique that is able to simplify the complexity in one analysis system. One analysis option is to use SEM analysis. SEM analysis is able to analyze how much influence each indicator (manifest) of soil physical and chemical properties (latent) has on productivity in one analysis unit [11]. The use of SEM is very helpful to determine

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

the effect on indicators and to produce a model that is better than other multivariate analyzes [12] [13]. Partial Least Square (PLS) is a variant of SEM which has a higher level of flexibility because PLS is based on variants, so that the number of samples used does not need to be large, ranges from 30-100, and does not require normal multivariate assumptions compared to CB-SEM. requires a large data sample size (> 100) and the data must be multivariate normal distribution [14] [15]. Therefore, a research on land quality that controls the productivity of composite maize was carried out using SEM-PLS analysis based on the consideration of complex land characteristics and quality, as well as limited data in the land unit in the study area. The purpose of this study was to determine the quality and characteristics of land that control the productivity of composite maize in Gorontalo.

MATERIALS AND METHOD

This research is located in the Sustainable Agriculture Area of Gorontalo Province (Figure 1) and the Soil Laboratory of the Department of Soil, Faculty of Agriculture, Brawijaya University. The timing of this research was started in December 2019 - May 2020. The tools used included the computer, SmatPLS version 2.0, SPSS, Microsoft Excel, and Microsoft Word. While the materials studied included the morphological data and soil characteristics, climate and terrain characteristics data that had been grouped into their respective land qualities, as well as composite maize productivity data from the study area.

Soil surveys and land observations were carried out to obtain morphological data and soil characteristics, climate and terrain characteristics data from the research area. Meanwhile, composite maize productivity data was obtained from the results of ubinan directly on the land of maize farmers and from direct interviews with maize farmers on 33 land units. Furthermore, the diversity of sizes and data units (ratio data) of land characteristics were converted in the form of interval data which were represented as follows were 1 (very low), 2 (low), 3 (medium), 4 (high), and 5 (very high). After the data is ready, the analysis process is continued using SEM-PLS (Figure 2).



Figure 2. Research Operational Framework

The latent variable in this study was the quality of the land consisting of temperature (X_1) , water availability (X_2) , oxygen availability (X_3) , root media (X_4) , nutrient retention (X_5) , available nutrients (X₆), sodicity (X₇), erosion hazard (X₈), flood hazard (X₉), and land preparation (X₁₀). While the manifest variable was the characteristic of the land which consists of temperature $(X_{1,1})$, rainfall $(X_{2,1})$, wet months (X_{2.2}), dry months (X_{2.3}), LGP (X_{2.4}), drainage. (X_{3.1}), texture (X_{4.1}), coarse material (X_{4.2}), effective soil depth (X_{4.3}), pH H₂O (X_{5.1}), pH KCI (X_{5.2}), C-organic (X_{5.3}), CEC (X_{5.4}), base saturation (X_{5.5}), total N (X_{6.1}), available P (X_{6.2}), K available (X_{6.3}), ESP (X_{7.1}), slope (X_{8.1}), soil erosion (X_{8.2}), inundation height (X_{9.1}), length of inundation (X_{9.2}), surface rock (X_{10.1}), and rock outcrop (X10.2). The use of SEM-PLS in this study consisted of testing the validity, reliability of the research variables, and testing the structural model. In summary, the test using SEM-PLS was described as follows:

a. Testing the Validity of Research Variables. The basic evaluation carried out in the SEM-PLS analysis is to evaluate the measurement model (outer model) with the aim of knowing the validity and reliability of indicators in measuring research latent variables through convergent validity, discriminant validity, and composite reliability. Convergent validity testing on SEM-PLS is seen from the size of the outer loading of each indicator on its latent variable. A loading factor value above 0.70 is highly recommended, but a loading factor value of 0.50-0.60 can still be tolerated with a tstatistic value of more than 1.96 or a small p-value of 0.05. The loading factor of an indicator with the highest value is the strongest or most important measure in reflecting the latent variable in question. Discriminant validity is an evaluation of the outer model in SEM-PLS using cross loading values to test valid and reliable indicators in explaining or reflecting latent variables. If the correlation of the latent variable with the measurement core of each indicator is greater than the other latent variables, then the latent variable is able to predict the indicator better than other latent variables and is said to be valid.

- b. Research Variable Reliability Testing. Composite reliability and alpha cronbach were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is> 0.60.
- c. Structural Model Testing. Testing of the structural model (*inner model*) is carried out after the relationship model is built in accordance with the observed data and the suitability of the overall model (goodness of fit model). Testing of structural models and hypotheses is carried out by looking at the estimated value of the path coefficient and the critical point value (t-statistic) which is significant at $\alpha = 0.05$. Testing the relationship model and hypothesis between variables can be done by testing the direct correlation coefficient between variables. The results of testing the relationship between the X variables and the Y variable in this study are shown by the correlation coefficient and t-statistic, and also seen in the path diagram.

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

RESULTS AND DISCUSSION

a. Research Variable Validity
The loading factor value of the research variables, where the loading factor value on the indicators was mostly more than the critical limit of 0.70 with a confidence level of 95% (Table 1). The value of the loading factor which is below the tolerance value of 0.5 at the 95% confidence level where the t-statistic value of each indicator is smaller than the t-table

(1.960) is found in the soil texture indicator of the latent variable root media (X₄) which is only 0.173 is also the CEC indicator (X_{5.4}) and the base saturation indicator (X_{5.5}) of the nutrient retention latent variable (X₅), which are only 0.399 and 0.482 respectively. This means that these indicators have not been able to properly form or explain their latent variables.

Table 1. Outer loading research variables

Effect of indi	Loading factors	Status		
Temperature (X _{1.1})	->	Temperature (X ₁)	1.000	Valid
Rainfall (X _{2.1})	->		0.981	Valid
Wet months (X _{2.2})	->	Water engile tite (V)	0.989	Valid
Dry months (X _{2.3})	->	water availability (A2)	0.827	Valid
LGP (X _{2.4})	->		0.968	Valid
Drainage (X _{3.1})	->	Oxygen availability (X ₃)	1.000	Valid
Texture (X _{4.1})	->		0.173	Not valid
Coarse material (X _{4.2})	->	Rooting media (X4)	-0.921	Valid
Effective depth (X _{4.3})	->		0.912	Valid
pH H ₂ O (X _{5.1})	->		0.768	Valid
pH KCI (X _{5.2})	->	Nutrient retention (X5)	0.772	Valid
C-Organic (X _{5.3})	->		0.710	Valid
CEC (X _{5.4})	->		0.399	Not valid
Base saturation (X _{5.5})	->		0.482	Not valid
N Total (X _{6.1})	->		0.799	Valid
Available P (X _{6.2})	->	Nutrient availability (X ₆)	0.521	Valid
Available K (X _{6.3})	->		0.886	Valid
ESP (X _{7.1})	->	Sodicity (X7)	1.000	Valid
Slope (X _{8.1})	->	Fracian hazard (Ya)	0.974	Valid
Soil erosion (X _{8.2})	->	El OSION NAZALU (A8)	0.957	Valid
Inundation height (X _{9.1})	->	Elooding hazard (Va)	0.993	Valid
Inundation period (X9.2)	->	Floounig flazar u (X9)	0.991	Valid
Surface rock (X _{10.1})	->	I and propagation (Vec)	0.998	Valid
Rock outcrop (X _{10.2})	->	Lanu preparation (A10)	0.998	Valid
Productivity (Y _{1.1})	->	Composite maize productivity (Y1)	1.000	Valid

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Indikator	Temperature (X1)	Water availability (X2)	Oxygen availability (X ₃)	Rooting media (X4)	Nutrient retention (X5)	Nutrient availability (X ₆)	Sodicity (X7)	Erosion hazard (X8)	Flooding hazard (X9)	Land preparation (X10)	Composite maize productivity (Y1)
Air temperature (X _{1.1})	1	0.952309	0.059098	-0.08736	-0.37805	-0.06653	0.38176	0.016269	-0.10297	0.19833	0.042282
Rainfall (X _{2.1})	0.968555	0.980906	0.114576	0.052348	-0.24379	0.058536	0.356547	-0.0379	-0.04621	0.056015	0.156751
Wet months (X _{2.2})	0.926635	0.989185	0.173659	-0.005903	-0.25644	0.062873	0.374745	-0.06373	-0.04367	0.060342	0.177251
Dry months (X _{2.3})	0.759123	0.82697	0.141078	-0.238735	-0.42612	-0.10563	0.47553	-0.11715	0.027746	0.215367	0.076041
LGP (X _{2.4})	0.900431	0.96821	0.13569	-0.003834	-0.28223	0.056251	0.459669	-0.12209	-0.04398	0.059938	0.193991
Drainage (X _{3.1})	0.059098	0.144225	1	0.129338	-0.24128	0.057861	0.084339	-0.50344	0.236555	-0.22277	0.400657
Texture (X _{4.1})	-0.02057	-0.01261	-0.16957	0.172551	0.242032	0.12283	0.217308	0.196875	-0.00074	-0.02261	0.09248
Coarse material (X _{4.2})	-0.00333	-0.1005	-0.13244	-0.921096	-0.38256	-0.6112	0.18822	0.322934	-0.26391	0.846957	-0.35202
Effective depth (X _{4.3})	-0.17758	-0.09256	0.165016	0.912088	0.3519	0.355112	-0.23141	-0.19005	0.095721	-0.76736	0.180089
pH H2O (X5.1)	-0.40346	-0.38437	-0.3719	0.29356	0.767791	0.27088	-0.17175	0.151553	-0.02966	-0.08478	0.186569
pH KCl (X _{5.2})	-0.25953	-0.22811	-0.44804	0.342269	0.771872	0.272936	-0.02729	0.167533	0.098977	-0.18312	0.268161
C-Organic (X _{5.3})	-0.29516	-0.13852	0.096529	0.248076	0.710022	0.612498	0.073184	-0.4692	0.063874	-0.1793	0.384332
CEC (X5.4)	0.066756	0.115697	0.003345	0.084182	0.399393	0.421251	0.373179	-0.05735	0.15285	-0.01387	0.281455
Base saturation (X _{5.5})	-0.30026	-0.25724	-0.10527	0.412102	0.481624	0.361795	-0.60079	-0.0895	-0.13592	-0.48759	0.136266
N Total (X _{6.1})	0.002878	0.137879	0.07154	0.268606	0.545283	0.798694	0.030267	-0.37884	-0.10212	-0.2485	0.427705
Available P (X _{6.2})	-0.09821	-0.09791	-0.44547	0.211821	0.409315	0.520984	-0.28705	-0.057	0.033581	-0.26033	-0.02547
Available K(X _{6.3})	-0.09732	-0.01031	0.06693	0.614343	0.51245	0.885686	-0.3292	-0.29441	0.237691	-0.6422	0.49531
ESP (X _{7.1})	0.38176	0.405078	0.084339	-0.186069	-0.06947	-0.21259	1	-0.01035	0.201152	0.361936	-0.0249
Slope (X _{8.1})	-0.02207	-0.12714	-0.51717	-0.295103	-0.1643	-0.40295	-0.03466	0.973779	-0.34215	0.324431	-0.64795
Soil erosion (X _{8.2})	0.064136	-0.00224	-0.44709	-0.166166	-0.11161	-0.32907	0.021581	0.956588	-0.12926	0.257787	-0.48649
Inundation height (X _{9.1})	-0.08956	-0.02635	0.225421	0.194354	0.082178	0.127762	0.193925	-0.26735	0.992798	-0.13415	0.175472
Inundation period (X _{9.2})	-0.11594	-0.06329	0.244833	0.199427	0.048584	0.078386	0.205739	-0.2425	0.991369	-0.11616	0.135302
Surface rock (X _{10.1})	0.212772	0.074279	-0.23401	-0.854273	-0.28568	-0.55023	0.376036	0.319248	-0.13208	0.997623	-0.28655
Rock outcrop (X _{10.2})	0.183196	0.051703	-0.21067	-0.868319	-0.29655	-0.55537	0.34638	0.290608	-0.12053	0.997697	-0.28228
Productivity (Y _{1.1})	0.042282	0.177277	0.400657	0.304774	0.418519	0.534535	-0.0249	-0.59733	0.157534	-0.28507	1

Table 2. Cross loading of research variables

LGP: Long growth period; CEC : Cation exchange capacity; ESP: Excangeable potassium percentage

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

b.

The standard of loading factor is greater equal to 0.50 [16] [17] [13]. However, in general, based on the indicated values, it can be concluded that the latent variables of land quality have been able to be well established or explained by each indicator and can be said to be convergent valid on these indicators. The cross-loading value for the indicators of latent variables on average is above the cross-loading value of the indicators for other latent variables (Table 2). That is, the greatest cross loading value on the indicator is found in the latent variable too, except for the texture indicator (X_{4.1}) of the root media variable (X₄), the CEC indicator (X_{5.4}) and base saturation (X_{5.5}) of the nutrient retention variable (X₅) whose cross loading value is still smaller (<0.5) than the cross loading value of other latent variables. The standard of loading factor is ≥0.50 [16] [17] [13]. Thus, the indicators of each latent variable are mostly able to explain the latent variable itself better than the other variables, so that the research variables are said to be discriminant valid.

Reliability of Research Variables

Composite reliability and Cronbach alpha were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is above 0.60 [18]. The composite reliability value on each research variable is more than the limit value (>0.6), except for the root media variable (Table 3). The composite reliability value and the Cronbach alpha value is greater than 0.6 so that the latent variable has good composite reliability and high reliability. A construct is said to be reliable if the Cronbach Alpha value must be >0.6 [19]. Thus, all indicators used in this study have met the criteria or are feasible to be used in the measurement of all latent variables because they have good validity and high reliability. The results of the evaluation of convergent validity and discriminant validity of indicators or variables as well as composite reliability and alpha Cronbach for indicators or variables can be concluded that the indicators as measures of latent variables are valid and reliable measures respectively.

Table 3. Composite reliability and Cronbach's Alpha values of research variables

Laten variables	Composite reliability	Alpha Cronbach
Temperature (X ₁)	1.000000	1.000000
Water availability (X ₂)	0.970030	0.965126
Oxygen availability (X ₃)	1.000000	1.000000
Rooting media (X ₄)	0.020314	-1.055192
Nutrient retention (X ₅)	0.770518	0.628062
Nutrient availability (X ₆)	0.788289	0.681393
Sodicity (X7)	1.000000	1.000000
Erosion hazard (X ₈)	0.964615	0.927731
Flooding hazard (X ₉)	0.992053	0.984010
Land preparation (X ₁₀)	0.997657	0.995304

b. Structural Model Testing

The structural model (inner model) is evaluated by looking at the coefficient value of the relationship path parameter between latent variables. It seems that the soil quality of the root media, nutrient retention, and available nutrients showed a positive correlation and had a significant effect on the productivity of composite maize (Table 4). The quality of land preparation shows a negative correlation and has a significant effect on the productivity of composite maize. This indicates that the better rooting media, available nutrient and nutrient retention and a decrease in the level of land preparation as the productivity of composite maize increases. The results of this study are slightly different from the research report [8] regarding the quality of soil rooting media which has not affected the productivity of maize in the Bogor area, but the quality of soil nutrient retention and available nutrients has a significant effect on the productivity of maize relatively the same as the results of this study.

Table 4. Path coefficient and significance testing

	Endogenous variables				
Exogenous variables	Composite maize produktivity (Y)				
	Path coeffisient	t-statistics (t _{critics} = 2.00)			
Temperature (X ₁)	0.086	1.531			
Water availability (X ₂)	0.457	-0.491			
Oxygen availability (X ₃)	0.099	0.371			
Rooting media (X ₄)	0.091*	2.250			
Nutrient retention (X ₅)	0.740*	2.291			
Nutrient availability (X ₆)	0.283**	6.509			
Sodicity (X ₇)	-0.194	-0.036			
Erosion hazard (X ₈)	-0.043	-1.043			
Flooding hazard (X ₉)	0.050	-0.050			
Land preparation (X ₁₀)	-0.386*	-2.339			

Significant on level test of 5%; ** Significant on level test of 1%

c. Land quality and characteristics that controlling of composite maize productivity

Based on the previous structural model testing, the most influential land quality and control of composite maize productivity based on the order of importance were nutrient retention, root media, land preparation, and available nutrients. This was also based on the results of multiple regression tests with the best equation (equation 1) of the land quality that affects composite maize productivity were: $Y = 5.892 + 0.430X_1 + 0.453X_2 + 0.248X_3 - 0.443X_4$ (1) r = 0.56

Where: X_1 = root medium, X_2 = nutrient retention, X_3 = nutrients availability, X_4 = land preparation. Furthermore, the land characteristics that most influenced the productivity of

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

composite maize based on the order of importance were pH KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. This was also based on the results of multiple regression tests with the best equation (equation 2) as follows:

$Y = 4.531 + 0.450X_1 - 0.351X_2 - 0.365X_3 + 0.321X_4 - 0.321X_4$	0.352X5
+ 0.351X ₆ + 0.337X ₁	(2)
	r = 0.63

Where: $X_1 = pH$ of KCl, $X_2 = coarse material$, $X_3 = rock outcrop$, $X_4 = effective depth$, $X_5 = surface rock$, $X_6 = K$ available, $X_7 = soil texture$.

The relationship of each land characteristic and its contribution to land quality in influencing the composite maize productivity was presented in Table 5 and Figure 3. The land characteristics consisting of texture, effective depth, pH of KCI, and available K has a positive relationship and significant effect on the composite maize productivity. This shows that the increasing of the land characteristics by 1%, the composite maize productivity will increase by 30% to

47%. In contrast, the content of coarse material, surface rock, and rock outcrops has a negative relationship and significant effect on the composite maize productivity. This indicates that the decreasing content of coarse material, surface rock, and rock outcrops was 1% each in line with the increase in the composite maize productivity by 42% to 44%. The correlation of each of these land characteristics was quite strong in influencing the composite maize productivity. Coarse material is rock fragments measuring 2 mm in diameter or more which affect soil moisture, infiltration, erosion, and land use [20]. Coarse material <15% is very suitable for maize, while > 55% is not suitable [21] [22] [23]. The most suitable soil texture for maize is a fine or loamy texture [24]. Meanwhile, the deeper effective depth affects root growth and development, so that plants can grow and develop well [25]. Surface rocks and rock outcrops are limiting factors in the suitability of maize land in Saentis Village [26].

Table 5. Coefficient of correlation and contribution level on land quality of the land characteristics and composite maize productivity

Coefficient of correlation	Contribution on land quality (%)	Coefficient of correlation
Temperature (X _{1.1})	0.127	0.20
Rainfall (X _{2.1})	0.279	17.2
Wet months (X _{2.2})	0.209	13.7
Dry months (X _{2.3})	-0.124	-13.2
LGP (X _{2.4})	0.166	12.2
Drainage (X _{3.1})	0.084	14.1
Texture (X _{4.1})	0.298*	18.4
Coarse material (X _{4.2})	-0.438**	-89.4
Effective depth (X _{4.3})	0.431**	76.1
pH H ₂ O (X _{5.1})	0.254	32.0
pH KCl (X5.2)	0.471**	43.2
C-Organic (X _{5.3})	0.264	41.5
CEC (X _{5.4})	0.123	24.7
Base saturation (X _{5.5})	0.216	47.3
N Total (X _{6.1})	0.158	46.7
Available P (X _{6.2})	0.012	33.2
Available K(X _{6.3})	0.368*	77.5
ESP (X7.1)	-0.024	-17.1
Slope (X _{8.1})	-0.266	-44.4
Soil erosion (X _{8.2})	-0.158	-28.3
Inundation height (X _{9.1})	0.014	23.1
Inundation period (X _{9.2})	0.010	20.1
Surface rock (X _{10.1})	-0.418**	-83.7
Rock outcrop (X _{10.2})	-0.436**	-85.0

*Significant on level test of 5%; ** Significant on level test of 1%.

CONCLUSION

Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of course materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

ACKNOWLEDGMENT

Thank you to the Brawijaya University, which provided PNBP research funding for the 2019 budget year. Thanks also went to Ecan Adam, SE, MM and Rival Rahman, SP, MSi for helping with SEM-PLS analysis and mapping of land resources.

REFERENCES

- 1. BPS RI. 2019. Maize production by province (tons), 1993-2018. Central Bureau of Statistics of the Republic of Indonesia, Jakarta.
- 2. Yasin, H. G. M., W. Langgo, dan Faesal. 2014. White seed maize as an alternative staple food. J. Food Crops Science and Technology **9**(2): 108 117.
- 3. BPS Kabupaten Gorontalo. 2019. Gorontalo regency in 2019 figures. Gorontalo Regencyt Statistics Agency, Limboto.

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

- 4. Zakaria, A. K. 2011. Anticipatory policy and farmers consolidating strategy toward national maize self-sufficiency. J. Agricultural Policy Analysis **9**(3): 261-274.
- 5. Bahtiar and B. Kumontoi. 2015. Challenges of composite maize seed production in Bolaang Mongondow regency, North Sulawesi. Proceedings of the National Cereals Seminar 2015: 596 - 604.
- 6. Murni, A. M and R. W. Arief. 2008. Maize cultivation technology. Center for Agricultural Studies and Development, Bogor.
- Swastika, D. K. S. 2002. Maize self-sufficiency in Indonesia: The past 30 years and future prospect. J. Indonesian Agricultural Research and Development 21(3): 75-83.
- 8. Subardja, D and Sudarsono. 2005. Effect of land quality on maize productivity in volcanic soils and sedimentary rocks in the Bogor area. J. Soils and Climate **23**: 38-47.
- 9. FAO. 1976. A Framework for land evaluation. Food and Agriculture Organization Soil Bull. No. 32, Rome.
- Subardja, D. 2005. Land suitability criteria for land use types based on maize and peanuts in the Bogor area. Dissertation. IPB Postgraduate School, Bogor.
- Syaf, H. 2014. Evaluation of the relationship between land quality, growth and production of aged cocoa in East Kolaka Regency, Southeast Sulawesi Province. J. Bioeducation 3(1): 267 – 276.
- Elisanti A. D, W. Purnomo and S. Melaniani. 2013. Implementation of partial least square health status of children under five in Indonesia. J. Biometrics and Population 2(2): 99 – 107.
- Ulum, M., I. M. Tirta, and D. Anggraeni. 2014. Analysis of structural equation modeling (SEM) for small samples using the partial least square (PLS) approach. Proceedings of the National Mathematics Seminar, University of Jember, 19 November 2014: 1 – 15.
- 14. Jaya, I. G. N. M and I. M. Sumertajaya. 2008. Structural equation modeling with partial least square. Proceedings of the National Seminar on Mathematics and Mathematics Education 2008: 119 – 132.
- 15. Hair, J. F., Ringle, C. M. and M. Sarstedt. 2013. Partial least squares structural equation modeling: Rigorous applications, better results and higher acceptance. J. Long Range Planning **46**: 1 - 12.
- Igbaria, M., N. Zinatelli, P. Cragg, and A. L. M. Cavaye. 1997. Personal computing acceptance factors in small firms: a structural equation model. J. MIS Quarterly 21(3): 279 – 305.
- 17. Mattjik, A. A and I. M. Sumertajaya. 2011. Investigate multiple variables using SAS. IPB Press, Bogor.
- 18. Sujarweni, V. W. 2014. SPSS for research. Pustaka Baru, Yogyakarta.
- 19. Abdilah, W., and J. Hartono. 2015. Partial least square (PLS): alternative structural equation modeling (SEM) in business research. Andi Offset, Yogyakarta.
- 20. Soil Research Institute. 2004. Technical guidelines for soil observation. Research and Development Center for Soil and Agro-climate, Bogor.
- 21. Djaenudin D, Marwan H, Subagjo H, and A. Hidayat. 2011. Technical guidelines for land evaluation for agricultural commodities. Center for R&D for agricultural land resources, Agricultural R&D Agency, Bogor. 36p.
- 22. Ritung S, K Nugroho, A Mulyani, and E Suryani. 2012. Technical guidelines for land evaluation for agricultural commodities. Center for Research and Development of Land Resources, Agricultural Research and Development Agency, Bogor. 166 p.

- 23. Wahyunto, Hikmatullah, E. Suryani, C. Tafakresnanto, S. Ritung, A. Mulyani, Sukarman, K. Nugroho, Y. Sulaeman, Y. Apriyana, Suciantini, A. Pramudia, Suparto, RE Subandiono, T. Sutriadi, D. Nursyamsi. 2016. Technical guidelines for land suitability assessment guidelines for semi-detailed strategic agricultural commodities at a scale of 1: 50,000. Center for Research and Development of Agricultural Land Resources, Agricultural Research and Development Agency, Bogor. 37 p.
- 24. Sudjana A, A. Rifin and M. Sudjadi. 1991. Maize. Agricultural research and development agency, Bogor.
- Wirosoedarmo R, A. T. Sutanhaji, E. Kurniati and R. Wijayanti. 2011. Evaluation of land suitability for maize using spatial analysis methods. J. Agritech **31**(1): 71-78.
- 26. Elfayetti and Herdi. 2015. Evaluation of land suitability for maize crops in Saentis Village, Percut Sei Tuan. J Social Sciences Education **7**(1): 33-40.

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maize Varieties in Gorontalo



Figure 1. Map of the Research Location

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Figure 3. Path coefficient diagram of land quality to productivity level of composite maize varieties

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Nurdin^{1*}, M. L Rayes², Soemarno², Sudarto²

¹Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 ³Soil Science Departmen, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 *E-mail: <u>nurdin@ung.ac.id</u>

ABSTRACT

The challenge of composite maize developing in the future is the low productivity because the maize is grown on land that is not suitable for land quality. This study aims to determine the land quality and land characteristics that control the composite maize productivity in Gorontalo Province. A total of 33 land units were surveyed and their land observed to obtain data on morphology and soil characteristics, climate and terrain characteristics, as well as composite maize productivity data through ubinan plots and direct interviews with maize farmers. Partial least square of structural equation models (PLS-SEM) analysis has been used to determine the land quality and land characteristics that control the composite maize productivity through variable validity and reliability tests, as well as structural model tests. The results showed that the manifest variables were air temperature, rainfall, wet months, dry months, LGP, drainage, coarse materials, effective depth, pH H₂O, pH KCl, C-organic, total N, available P, available K, ESP, slopes, soil erosion, inundation height, inundation time, surface rock, and rock outcrops were valid and able to explain well the latent variables. Furthermore, the latent variables were temperature, water availability, oxygen availability, nutrient retention, nutrients availability, sodicity, erosion hazard, flood hazard, and land preparation used has good composite reliability and high reliability because of the composite reliability and alpha cronbach >0.6. Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of course materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

INTRODUCTION

Low maize productivity is still a major problem in efforts to increase maize production in Indonesia. Until 2018, maize productivity had only reached 5.24 tons/ha [1], while the potential for maize productivity in Indonesia could reach 10-11 tons/ha [2]. Even though the government has rolled out various programs to increase maize production and productivity in order to achieve national maize selfsufficiency.

Gorontalo Province is one of the maizes producing centers in Indonesia with an average maize productivity achievement until 2019 of 5.03 tons/ha [3] or still far below the average national maize productivity. All this time, maize farmers has been more dominant in planting maize with hybrid and composite varieties. There are no references to the productivity of hybrid or composite maize in this area, so it is assumed that the maize productivity achievements are relatively the same at 5.24 tons/ ha. In fact, specifically the productivity of composite maize can reach 5-6 tons/ha [4] [5] or as much as 8.5 tons/ha [6]. Composite maize, besides its productivity potential is relatively similar to that of hybrid maize, it also has advantages, including being more adaptive in acid soils [5] and can be used as seeds for the next growing season, while hybrid maize can't be planted again. The use of composite maize can reduce the dependence of maize farmers on hybrid maize seed assistance from the government, so that if the maize seed subsidy is stopped, the farmers can plant the composite maize again.

Keywords: Quality, characteristic, land, productivity, maize, composite.

Correspondence:

Nurdin

Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 E-mail: nurdin@ung.ac.id

The challenge ahead in developing composite maize is the low productivity of composite maize, so it is necessary to address the root of the problem. Maize planted on land with low productivity potential is one of the causes for the low productivity of maize [7]. Meanwhile, land characteristics and quality have a close relationship with maize productivity [8] and each land quality has a significant effect on land suitability for certain uses [9], especially for maize. Research on land quality that controls the productivity of composite maize has been conducted in the Bogor area using stepwise regression analysis [10]. The use of structural equation modeling (SEM) analysis in determining the characteristics and quality of land that control plant productivity has not been widely published, except for [11] who used SEM analysis on older cocoa plants in Kolaka Timur Regency, Southeast Sulawesi Province. Meanwhile, the use of SEM analysis specifically to determine the relationship between land quality and maize productivity has not been found. The response of maize plants to the diversity of

characteristics and quality of land will vary, so it is important to know the quality and characteristics of the land that control the productivity of maize. The diversity of characteristics and complex quality of land in the field really requires a comprehensive analysis technique that is able to simplify the complexity in one analysis system. One analysis option is to use SEM analysis. SEM analysis is able to analyze how much influence each indicator (manifest) of soil physical and chemical properties (latent) has on productivity in one analysis unit [11]. The use of SEM is very helpful to determine

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

the effect on indicators and to produce a model that is better than other multivariate analyzes [12] [13]. Partial Least Square (PLS) is a variant of SEM which has a higher level of flexibility because PLS is based on variants, so that the number of samples used does not need to be large, ranges from 30-100, and does not require normal multivariate assumptions compared to CB-SEM. requires a large data sample size (> 100) and the data must be multivariate normal distribution [14] [15]. Therefore, a research on land quality that controls the productivity of composite maize was carried out using SEM-PLS analysis based on the consideration of complex land characteristics and quality, as well as limited data in the land unit in the study area. The purpose of this study was to determine the quality and characteristics of land that control the productivity of composite maize in Gorontalo.

MATERIALS AND METHOD

This research is located in the Sustainable Agriculture Area of Gorontalo Province (Figure 1) and the Soil Laboratory of the Department of Soil, Faculty of Agriculture, Brawijaya University. The timing of this research was started in December 2019 - May 2020. The tools used included the computer, SmatPLS version 2.0, SPSS, Microsoft Excel, and Microsoft Word. While the materials studied included the morphological data and soil characteristics, climate and terrain characteristics data that had been grouped into their respective land qualities, as well as composite maize productivity data from the study area.

Soil surveys and land observations were carried out to obtain morphological data and soil characteristics, climate and terrain characteristics data from the research area. Meanwhile, composite maize productivity data was obtained from the results of ubinan directly on the land of maize farmers and from direct interviews with maize farmers on 33 land units. Furthermore, the diversity of sizes and data units (ratio data) of land characteristics were converted in the form of interval data which were represented as follows were 1 (very low), 2 (low), 3 (medium), 4 (high), and 5 (very high). After the data is ready, the analysis process is continued using SEM-PLS (Figure 2).

Figure 2. Research Operational Framework

The latent variable in this study was the quality of the land consisting of temperature (X_1) , water availability (X_2) , oxygen availability (X_3) , root media (X_4) , nutrient retention (X_5) , available nutrients (X₆), sodicity (X₇), erosion hazard (X₈), flood hazard (X₉), and land preparation (X₁₀). While the manifest variable was the characteristic of the land which consists of temperature $(X_{1,1})$, rainfall $(X_{2,1})$, wet months (X_{2.2}), dry months (X_{2.3}), LGP (X_{2.4}), drainage. (X_{3.1}), texture (X_{4.1}), coarse material (X_{4.2}), effective soil depth (X_{4.3}), pH H₂O (X_{5.1}), pH KCI (X_{5.2}), C-organic (X_{5.3}), CEC (X_{5.4}), base saturation (X_{5.5}), total N (X_{6.1}), available P (X_{6.2}), K available (X_{6.3}), ESP (X_{7.1}), slope (X_{8.1}), soil erosion (X_{8.2}), inundation height (X_{9.1}), length of inundation (X_{9.2}), surface rock (X_{10.1}), and rock outcrop (X10.2). The use of SEM-PLS in this study consisted of testing the validity, reliability of the research variables, and testing the structural model. In summary, the test using SEM-PLS was described as follows:

a. Testing the Validity of Research Variables. The basic evaluation carried out in the SEM-PLS analysis is to evaluate the measurement model (outer model) with the aim of knowing the validity and reliability of indicators in measuring research latent variables through convergent validity, discriminant validity, and composite reliability. Convergent validity testing on SEM-PLS is seen from the size of the outer loading of each indicator on its latent variable. A loading factor value above 0.70 is highly recommended, but a loading factor value of 0.50-0.60 can still be tolerated with a tstatistic value of more than 1.96 or a small p-value of 0.05. The loading factor of an indicator with the highest value is the strongest or most important measure in reflecting the latent variable in question. Discriminant validity is an evaluation of the outer model in SEM-PLS using cross loading values to test valid and reliable indicators in explaining or reflecting latent variables. If the correlation of the latent variable with the measurement core of each indicator is greater than the other latent variables, then the latent variable is able to predict the indicator better than other latent variables and is said to be valid.

- b. Research Variable Reliability Testing. Composite reliability and alpha cronbach were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is> 0.60.
- c. Structural Model Testing. Testing of the structural model (*inner model*) is carried out after the relationship model is built in accordance with the observed data and the suitability of the overall model (goodness of fit model). Testing of structural models and hypotheses is carried out by looking at the estimated value of the path coefficient and the critical point value (t-statistic) which is significant at $\alpha = 0.05$. Testing the relationship model and hypothesis between variables can be done by testing the direct correlation coefficient between variables. The results of testing the relationship between the X variables and the Y variable in this study are shown by the correlation coefficient and t-statistic, and also seen in the path diagram.

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

RESULTS AND DISCUSSION

a. Research Variable Validity
The loading factor value of the research variables, where the loading factor value on the indicators was mostly more than the critical limit of 0.70 with a confidence level of 95% (Table 1). The value of the loading factor which is below the tolerance value of 0.5 at the 95% confidence level where the t-statistic value of each indicator is smaller than the t-table

(1.960) is found in the soil texture indicator of the latent variable root media (X₄) which is only 0.173 is also the CEC indicator (X_{5.4}) and the base saturation indicator (X_{5.5}) of the nutrient retention latent variable (X₅), which are only 0.399 and 0.482 respectively. This means that these indicators have not been able to properly form or explain their latent variables.

Table 1. Outer loading research variables

Effect of indicators on latent variables			Loading factors	Status
Temperature (X _{1.1})	->	Temperature (X ₁)	1.000	Valid
Rainfall (X _{2.1})	->		0.981	Valid
Wet months (X _{2.2})	->	Water engile tite (V)	0.989	Valid
Dry months (X _{2.3})	->	water availability (X ₂)	0.827	Valid
LGP (X _{2.4})	->		0.968	Valid
Drainage (X _{3.1})	->	Oxygen availability (X ₃)	1.000	Valid
Texture (X _{4.1})	->		0.173	Not valid
Coarse material (X _{4.2})	->	Rooting media (X4)	-0.921	Valid
Effective depth (X _{4.3})	->		0.912	Valid
pH H ₂ O (X _{5.1})	->		0.768	Valid
pH KCI (X _{5.2})	->		0.772	Valid
C-Organic (X _{5.3})	->	Nutrient retention (X ₅)	0.710	Valid
CEC (X _{5.4})	->		0.399	Not valid
Base saturation ($X_{5.5}$)	->		0.482	Not valid
N Total (X _{6.1})	->		0.799	Valid
Available P (X _{6.2})	->	Nutrient availability (X ₆)	0.521	Valid
Available K (X _{6.3})	->		0.886	Valid
ESP (X _{7.1})	->	Sodicity (X7)	1.000	Valid
Slope (X _{8.1})	->	Fracian hazard (Ya)	0.974	Valid
Soil erosion (X _{8.2})	->		0.957	Valid
Inundation height (X _{9.1})	->	Elooding bazard (Ye)	0.993	Valid
Inundation period (X _{9.2})	->	rioouing nazai u (xy)	0.991	Valid
Surface rock (X _{10.1})	->	I and proparation (Yes)	0.998	Valid
Rock outcrop (X _{10.2})	->	במות או לאמו מנוסוו (גוס)	0.998	Valid
Productivity (Y _{1.1})	->	Composite maize productivity (Y1)	1.000	Valid

Indikator	Temperature (X1)	Water availability (X2)	Oxygen availability (X ₃)	Rooting media (X4)	Nutrient retention (X5)	Nutrient availability (X6)	Sodicity (X7)	Erosion hazard (X8)	Flooding hazard (X9)	Land preparation (X10)	Composite maize productivity (Y1)
Air temperature (X _{1.1})	1	0.952309	0.059098	-0.08736	-0.37805	-0.06653	0.38176	0.016269	-0.10297	0.19833	0.042282
Rainfall (X _{2.1})	0.968555	0.980906	0.114576	0.052348	-0.24379	0.058536	0.356547	-0.0379	-0.04621	0.056015	0.156751
Wet months (X _{2.2})	0.926635	0.989185	0.173659	-0.005903	-0.25644	0.062873	0.374745	-0.06373	-0.04367	0.060342	0.177251
Dry months (X _{2.3})	0.759123	0.82697	0.141078	-0.238735	-0.42612	-0.10563	0.47553	-0.11715	0.027746	0.215367	0.076041
LGP (X _{2.4})	0.900431	0.96821	0.13569	-0.003834	-0.28223	0.056251	0.459669	-0.12209	-0.04398	0.059938	0.193991
Drainage (X _{3.1})	0.059098	0.144225	1	0.129338	-0.24128	0.057861	0.084339	-0.50344	0.236555	-0.22277	0.400657
Texture (X _{4.1})	-0.02057	-0.01261	-0.16957	0.172551	0.242032	0.12283	0.217308	0.196875	-0.00074	-0.02261	0.09248
Coarse material (X _{4.2})	-0.00333	-0.1005	-0.13244	-0.921096	-0.38256	-0.6112	0.18822	0.322934	-0.26391	0.846957	-0.35202
Effective depth (X _{4.3})	-0.17758	-0.09256	0.165016	0.912088	0.3519	0.355112	-0.23141	-0.19005	0.095721	-0.76736	0.180089
pH H ₂ O (X _{5.1})	-0.40346	-0.38437	-0.3719	0.29356	0.767791	0.27088	-0.17175	0.151553	-0.02966	-0.08478	0.186569
pH KCl (X _{5.2})	-0.25953	-0.22811	-0.44804	0.342269	0.771872	0.272936	-0.02729	0.167533	0.098977	-0.18312	0.268161
C-Organic (X _{5.3})	-0.29516	-0.13852	0.096529	0.248076	0.710022	0.612498	0.073184	-0.4692	0.063874	-0.1793	0.384332
CEC (X _{5.4})	0.066756	0.115697	0.003345	0.084182	0.399393	0.421251	0.373179	-0.05735	0.15285	-0.01387	0.281455
Base saturation (X _{5.5})	-0.30026	-0.25724	-0.10527	0.412102	0.481624	0.361795	-0.60079	-0.0895	-0.13592	-0.48759	0.136266
N Total (X _{6.1})	0.002878	0.137879	0.07154	0.268606	0.545283	0.798694	0.030267	-0.37884	-0.10212	-0.2485	0.427705
Available P (X _{6.2})	-0.09821	-0.09791	-0.44547	0.211821	0.409315	0.520984	-0.28705	-0.057	0.033581	-0.26033	-0.02547
Available K(X _{6.3})	-0.09732	-0.01031	0.06693	0.614343	0.51245	0.885686	-0.3292	-0.29441	0.237691	-0.6422	0.49531
ESP (X _{7.1})	0.38176	0.405078	0.084339	-0.186069	-0.06947	-0.21259	1	-0.01035	0.201152	0.361936	-0.0249
Slope (X _{8.1})	-0.02207	-0.12714	-0.51717	-0.295103	-0.1643	-0.40295	-0.03466	0.973779	-0.34215	0.324431	-0.64795
Soil erosion (X _{8.2})	0.064136	-0.00224	-0.44709	-0.166166	-0.11161	-0.32907	0.021581	0.956588	-0.12926	0.257787	-0.48649
Inundation height (X _{9.1})	-0.08956	-0.02635	0.225421	0.194354	0.082178	0.127762	0.193925	-0.26735	0.992798	-0.13415	0.175472
Inundation period (X _{9.2})	-0.11594	-0.06329	0.244833	0.199427	0.048584	0.078386	0.205739	-0.2425	0.991369	-0.11616	0.135302
Surface rock (X _{10.1})	0.212772	0.074279	-0.23401	-0.854273	-0.28568	-0.55023	0.376036	0.319248	-0.13208	0.997623	-0.28655
Rock outcrop (X _{10.2})	0.183196	0.051703	-0.21067	-0.868319	-0.29655	-0.55537	0.34638	0.290608	-0.12053	0.997697	-0.28228
Productivity (Y _{1.1})	0.042282	0.177277	0.400657	0.304774	0.418519	0.534535	-0.0249	-0.59733	0.157534	-0.28507	1

Table 2. Cross loading of research variables

LGP: Long growth period; CEC : Cation exchange capacity; ESP: Excangeable potassium percentage

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

The standard of loading factor is greater equal to 0.50 [16] [17] [13]. However, in general, based on the indicated values, it can be concluded that the latent variables of land quality have been able to be well established or explained by each indicator and can be said to be convergent valid on these indicators. The cross-loading value for the indicators of latent variables on average is above the cross-loading value of the indicators for other latent variables (Table 2). That is, the greatest cross loading value on the indicator is found in the latent variable too, except for the texture indicator (X_{4.1}) of the root media variable (X₄), the CEC indicator (X_{5.4}) and base saturation (X_{5.5}) of the nutrient retention variable (X₅) whose cross loading value is still smaller (<0.5) than the cross loading value of other latent variables. The standard of loading factor is ≥0.50 [16] [17] [13]. Thus, the indicators of each latent variable are mostly able to explain the latent variable itself better than the other variables, so that the research variables are said to be discriminant valid.

b. Reliability of Research Variables

Composite reliability and Cronbach alpha were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is above 0.60 [18]. The composite reliability value on each research variable is more than the limit value (>0.6), except for the root media variable (Table 3). The composite reliability value and the Cronbach alpha value is greater than 0.6 so that the latent variable has good composite reliability and high reliability. A construct is said to be reliable if the Cronbach Alpha value must be >0.6 [19]. Thus, all indicators used in this study have met the criteria or are feasible to be used in the measurement of all latent variables because they have good validity and high reliability. The results of the evaluation of convergent validity and discriminant validity of indicators or variables as well as composite reliability and alpha Cronbach for indicators or variables can be concluded that the indicators as measures of latent variables are valid and reliable measures respectively.

Laten variables	Composite reliability	Alpha Cronbach
Temperature (X1)	1.000000	1.000000
Water availability (X ₂)	0.970030	0.965126
Oxygen availability (X ₃)	1.000000	1.000000
Rooting media (X ₄)	0.020314	-1.055192
Nutrient retention (X ₅)	0.770518	0.628062
Nutrient availability (X ₆)	0.788289	0.681393
Sodicity (X7)	1.000000	1.000000
Erosion hazard (X ₈)	0.964615	0.927731
Flooding hazard (X ₉)	0.992053	0.984010
Land preparation (X ₁₀)	0.997657	0.995304

b. Structural Model Testing

The structural model (inner model) is evaluated by looking at the coefficient value of the relationship path parameter between latent variables. It seems that the soil quality of the root media, nutrient retention, and available nutrients showed a positive correlation and had a significant effect on the productivity of composite maize (Table 4). The quality of land preparation shows a negative correlation and has a significant effect on the productivity of composite maize. This indicates that the better rooting media, available nutrient and nutrient retention and a decrease in the level of land preparation as the productivity of composite maize increases. The results of this study are slightly different from the research report [8] regarding the quality of soil rooting media which has not affected the productivity of maize in the Bogor area, but the quality of soil nutrient retention and available nutrients has a significant effect on the productivity of maize relatively the same as the results of this study.

Table 4.	Path	coefficient	and	signific	ance	testing
rubic ii	I uuii	coefficient	unu	Signific	ance	cesting

	Endogenous variables			
Exogenous variables	Composite maize produktivity (Y)			
	Path coeffisient	t-statistics ($t_{critics} = 2.00$)		
Temperature (X ₁)	0.086	1.531		
Water availability (X ₂)	0.457	-0.491		
Oxygen availability (X ₃)	0.099	0.371		
Rooting media (X ₄)	0.091*	2.250		
Nutrient retention (X ₅)	0.740*	2.291		
Nutrient availability (X ₆)	0.283**	6.509		
Sodicity (X7)	-0.194	-0.036		
Erosion hazard (X ₈)	-0.043	-1.043		
Flooding hazard (X ₉)	0.050	-0.050		
Land preparation (X ₁₀)	-0.386*	-2.339		

Significant on level test of 5%; ** Significant on level test of 1%

c. Land quality and characteristics that controlling of composite maize productivity

Based on the previous structural model testing, the most influential land quality and control of composite maize productivity based on the order of importance were nutrient retention, root media, land preparation, and available nutrients. This was also based on the results of multiple regression tests with the best equation (equation 1) of the land quality that affects composite maize productivity were: $Y = 5.892 + 0.430X_1 + 0.453X_2 + 0.248X_3 - 0.443X_4 \dots (1)$ r = 0.56

Where: X_1 = root medium, X_2 = nutrient retention, X_3 = nutrients availability, X_4 = land preparation. Furthermore, the land characteristics that most influenced the productivity of

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

composite maize based on the order of importance were pH KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. This was also based on the results of multiple regression tests with the best equation (equation 2) as follows:

$Y = 4.531 + 0.450X_1 - 0.351X_2 - 0.365X_3 + 0.321X_4 -$	0.352X5
+ 0.351X ₆ + 0.337X ₁	(2)
	r = 0.63

Where: $X_1 = pH$ of KCl, $X_2 = coarse material$, $X_3 = rock outcrop$, $X_4 = effective depth$, $X_5 = surface rock$, $X_6 = K$ available, $X_7 = soil texture$.

The relationship of each land characteristic and its contribution to land quality in influencing the composite maize productivity was presented in Table 5 and Figure 3. The land characteristics consisting of texture, effective depth, pH of KCI, and available K has a positive relationship and significant effect on the composite maize productivity. This shows that the increasing of the land characteristics by 1%, the composite maize productivity will increase by 30% to

47%. In contrast, the content of coarse material, surface rock, and rock outcrops has a negative relationship and significant effect on the composite maize productivity. This indicates that the decreasing content of coarse material, surface rock, and rock outcrops was 1% each in line with the increase in the composite maize productivity by 42% to 44%. The correlation of each of these land characteristics was quite strong in influencing the composite maize productivity. Coarse material is rock fragments measuring 2 mm in diameter or more which affect soil moisture, infiltration, erosion, and land use [20]. Coarse material <15% is very suitable for maize, while > 55% is not suitable [21] [22] [23]. The most suitable soil texture for maize is a fine or loamy texture [24]. Meanwhile, the deeper effective depth affects root growth and development, so that plants can grow and develop well [25]. Surface rocks and rock outcrops are limiting factors in the suitability of maize land in Saentis Village [26].

Table 5. Coefficient of correlation and contribution level on land quality of the land characteristics and composite maize productivity

Coefficient of correlation	Contribution on land quality (%)	Coefficient of correlation
Temperature (X _{1.1})	0.127	0.20
Rainfall (X _{2.1})	0.279	17.2
Wet months (X _{2.2})	0.209	13.7
Dry months (X _{2.3})	-0.124	-13.2
LGP (X _{2.4})	0.166	12.2
Drainage (X _{3.1})	0.084	14.1
Texture (X _{4.1})	0.298*	18.4
Coarse material (X _{4.2})	-0.438**	-89.4
Effective depth (X _{4.3})	0.431**	76.1
pH H ₂ O (X _{5.1})	0.254	32.0
pH KCl (X5.2)	0.471**	43.2
C-Organic (X _{5.3})	0.264	41.5
CEC (X _{5.4})	0.123	24.7
Base saturation (X _{5.5})	0.216	47.3
N Total (X _{6.1})	0.158	46.7
Available P (X _{6.2})	0.012	33.2
Available K(X _{6.3})	0.368*	77.5
ESP (X _{7.1})	-0.024	-17.1
Slope (X _{8.1})	-0.266	-44.4
Soil erosion (X _{8.2})	-0.158	-28.3
Inundation height (X _{9.1})	0.014	23.1
Inundation period (X _{9.2})	0.010	20.1
Surface rock (X _{10.1})	-0.418**	-83.7
Rock outcrop (X _{10.2})	-0.436**	-85.0

*Significant on level test of 5%; ** Significant on level test of 1%.

CONCLUSION

Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of course materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

ACKNOWLEDGMENT

Thank you to the Brawijaya University, which provided PNBP research funding for the 2019 budget year. Thanks also went to Ecan Adam, SE, MM and Rival Rahman, SP, MSi for helping with SEM-PLS analysis and mapping of land resources.

REFERENCES

- 1. BPS RI. 2019. Maize production by province (tons), 1993-2018. Central Bureau of Statistics of the Republic of Indonesia, Jakarta.
- 2. Yasin, H. G. M., W. Langgo, dan Faesal. 2014. White seed maize as an alternative staple food. J. Food Crops Science and Technology **9**(2): 108 117.
- 3. BPS Kabupaten Gorontalo. 2019. Gorontalo regency in 2019 figures. Gorontalo Regencyt Statistics Agency, Limboto.

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maíze Varieties in Gorontalo

- Zakaria, A. K. 2011. Anticipatory policy and farmers consolidating strategy toward national maize selfsufficiency. J. Agricultural Policy Analysis 9(3): 261-274.
- Bahtiar and B. Kumontoi. 2015. Challenges of composite maize seed production in Bolaang Mongondow regency, North Sulawesi. Proceedings of the National Cereals Seminar 2015: 596 - 604.
- 6. Murni, A. M and R. W. Arief. 2008. Maize cultivation technology. Center for Agricultural Studies and Development, Bogor.
- Swastika, D. K. S. 2002. Maize self-sufficiency in Indonesia: The past 30 years and future prospect. J. Indonesian Agricultural Research and Development 21(3): 75-83.
- 8. Subardja, D and Sudarsono. 2005. Effect of land quality on maize productivity in volcanic soils and sedimentary rocks in the Bogor area. J. Soils and Climate **23**: 38-47.
- 9. FAO. 1976. A Framework for land evaluation. Food and Agriculture Organization Soil Bull. No. 32, Rome.
- 10. Subardja, D. 2005. Land suitability criteria for land use types based on maize and peanuts in the Bogor area. Dissertation. IPB Postgraduate School, Bogor.
- Syaf, H. 2014. Evaluation of the relationship between land quality, growth and production of aged cocoa in East Kolaka Regency, Southeast Sulawesi Province. J. Bioeducation 3(1): 267 – 276.
- Elisanti A. D, W. Purnomo and S. Melaniani. 2013. Implementation of partial least square health status of children under five in Indonesia. J. Biometrics and Population 2(2): 99 – 107.
- Ulum, M., I. M. Tirta, and D. Anggraeni. 2014. Analysis of structural equation modeling (SEM) for small samples using the partial least square (PLS) approach. Proceedings of the National Mathematics Seminar, University of Jember, 19 November 2014: 1 – 15.
- 14. Jaya, I. G. N. M and I. M. Sumertajaya. 2008. Structural equation modeling with partial least square. Proceedings of the National Seminar on Mathematics and Mathematics Education 2008: 119 132.
- 15. Hair, J. F., Ringle, C. M. and M. Sarstedt. 2013. Partial least squares structural equation modeling: Rigorous applications, better results and higher acceptance. J. Long Range Planning **46**: 1 12.
- Igbaria, M., N. Zinatelli, P. Cragg, and A. L. M. Cavaye. 1997. Personal computing acceptance factors in small firms: a structural equation model. J. MIS Quarterly 21(3): 279 – 305.
- 17. Mattjik, A. A and I. M. Sumertajaya. 2011. Investigate multiple variables using SAS. IPB Press, Bogor.
- 18. Sujarweni, V. W. 2014. SPSS for research. Pustaka Baru, Yogyakarta.
- 19. Abdilah, W., and J. Hartono. 2015. Partial least square (PLS): alternative structural equation modeling (SEM) in business research. Andi Offset, Yogyakarta.
- 20. Soil Research Institute. 2004. Technical guidelines for soil observation. Research and Development Center for Soil and Agro-climate, Bogor.
- Djaenudin D, Marwan H, Subagjo H, and A. Hidayat. 2011. Technical guidelines for land evaluation for agricultural commodities. Center for R&D for agricultural land resources, Agricultural R&D Agency, Bogor. 36p.
- 22. Ritung S, K Nugroho, A Mulyani, and E Suryani. 2012. Technical guidelines for land evaluation for agricultural commodities. Center for Research and Development of Land Resources, Agricultural Research and Development Agency, Bogor. 166 p.

- 23. Wahyunto, Hikmatullah, E. Suryani, C. Tafakresnanto, S. Ritung, A. Mulyani, Sukarman, K. Nugroho, Y. Sulaeman, Y. Apriyana, Suciantini, A. Pramudia, Suparto, RE Subandiono, T. Sutriadi, D. Nursyamsi. 2016. Technical guidelines for land suitability assessment guidelines for semi-detailed strategic agricultural commodities at a scale of 1: 50,000. Center for Research and Development of Agricultural Land Resources, Agricultural Research and Development Agency, Bogor. 37 p.
- 24. Sudjana A, A. Rifin and M. Sudjadi. 1991. Maize. Agricultural research and development agency, Bogor.
- 25. Wirosoedarmo R, A. T. Sutanhaji, E. Kurniati and R. Wijayanti. 2011. Evaluation of land suitability for maize using spatial analysis methods. J. Agritech **31**(1): 71-78.
- 26. Elfayetti and Herdi. 2015. Evaluation of land suitability for maize crops in Saentis Village, Percut Sei Tuan. J Social Sciences Education **7**(1): 33-40.

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Figure 1. Map of the Research Location

Study of Land Quality and Land Characterístics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Figure 3. Path coefficient diagram of land quality to productivity level of composite maize varieties

15 November, 2020 SRY_1289_ November -2020

Dear Authors

Nurdin¹*, M. L Rayes², Soemarno³, Sudarto⁴

¹Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145

^{2,3,4} Soil Science Departmen, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145

Email: nurdin@ung.ac.id

We would like to inform you that your manuscript id SRP-October 1289 has been accepted for publication in **SYSTEMATIC REVIEWS IN PHARMACY (ISSN 0975-8453).** Your paper will be published in Volume 12 Issue 1 of 2021.

TITLE: Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Thanks for submission of your work with us.

Regards, Professor Dr. Lucius,

Systematic Reviews in Pharmacy Pharmaceutical Science best quartile SJR 2018 0.3 powered by scimagojr.com

Associate Editor SYSTEMATIC REVIEWS IN PHARMACY (ISSN 0975-8453) http://www.sysrevpharm.org/

PUBLISH

COLUMN STREET, STREET,

WWW.SVST

al and the same

ALC: NOT THE OWNER.

Systematic Reviews Pharmacy

Andre S. Marten I. Control of Con

2021, Vol: 12, Issue: 2

Current Issue (http://www.sysrevpharm.org//?sec=cissue)	12 / 2
Online First (http://www.sysrevpharm.org//?sec=aip)	
Archive (http://www.sysrevpharm.org//?sec=archive)	
Aims and Scope (http://www.sysrevpharm.org//?sec=aimsscope)	
Abstracting & Indexing (http://www.sysrevpharm.org//?sec=jindex)	
Most Accessed Articles (http://www.sysrevpharm.org//?sec=mosta)	
Most Downloaded Articles (http://www.sysrevpharm.org//?sec=mostd)	
Most Cited Articles (http://www.sysrevpharm.org//?sec=mostc)	

Systematic Reviews in Pharmacy (Sys Rev Pharm.), (SRP) (Print ISSN: 0975-8453, E-ISSN: 0976-2779) a halfyearly publication, serves the need of different scientists and others involved in Pharmaceutical research and development. Each issue covers review articles on Drug discovery topics, and also publishes full-length reviews related to different subjects in pharmacy and that are of broad readership interest to users in industry, academia, and government. The first issue was published online in December 2009.

All contributions to Sys. Rev Pharm are reviewed by the peer review process and copyediting process with the understanding that they have not been published previously and are not under consideration for publication elsewhere. Author/s is/are responsible for all statements made in their work and obtaining necessary permission to republish any previously published illustrations and/or other relevant materials. The journal's full text is available online at www.sysrevpharm.org (http://www.sysrevpharm.org/). The journal allows free access (Open Access) to its contents and permits authors to self-archive the final accepted version of the articles on any OAI-compliant institutional / subject-based repository.

Scope of the journal

The journal covers and publishes **all articles** related to Pharmacy subjects including some of the allied subjects. Articles with timely interest and newer research concepts will be given more preference.

Systematic Reviews in Pharmacy

This journal also publishes manuscripts related to agriculture and agriculture sciences. Topics include all aspects of crop and animal physiology, modeling of crop and animal systems, the scientific underpinning of agronomy and husbandry, animal welfare and behavior, soil science, plant and animal product quality, plant and animal nutrition, medical law, law and education, engineering solutions, decision support systems, land use, environmental impacts of agriculture and forestry, impacts of climate change, rural biodiversity, experimental design and statistical analysis, and the application of new analytical and study methods (including genetic diversity and molecular biology approaches).

(https://www.scimagojr.com/journalsearch.php?

q=19700201140&tip=sid&exact=no)

2021, Vol: 12, Issue: 2

« Previous Issue (index.php?iid=2021-12-1.000&jid=196)

SRP. Year: 2021, Volume: 12

Systematic Review

1. Contagious Viral Diseases in the Operating Room: A Review of Evolving Safety Protocols at Three Levels Li Zihan

SRP. 2021; 12(2): 1-9

» Abstract (?mno=135420) » PDF (index.php?fulltxt=135420&fulltxtj=196&fulltxtp=196-1600235518.pdf) » doi: 10.31838/srp.2021.2.1 (http://dx.doi.org/10.31838/srp.2021.2.1)

2. Chinese, European and Persian Medicine Recommendations for the Prevention of Pandemics (COVID-19) Related to the Respiratory System: A Systematic Review

Yingwen ZHAO

SRP. 2021; 12(2): 10-23

```
» Abstract (?mno=135421) » PDF (index.php?fulltxt=135421&fulltxtj=196&fulltxtp=196-1600235656.pdf) » doi: 10.31838/srp.2021.2.2 (http://dx.doi.org/10.31838/srp.2021.2.2)
```

3. Proline Rich Antimicrobial Peptides-----A tool to Counter Antimicrobial Resistance

```
Amjid lqbal

SRP. 2021; 12(2): 24-31

» Abstract (?mno=135423) » PDF (index.php?fulltxt=135423&fulltxtj=196&fulltxtp=196-1600236112.pdf) » doi:

10.31838/srp.2021.2.3 (http://dx.doi.org/10.31838/srp.2021.2.3)
```

And A COMPANY

ISVS WWW

al and the same

ALC: NOT THE OWNER.

Systematic Reviews Pharmacy

Andre de Andre i Connes y Connes de Andre de Andre i Connes de Barrier Andre de Andre i Connes de Andre de Andre i Connes de Andre i Connes de Andre I Californi Connes de Andre Connes de Andre i Californi de Andre i Connes de Andre i Californi de Andre i Connes de Andre i Californi de Andre i Andre i Californi de Andre i C

2021, Vol: 12, Issue: 2

Current Issue (http://www.sysrevpharm.org//?sec=cissue)	12 / 2
Online First (http://www.sysrevpharm.org//?sec=aip)	
Archive (http://www.sysrevpharm.org//?sec=archive)	
Aims and Scope (http://www.sysrevpharm.org//?sec=aimsscope)	
Abstracting & Indexing (http://www.sysrevpharm.org//?sec=jindex)	
Most Accessed Articles (http://www.sysrevpharm.org//?sec=mosta)	
Most Downloaded Articles (http://www.sysrevpharm.org//?sec=mostd)	
Most Cited Articles (http://www.sysrevpharm.org//?sec=mostc)	
ORCID (https://orcid.org/register) Crossref (https://www.crossref.org/)	

« Previous Issue (index.php?iid=2020-11-11.000&jid=196)

Next Issue » (index.php?iid=2021-12-1.000&jid=196)

SRP. Year: 2020, Volume: 11, Issue: 12

Review Article

1. Attachment Differences of Candida albicans Colonies on Self Cured and Heat Cured Acrylic Resin Plates Used as Removable Orthodontic Appliances Donald R. Nahusona, Rika Damayanti Syarif, Nurlina Nurdin SRP. 2020; 11(12): 1-5 » Abstract (?mno=126838) » PDF (index.php?fulltxt=126838&fulltxtj=196&fulltxtp=196-1597863363.pdf) » doi: 10.31838/srp.2020.12.1 (http://dx.doi.org/10.31838/srp.2020.12.1) 2. Chlorella Vulgaris Extract Cream as A Formulation of Bone Remodelling Edy Machmud, Nina Permatasari

SRP. 2020; 11(12): 6-15

» Abstract (?mno=127092) » PDF (index.php?fulltxt=127092&fulltxtj=196&fulltxtp=196-1597921104.pdf) » doi: 10.31838/srp.2020.12.2 (http://dx.doi.org/10.31838/srp.2020.12.2)

3. The Hard Tissue Lesions Associated with Impacted Teeth	
Dwi Putri Wulansari, Irfan Sugianto, Muliaty Yunus, Rafikah Hasyim, Nasyrah Hidayati, Sri Naca Hardiana	
SRP. 2020; 11(12): 16-18	
» Abstract (?mno=127098) » PDF (index.php?fulltxt=127098&fulltxtj=196&fulltxtp=196-1597921426.pdf)	» doi:
10.31838/srp.2020.12.3 (http://dx.doi.org/10.31838/srp.2020.12.3)	
4. The Relationship between Service Quality, Culture Similarity to Satisfaction and Loyalty of Medical (Dentation and Loyalty of Medical (Dent	al)
Tourism	
Fuad Husain Akbar, Burhanuddin Daeng Pasiga, Rasmidar Samad, Fridawati Rivai, Andi Zulkifli Abdullah, Abd H	air
Awang, Syahrir A. Pasinringi, Lalu Muhammad Saleh, Irwandy, Dian Pratiwi	
SRP. 2020; 11(12): 19-30	
» Abstract (?mno=127100) » PDF (index.php?fulltxt=127100&fulltxtj=196&fulltxtp=196-1597921573.pdf)	» doi:
10.31838/srp.2020.12.4 (http://dx.doi.org/10.31838/srp.2020.12.4)	
5. The Use of Marine Biota in Bone Tissue Regeneration: A Systematic Review	
Edy Machmud, Muhammad Ikbal, Irfan Dammar, Darmiaty	
SRP. 2020; 11(12): 31-34	
» Abstract (?mno=127104) » PDF (index.php?fulltxt=127104&fulltxtj=196&fulltxtp=196-1597921751.pdf)	» doi:
10.31838/srp.2020.12.5 (http://dx.doi.org/10.31838/srp.2020.12.5)	
6. Implant Coating Materials to Increase Osseointegration of Dental Implant: A Systematic Review	
Eri Hendra Jubhari, Irfan Dammar, Vinsensia Launardo, Yonatan Goan	
SRP. 2020; 11(12): 35-41	
» Abstract (?mno=127107) » PDF (index.php?fulltxt=127107&fulltxtj=196&fulltxtp=196-1597921882.pdf)	» doi:
10.31838/srp.2020.12.6 (http://dx.doi.org/10.31838/srp.2020.12.6)	
7. Role of Green Tea Catechins Toothpaste on Transforming Growth Factor- β 1 (TGF- β 1) and Bone Morphog	enetic-2
(BMP-2) on Early Childhood Caries	
Fajriani1, Sartini, Sherly Horax, Adam Malik, Asmawati, Atika Balqis, Dwi Putri Wulansari	
SRP. 2020; 11(12): 42-47	
» Abstract (?mno=127111) » PDF (index.php?fulltxt=127111&fulltxtj=196&fulltxtp=196-1597922044.pdf)	» doi:
10.31838/srp.2020.12.7 (http://dx.doi.org/10.31838/srp.2020.12.7)	
8. The Rationale For A Method Of Auditing Medical Organizations	
Stasevich N. Y., Basilyeva T. P., Grizanchuk A.M., Latynin E.O.	
SRP. 2020; 11(12): 48-52	
» Abstract (?mno=129976) » PDF (index.php?fulltxt=129976&fulltxtj=196&fulltxtp=196-1598742185.pdf)	» doi:
10.31838/srp.2020.12.8 (http://dx.doi.org/10.31838/srp.2020.12.8)	
9. Clinical and Morphological Characteristics of Gunshot Wound of Soft Tissues	
Julia V. Zbrueva, Dmitry V. Bogomolov	
SRP. 2020; 11(12): 53-56	
» Abstract (?mno=129977) » PDF (index.php?fulltxt=129977&fulltxtj=196&fulltxtp=196-1598742285.pdf)	» doi:
10.31838/srp.2020.12.9 (http://dx.doi.org/10.31838/srp.2020.12.9)	

2/2020	Systematic Reviews in Pharmacy	
10.	. Evaluation of commercial Linked immune-sorbent assay (ELISA) for detecting sero-prevalence of Toxopla	sma
	gondii antibodies in Iraqi women	
	Rasha Amer Noori Al-Tufaili	
	SRP. 2020; 11(12): 57-62	
	» Abstract (?mno=129992) » PDF (index.php?fulltxt=129992&fulltxtj=196&fulltxtp=196-1598756661.pdf)	» doi:
	10.31838/srp.2020.12.10 (http://dx.doi.org/10.31838/srp.2020.12.10)	
11.	Impact of Interleukin-1 beta gene allelic polymorphisms in diabetic and non-diabetic hemodialysis Iragi pa	atients
	Rand Muhammed Abdul-Hussein Al-Husseini	
	SRP. 2020: 11(12): 63-69	
	» Abstract (?mno=129979) » PDF (index php?fulltxt=129979&fulltxti=196&fulltxtp=196-1598742478 pdf)	» doi:
	10.31838/srp 2020 12 11 (http://dx.doi.org/10.31838/srp 2020 12 11)	<i>"</i> doi:
	10.01000/01p.2020.12.11 (http://dx.doi.org/10.01000/01p.2020.12.11)	
12	Isolation. Identification, and Antifungal sensitivity Testing of Epidermophyton floccosum from Clinical Sa	mples
	Raed Ali Hussain Shabaa	
	SRP 2020: 11(12): 70-77	
	» Abstract (2mno=129980) » PDF (index nhn?fulltyt=129980&fulltyti=196&fulltytn=196-1598742701 ndf)	» doi:
	10.31838/srn 2020.12.12.(http://dx.doi.org/10.31838/srn 2020.12.12)	<i>"</i> doi.
	10.51050/stp.2020.12.12 (http://dx.doi.org/10.51050/stp.2020.12.12)	
13	Estimation of Truffles extracts and study the biological efficacy of Ethyl 6-methyl-2-oxo-4-(2-thienyl)-1.2.3	4-
	tetrahydronyrimidine-5-carboxylate (EMOTTC) as one of the extracts against onbthalmic bacteria	•
	Rana D. S. Al kamil. Dawood CH. Al-Babadily. Rasool Chaloob. Falab Hassan Shari, H. N. K. Al -Salman	
	SRP 2020: 11(12): 78-87	
	NAstract (2mno=120081)	» doi:
	= 10.31838/srn 2020 12.13 (http://dv.doi.org/10.31838/srn 2020 12.13)	<i>"</i> doi.
	10.51050/stp.2020.12.15 (http://dx.doi.org/10.51050/stp.2020.12.15)	
14	Assessment Serum Levels of Neopterin, IL-6, IL-18, hs-CRP, TNF-α and MMP 9 in Iraαi Rheumatoid Arthrit	is
• • •	Patients	
	Shakir F T Alaaraji Miid A Mohisen Muthanna M Awad	
	SRP 2020: 11(12): 88-93	
	» Abstract (2mpg=120082) » PDE (index php2fulltyt=120082&fulltyti=106&fulltytn=106_15087/2076 pdf)	» doi:
	10.31838/srp.2020.12.14 (http://dv.doi.org/10.31838/srp.2020.12.14)	<i>"</i> uoi.
	10.51656/sip.2020.12.14 (http://dx.doi.org/10.51656/sip.2020.12.14)	
15	Assessment Serum Levels of Neonterin, II -6, II -18, hs-CRP, TNF-q and MMP 9 in Iraqi Rheumatoid Arthrit	is
10.	Patiente	15
	Shakir E T Alaaraji Mijd A Mohisen Muthanna M Awad	
	» Abstract (2mpg-120000) » PDE (index phg2fulltyt=1200008fulltyti=1068fulltyte=106 1508756280 pdf)	» doi:
	* Abstract (11110-129990) * FDT (Index.php 101111-129990a1011xtj-190a1011xtp-190-1090700200.pdf)	<i>»</i> uoi.
	10.31838/srp.2020.12.14 (http://dx.doi.org/10.31838/srp.2020.12.14)	
16	Statin therapy improvement is a vital necessity for Kazakhstan healtheare	
10.	Altyn A Nurneisova, Raushan Karabayeya, Valoriy V, Ronhorin Larica C, Makalkina	
	CIUTA. HUIPEISUVA, NAUSHAH NATADAYEVA, VAIETTY V. DEHDEHH, LAHSA G. IVIANAINIIA	
	SRF. 2020 , 11(12): 34-30 N Abstract (2mpa=122600) Addit 10.21020/am 2020.10.15 (http://dx.doi.org/10.21020/am 2020.10.15)	
	» Abstract (11110-155009) vul. 10.51858/Stp.2020.12.15 (111p://dx.doi.org/10.31838/Stp.2020.12.15)	
47	Ctatin the same improvement is a witch passagity for Kanal-hatan baskbases	
17.	. Statin therapy improvement is a vital necessity for Nazakinstan nealthcare	

Altyn A. Nurpeisova, Raushan Karabayeva, Valeriy V. Benberin, Larisa G. Makalkina

SRP. 2020; 11(12): 94-98

» Abstract (?mno=133692) » PDF (index.php?fulltxt=133692&fulltxtj=196&fulltxtp=196-1599736408.pdf) » doi: 10.31838/srp.2020.12.15 (http://dx.doi.org/10.31838/srp.2020.12.15)

18. Visualizing impact of Goods and Service Tax on different corners of Pharmaceutical industry- A study of Jaipur District of Rajasthan

Mr. Arun Gautam, Gaurav Lodha, Manish Vadera, Pawan Kumar Verma

SRP. 2020; 11(12): 99-108

» Abstract (?mno=133695) » PDF (index.php?fulltxt=133695&fulltxtj=196&fulltxtp=196-1599736598.pdf) » doi: 10.31838/srp.2020.12.16 (http://dx.doi.org/10.31838/srp.2020.12.16)

19. Detection of Malassezia Species Isolated from Pityriasis Versicolor Infections in An Najaf, Iraq

Raed Ali Hussain Shabaa

SRP. 2020; 11(12): 109-112

» Abstract (?mno=133698) » PDF (index.php?fulltxt=133698&fulltxtj=196&fulltxtp=196-1599736696.pdf) » doi: 10.31838/srp.2020.12.17 (http://dx.doi.org/10.31838/srp.2020.12.17)

20. EPIDEMIOLOGICAL CHARACTERISTICS OF DENGUE AMONG VIETNAMESE CHILDREN IN THE 2017 DENGUE OUTBREAK

Tuan Minh Nguyen, Hung Thanh Nguyen, Minh Ngoc Quang Ngo, Lien Bich Le, Quang Van Pham, Nhan Nguyen Thanh Le, Kinh Van Nguyen, Huy Vu Bui, Thuy Thi Dang, Duyet Van Le

SRP. 2020; 11(12): 113-118

» Abstract (?mno=133700) » PDF (index.php?fulltxt=133700&fulltxtj=196&fulltxtp=196-1599736840.pdf) » doi: 10.31838/srp.2020.12.18 (http://dx.doi.org/10.31838/srp.2020.12.18)

21. Sewage Water Treatment of Chemistry Department in College of Science-Diyala University

Marwah H. Abdulateef, Faihaa F. Hadi, Ahmed Hatem Alwan, Ahmed Najem Abd

SRP. 2020; 11(12): 119-123

» Abstract (?mno=133701) » PDF (index.php?fulltxt=133701&fulltxtj=196&fulltxtp=196-1599736964.pdf) » doi: 10.31838/srp.2020.12.19 (http://dx.doi.org/10.31838/srp.2020.12.19)

22. Quality of Life of Medical University Teachers

Raushangul Magzumova, Indira Bozheyeva, Zaur Mustafayev, Dinara Utepova, Dinara Turzhanova SRP. 2020; 11(12): 124-126

» Abstract (?mno=134329) » PDF (index.php?fulltxt=134329&fulltxtj=196&fulltxtp=196-1599892801.pdf) » doi: 10.31838/srp.2020.12.20 (http://dx.doi.org/10.31838/srp.2020.12.20)

23. Clean and Healthy Living Behavior with The Stunting Events in Children in Central Java, Indonesia

Louisa Ariantjelangi

SRP. 2020; 11(12): 127-133

» Abstract (?mno=134330) » PDF (index.php?fulltxt=134330&fulltxtj=196&fulltxtp=196-1599892888.pdf) » doi: 10.31838/srp.2020.12.21 (http://dx.doi.org/10.31838/srp.2020.12.21)

24. CHANGES IN LABORATORY DATA AFTER LIVER SURGERY, EARLY POSTOPERATIVE PERIOD

Nurbek Ilyassov, Zhaksylyk Doskaliev, Anuar Abdikarimov, Berdybek Yelchibaev, Yerlan Nurgaliyev, Ruslan Kanaev, Ainara Darbayeva

SRP. 2020; 11(12): 134-137

» Abstract (?mno=134544) » PDF (index.php?fulltxt=134544&fulltxtj=196&fulltxtp=196-1599977008.pdf) 10.31838/srp.2020.12.22 (http://dx.doi.org/10.31838/srp.2020.12.22)	» doi:
25. Implementation of Development Policy for Livestock Farming Business in Gorontalo Regency, Gorontalo Indonesia	i
Irawaty Igirisa, Misran Rahman, Juriko Abdussamad, Zuchri Abdusamad, Abd. Kadir Husain, Suparman SRP. 2020; 11(12): 138-142	
» Abstract & References (?mno=28462)	
1604934446.pdf) » doi: 10.31838/srp.2020.12.23 (http://dx.doi.org/10.31838/srp.2020.12.23)	
26. Mobile Health and Users Demographic Characteristics and Preferences. A case Study from the UAE	
Halimah Al-blooshi, Samer Ali Al-shami, Safiah Sidek	
SRP. 2020; 11(12): 143-149	
» Abstract (?mno=134545)	» doi:
27. Mobile Health and Users Demographic Characteristics and Preferences. A case Study from the UAE Halimah Al-blooshi, Samer Ali Al-shami, Safiah Sidek	
SRP. 2020: 11(12): 143-149	
» Abstract (?mno=134333) » PDF (index.php?fulltxt=134333&fulltxti=196&fulltxtp=196-1599893281.pdf)	» doi:
10.31838/srp.2020.12.24 (http://dx.doi.org/10.31838/srp.2020.12.24)	
28. A Review Paper on mhealth Continuous Intention Among Diabetes Users in The Developing Countries	
Halimah Al-blooshi, Samer Ali Al-shami, Hayder Adil, Safiah Sidek	
SRP. 2020; 11(12): 150-159	
» Abstract (?mno=134334) » PDF (index.php?fulltxt=134334&fulltxtj=196&fulltxtp=196-1599893386.pdf)	» doi:
10.31838/srp.2020.12.25 (http://dx.doi.org/10.31838/srp.2020.12.25)	
29. Integrated Model of Factors Affecting Drug Addiction Among Juveniles in UAE	
SRP. 2020; 11(12): 160-166	
» Abstract (?mno=134335)	» doi:
10.31838/srp.2020.12.26 (http://dx.doi.org/10.31838/srp.2020.12.26)	
30. Catfish (Silurus Glanis) Remaining Parts as A Source of Biodiesel Preparation	
Kadhim F. Kadhim, Inaam M. N. Alrubayae, Ali M. Amer	
SRP. 2020; 11(12): 167-171	
» Abstract (?mno=134336) » PDF (index.php?fulltxt=134336&fulltxtj=196&fulltxtp=196-1599893595.pdf)	» doi:
10.31838/srp.2020.12.27 (http://dx.doi.org/10.31838/srp.2020.12.27)	
31. A Review Paper on MOOCs Development Stages, Types, and Opportunities and Challenges	
Salem Aldahmani, Samer Ali Al-shami, Hayder Adil, Safiah Sidek	
SRP. 2020; 11(12): 172-179	
» Abstract (?mno=134337) » PDF (index.php?fulltxt=134337&fulltxtj=196&fulltxtp=196-1599893674.pdf)	» doi:
10.31838/srp.2020.12.28 (http://dx.doi.org/10.31838/srp.2020.12.28)	

/2020	Systematic Reviews in Fharmacy	
32.	Determination of Genetic Relationships and Pathogenicity of Oral Candidiasis Etiological Agents in Pedi	atric
	Malignant Patients in Basrah Province, Iraq	
	Inaam M. N. Alrubayae, Ayat Al-laaeiby, Mohammed Hussein Minati, Shrouk A. H. ALibraheem	
	SRP. 2020; 11(12): 180-188	
	» Abstract (?mno=134338) » PDF (index.php?fulltxt=134338&fulltxtj=196&fulltxtp=196-1599893760.pdf)	» doi:
	10.31838/srp.2020.12.29 (http://dx.doi.org/10.31838/srp.2020.12.29)	
33.	TO BE CREATIVE, OR NOT TO BE CREATIVE; THAT IS THE QUESTION!	
	Suhair Al-Alami	
	SRP. 2020; 11(12): 189-191	
	» Abstract (?mno=134761) » PDF (index.php?fulltxt=134761&fulltxtj=196&fulltxtp=196-1600060460.pdf)	» doi:
	10.31838/srp.2020.12.30 (http://dx.doi.org/10.31838/srp.2020.12.30)	
34.	THE DETERMINANTS OF FINANCIAL INCLUSION IN LATIN AMERICA AND EUROPE (BRAZIL AND ROMA	NIA
	CASE)	
	Yaseen Altarawneh, Mohamed Al-Nuaimi, Ade Al-Nimri	
	SRP. 2020; 11(12): 192-196	
	» Abstract (?mno=134762) » PDF (index.php?fulltxt=134762&fulltxtj=196&fulltxtp=196-1600060652.pdf)	» doi:
	10.31838/srp.2020.12.31 (http://dx.doi.org/10.31838/srp.2020.12.31)	
35.	IMPACT OF PREFERENTIAL TRADE AGREEMENT (PTA) ON THE EXPORT OF ASEAN+4	
	Rossanto Dwi Handoyo, Wisnu Wibowo, Angga Erlando, and Retno Putri Nurkumalasari	
	SRP. 2020; 11(12): 197-204	
	» Abstract (?mno=134764) » PDF (index.php?fulltxt=134764&fulltxtj=196&fulltxtp=196-1600060851.pdf)	» doi:
	10.31838/srp.2020.12.32 (http://dx.doi.org/10.31838/srp.2020.12.32)	
36.	THE ADOPTION OF SOCIAL MEDIA BY PR PRACTITIONERS IN THE UNITED ARAB EMIRATES	
	Tahseen Mansour, Ibrahim Ahmad Abu-Arqoub	
	SRP. 2020; 11(12): 205-209	
	» Abstract (?mno=134765) » PDF (index.php?fulltxt=134765&fulltxtj=196&fulltxtp=196-1600060987.pdf)	» doi:
	10.31838/srp.2020.12.33 (http://dx.doi.org/10.31838/srp.2020.12.33)	
37.	HALAL CERTIFICATION OF PATENTED MEDICINES IN INDONESIA IN DIGITAL AGE: A PANACEA FOR TH	IE PAIN?
	Mas Rahmah Nurul Barizah	
	SRP. 2020; 11(12): 210-217	
	» Abstract (?mno=134766) » PDF (index.php?fulltxt=134766&fulltxtj=196&fulltxtp=196-1600061076.pdf)	» doi:
	10.31838/srp.2020.12.34 (http://dx.doi.org/10.31838/srp.2020.12.34)	
38.	THE EFFECTIVENESS OF SOCIAL MEDIA USE FOR LOCAL GOVERNANCE DEVELOPMENT	
	Sataporn Roengtam	
	SRP. 2020; 11(12): 218-225	
	» Abstract (?mno=134768) » PDF (index.php?fulltxt=134768&fulltxtj=196&fulltxtp=196-1600061258.pdf)	» doi:
	10.31838/srp.2020.12.35 (http://dx.doi.org/10.31838/srp.2020.12.35)	
30	REGULATORY FOCUS AND EMPLOYEE CREATIVITY: THE ROLE OF INDIVIDUAL PARTICIPATION AND	
00.	INTELLECTUAL STIMULATION	

Jovi Sulistiawana, Dian Ekowati, Poppy Sagita Putri

SRP. 2020; 11(12): 226-234

» Abstract (?mno=134769)	» PDF (index.php?fulltxt=134769&fulltxtj=196&fulltxtp=196-1600061364.pdf)	» doi:
10.31838/srp.2020.12.36 (http://	dx.doi.org/10.31838/srp.2020.12.36)	

40. THE IMPACT OF ZAKAT, EDUCATION EXPENDITURE, AND HEALTH EXPENDITURE TOWARDS POVERTY REDUCTION

Sri Herianingrum, Rihfenti Ernayani, Haryo Seto, M.N.H. Rayandono, M.Q.Fauzy SRP. 2020; 11(12): 235-239

» Abstract (?mno=134770) » PDF (index.php?fulltxt=134770&fulltxtj=196&fulltxtp=196-1600061572.pdf) » doi: 10.31838/srp.2020.12.37 (http://dx.doi.org/10.31838/srp.2020.12.37)

41. THE ROLE OF KNOWLEDGE MANAGEMENT IN IMPROVING THE PERFORMANCE OF MEDIA INSTITUTIONS: A CASE STUDY OF ABU DHABI TV

Tahseen Mansour, Ibrahim Ahmad Abuarqoub

SRP. 2020; 11(12): 240-244

» Abstract (?mno=134771) » PDF (index.php?fulltxt=134771&fulltxtj=196&fulltxtp=196-1600061674.pdf) » doi: 10.31838/srp.2020.12.38 (http://dx.doi.org/10.31838/srp.2020.12.38)

42. DENTAL STUDENTS' PERCEPTION OF ONLINE LECTURE USING VIDEO CONFERENCING

Aqsa Sjuhada Oki, Yuliati, Jenny Sunariani, Anis Irmawati, Muhammad Luthfi

SRP. 2020; 11(12): 245-248

» Abstract (?mno=134837) » PDF (index.php?fulltxt=134837&fulltxtj=196&fulltxtp=196-1600067825.pdf) » doi: 10.31838/srp.2020.12.39 (http://dx.doi.org/10.31838/srp.2020.12.39)

43. VALIDATION OF FOREST FIRE HOTSPOT ANALYSIS IN GIS USING FOREST FIRE CONTRIBUTORY FACTORS

El-Said Mamdouh Mahmoud Zahran, Shahriar Shams, Safwanah Ni'matullah Binti Mohd Said

SRP. 2020; 11(12): 249-255

» Abstract (?mno=134776) » PDF (index.php?fulltxt=134776&fulltxtj=196&fulltxtp=196-1600062015.pdf) » doi: 10.31838/srp.2020.12.40 (http://dx.doi.org/10.31838/srp.2020.12.40)

44. THE CHALLENGES OF ACADEMIC LIBRARY LEADERS IN DISRUPTION ERA

Dyah Puspitasari Srirahayu, Anis Eliyana, Indrianawati Usman

SRP. 2020; 11(12): 256-258

» Abstract (?mno=134778) » PDF (index.php?fulltxt=134778&fulltxtj=196&fulltxtp=196-1600062122.pdf) » doi: 10.31838/srp.2020.12.41 (http://dx.doi.org/10.31838/srp.2020.12.41)

45. "GREEN VILLAGE DESTINATION" DEVELOPMENT STRATEGY BASED ON ECGO BIKE IN PELAGA DISTRICT PETANG BADUNG BALI INDONESIA

I Nyoman Sunarta, Putu Eka Wirawan, A.A Ayu Arun Suwi Arianty, Komang Ratih Tunjungsari, I Made Trisna Semara, Diena Mutiara Lemy, Sri Pujiastuti, Nuruddin

SRP. 2020; 11(12): 259-266

» Abstract (?mno=134784) » PDF (index.php?fulltxt=134784&fulltxtj=196&fulltxtp=196-1600062359.pdf) » doi: 10.31838/srp.2020.12.42 (http://dx.doi.org/10.31838/srp.2020.12.42)

46. PUBLIC RELATIONS: THE CASE OF JORDAN'S ACCESSION TO THE GULF COOPERATION

Faisal Al Serhan

SRP. 2020; 11(12): 267-275

» Abstract (?mno=134787) » PDF (index.php?fulltxt=134787&fulltxtj=196&fulltxtp=196-1600062456.pdf) » doi: 10.31838/srp.2020.12.43 (http://dx.doi.org/10.31838/srp.2020.12.43)

47. MOTIVATION OF INDONESIAN CULINARY SMES TO ENGAGE IN INTERNATIONAL MARKETS

Evy Rachmawati Chalduna, Gatot Yudokob, Eko Agus Prasetioc, Jessica Setiawand, Sri Herlianae, Qori Ainaf, Sri Hartatig

SRP. 2020; 11(12): 276-282

» Abstract (?mno=134789) » PDF (index.php?fulltxt=134789&fulltxtj=196&fulltxtp=196-1600062558.pdf) » doi: 10.31838/srp.2020.12.44 (http://dx.doi.org/10.31838/srp.2020.12.44)

48. CULTURAL HERITAGE TOURISM IN INDONESIA POTENTIAL OF "GUNUNG GANGSIR TEMPLE" AS A TOURIST ATTRACTION

Nuruddin, Akhmad Fajar Ma'rufin, Putu Eka Wirawan, I Wayan Pantiyasa, I Made Trisna Semara, Estikowati, Fitria Earlike Anwar Sani

SRP. 2020; 11(12): 283-289

» Abstract (?mno=134792) » PDF (index.php?fulltxt=134792&fulltxtj=196&fulltxtp=196-1600062668.pdf) » doi: 10.31838/srp.2020.12.45 (http://dx.doi.org/10.31838/srp.2020.12.45)

49. EFFECTS OF A LEADER PASSIONATE PERFORMANCE MODERATION ON THE PERFORMANCE OF FEMALE MARKETING EMPLOYEES IN STATE-OWNED COMMERCIAL BANKS IN MEDAN CITY — INDONESIA

Elperida Juniarni Sinurat, Prihatin Lumbanraja, Yenni Absah and Parapat Gultom

SRP. 2020; 11(12): 290-295

» Abstract (?mno=134797) » PDF (index.php?fulltxt=134797&fulltxtj=196&fulltxtp=196-1600062814.pdf) » doi: 10.31838/srp.2020.12.46 (http://dx.doi.org/10.31838/srp.2020.12.46)

50. EFFECTS OF A LEADER PASSIONATE PERFORMANCE MODERATION ON THE PERFORMANCE OF FEMALE MARKETING EMPLOYEES IN STATE-OWNED COMMERCIAL BANKS IN MEDAN CITY — INDONESIA

Elperida Juniarni Sinurat, Prihatin Lumbanraja, Yenni Absah and Parapat Gultom

SRP. 2020; 11(12): 290-295

» Abstract (?mno=134836) » PDF (index.php?fulltxt=134836&fulltxtj=196&fulltxtp=196-1600067550.pdf) » doi: 10.31838/srp.2020.12.46 (http://dx.doi.org/10.31838/srp.2020.12.46)

51. CONFLICTING IDEAS ON MUSIC AMONG INDONESIAN ISLAMIC FIGURES

Suharno Suharno, Milwan Milwan

SRP. 2020; 11(12): 296-301

» Abstract (?mno=135265) » PDF (index.php?fulltxt=135265&fulltxtj=196&fulltxtp=196-1600182207.pdf) » doi: 10.31838/srp.2020.12.47 (http://dx.doi.org/10.31838/srp.2020.12.47)

52. ANALYSIS DESIGN STUDENT CRITICAL ABILITY WITH PROBLEM-BASE LEARNING AND PROJECT-BASED LEARNING MODELS: CASE STUDY IN JOMBANG INDONESIA

Munawaroh, Qomariyatus Sholihah, Kumoro Asto Lenggono

SRP. 2020; 11(12): 302-308

» Abstract (?mno=135268) » PDF (index.php?fulltxt=135268&fulltxtj=196&fulltxtp=196-1600182386.pdf) » doi: 10.31838/srp.2020.12.48 (http://dx.doi.org/10.31838/srp.2020.12.48)

53. POSTMODERN THOUGHT OF CURRICULUM DEVELOPMENT IN A PHILOSOPHICAL FOUNDATION

Zakarias S. Soeteja

SRP. 2020; 11(12): 309-312

» Abstract (?mno=135269) » PDF (index.php?fulltxt=135269&fulltxtj=196&fulltxtp=196-1600182483.pdf) » doi: 10.31838/srp.2020.12.49 (http://dx.doi.org/10.31838/srp.2020.12.49)

54. ANALYSIS OF PERCEIVED USEFULNESS, PERCEIVED EASE OF USE, AND PERCEIVED RISK TOWARD ONLINE SHOPPING IN THE ERA OF COVID-19 PANDEMIC

Sri Setyo Iriani, Anik Lestari Andjarwati

SRP. 2020; 11(12): 313-320

» Abstract (?mno=135270) » PDF (index.php?fulltxt=135270&fulltxtj=196&fulltxtp=196-1600182590.pdf) » doi: 10.31838/srp.2020.12.50 (http://dx.doi.org/10.31838/srp.2020.12.50)

55. UTILIZATION OF FILM TECHNOLOGY FOR ANTICORRUPTION LEARNING AMONG STUDENTS IN LAW EDUCATION SUBJECT

Nelvitia Purba, Sri Sulistyawaty, Yulia Arfanti

SRP. 2020; 11(12): 321-324

» Abstract (?mno=135272) » doi: 10.31838/srp.2020.12.51 (http://dx.doi.org/10.31838/srp.2020.12.51)

56. UTILIZATION OF FILM TECHNOLOGY FOR ANTICORRUPTION LEARNING AMONG STUDENTS IN LAW EDUCATION SUBJECT

Nelvitia Purba, Sri Sulistyawaty, Yulia Arfanti

SRP. 2020; 11(12): 321-324

» Abstract (?mno=135274) » PDF (index.php?fulltxt=135274&fulltxtj=196&fulltxtp=196-1600182819.pdf) » doi: 10.31838/srp.2020.12.51 (http://dx.doi.org/10.31838/srp.2020.12.51)

57. IMPACT OF CYBERBULLYING ON THE DEVELOPMENT OF STUDENT SOFT SKILLS IN ENGINEERING EDUCATION

Yasdin Yasdin, Amirullah Abduh, Muhammad Iskandar Musa, Bakhrani Rauf

SRP. 2020; 11(12): 325-329

» Abstract (?mno=135276) » PDF (index.php?fulltxt=135276&fulltxtj=196&fulltxtp=196-1600182918.pdf) » doi: 10.31838/srp.2020.12.52 (http://dx.doi.org/10.31838/srp.2020.12.52)

58. PERCEIVED LEARNING ASSISTANCE AND PERCEIVED COMMUNITY BUILDING ASSISTANCE: STUDY ON E-LEARNING SYSTEM

Dodik Ariyanto, Ayu Aryista Dewi, I Made Gilang Jhuniantara

SRP. 2020; 11(12): 330-339

» Abstract (?mno=135278) » PDF (index.php?fulltxt=135278&fulltxtj=196&fulltxtp=196-1600183002.pdf) » doi: 10.31838/srp.2020.12.53 (http://dx.doi.org/10.31838/srp.2020.12.53)

59. CHARACTER EDUCATION FOR EDUCATORS THROUGH SCHOOL BASED ON ENVIRONMENTAL

CONSERVATION IN MALANG DISTRICT

Mohamad Syahri, Ahmad Sofwani, Defrian Marza Arisandi

SRP. 2020; 11(12): 340-347

» Abstract (?mno=135279) » PDF (index.php?fulltxt=135279&fulltxtj=196&fulltxtp=196-1600183115.pdf) » doi: 10.31838/srp.2020.12.54 (http://dx.doi.org/10.31838/srp.2020.12.54)

60. MATHEMATICS LANGUAGE UNDERSTANDING OF TEACHERS' CANDIDATE IN MATHEMATICS LEARNING

Baiduri, Yus Mochamad Cholily, Dina Amalya Lapele

SRP. 2020; 11(12): 348-355

» Abstract (?mno=135280) » PDF (index.php?fulltxt=135280&fulltxtj=196&fulltxtp=196-1600183199.pdf) » doi: 10.31838/srp.2020.12.55 (http://dx.doi.org/10.31838/srp.2020.12.55)

61. Fibroblast Viability Test Toward Red Dragon Fruit (Hylocereus polyrhizus) Peel Ethanolic Extract

Annabella Natasha Kylanel, Vinna Kurniawati Sugiaman, Natallia Pranata

SRP. 2020; 11(12): 356-360

» Abstract (?mno=26203) » PDF (index.php?fulltxt=26203&fulltxtj=196&fulltxtp=196-1604498390.pdf) » doi: 10.31838/srp.2020.12.56 (http://dx.doi.org/10.31838/srp.2020.12.56)

62. The Effectiveness Analysis Of Medical Care Management In The Regional Vascular Center For Patients Of Elderly And Senile Age With Acute Myocardial Infarction

A.M. Gritsanchuk, O. V. Gridnev, V.V. Fomin, E. V. Pesennikova, S.S. Kuchits **SRP. 2020; 11(12): 361-365**

» Abstract (?mno=27406) » PDF (index.php?fulltxt=27406&fulltxtj=196&fulltxtp=196-1604668589.pdf) » doi: 10.31838/srp.2020.12.57 (http://dx.doi.org/10.31838/srp.2020.12.57)

63. Features Of The Drug Labeling System Implementation In The Russian Federation: Positive And Negative Experience

Ekaterina O. Kuchits, Oleg V. Gridnev, Gelya N. Gildeeva, Elena V. Pesennikova, Sergey S. Kuchits SRP. 2020; 11(12): 366-369

» Abstract & References (?mno=27408)
 » PDF (index.php?fulltxt=27408&fulltxtj=196&fulltxtp=196-1604668662.pdf)
 » doi: 10.31838/srp.2020.12.58 (http://dx.doi.org/10.31838/srp.2020.12.58)

64. Morphological, Immunophysiological And Clinical Features Of Respiration And Blood Circulation Under The Influence Of Combined Stress In The Covid-19 Conditions

K. M. Khamchiyev, F. M. Suleimenova, G.K. Sagimova, K.T. Adaibayev, A. F. Suchshenko, Ye.M. Serkesh SRP. 2020; 11(12): 370-374

» Abstract (?mno=28327) » PDF (index.php?fulltxt=28327&fulltxtj=196&fulltxtp=196-1604916558.pdf) » doi: 10.31838/srp.2020.12.59 (http://dx.doi.org/10.31838/srp.2020.12.59)

65. Assessment Of Drug Nephrocisin's Hypoazotemic Efficiency At Various Stages Of Chronic Kidney Disease In Clinical Conditions

Burkhonzhon A. Munavvarov, Maksud Atabaevich Sabirov, Anastasia P. Markushina, Khabib B. Barnoev, Sharof N. Eshonov

SRP. 2020; 11(12): 375-380

» Abstract & References (?mno=27415)
 » PDF (index.php?fulltxt=27415&fulltxtj=196&fulltxtp=196-1604668898.pdf)
 » doi: 10.31838/srp.2020.12.60 (http://dx.doi.org/10.31838/srp.2020.12.60)

66. Efficacy And Safety Of Trombless® Neo And Detralex® In The Treatment Of Chronic Venous Diseases Are The Results Of An Open-Label, Comparative, Multicenter Randomized Prospective Study

G.N. Gildeeva, A.V. Belostotsky, O.V. Gridnev, V.I. Yurkov

SRP. 2020; 11(12): 381-384

» Abstract & References (?mno=27416)
 » PDF (index.php?fulltxt=27416&fulltxtj=196&fulltxtp=196-1604669023.pdf)
 » doi: 10.31838/srp.2020.12.61 (http://dx.doi.org/10.31838/srp.2020.12.61)

67. Cas Aiss SRP	e Report. Late-Diagnosec ulu Kapassova, Gulmira De 2 2020: 11(12): 385-387	d Celiac Disease As A Trigger Of Family History Of Gluten Related Disorde erbissalina, Dinara Zhunussova,Elmira Kelimberdiyeva, Baurzhan Iskakov	rs
» Ab	estract (?mno=28324)	» PDF (index.php?fulltxt=28324&fulltxtj=196&fulltxtp=196-1604916074.pdf) //dx doi.org/10.31838/srp.2020.12.62)	» doi:
10.5	1000/31p.2020.12.02 (http:	//dx.doi.org/10.01000/sip.2020.12.02)	
68. The	Role Of The Tutor And S	tudents In The Process Of Problem-Based Learning (Pbl)	
К.М. SRP	Khamchiyev, Y.Kh. Batyay 2 2020; 11(12): 388-393	yeva, A.Kh. Shandaulov, A.M. Zhashkeyeva, F.M. Suleimenova, G.K.Sagimova	
» Ab 10.3	stract (?mno=28326) 1838/srp.2020.12.63 (http:	» PDF (index.php?fulltxt=28326&fulltxtj=196&fulltxtp=196-1604916180.pdf) //dx.doi.org/10.31838/srp.2020.12.63)	» doi:
69. Gree	en Supply Chain, Green C	Communication and Firm Performance: Empirical Evidence from Thailand	
Lime SRP	ega Candrasa, Cia Cai Cen 2 2020: 11(12): 398-406	n, Willy Cahyadi, Lukieto Cahyadi, Ikbar Pratama	
» Ab	stract (?mno=28320)	» PDF (index.php?fulltxt=28320&fulltxtj=196&fulltxtp=196-1604915747.pdf)	» doi:
10.3	1838/srp.2020.12.65 (http:	//dx.doi.org/10.31838/srp.2020.12.65)	
70. EFF URE Evi k SRP » Ab 10.3	ECT OF GIVING ETHANO CUM AND CREATININE LE Kurniawaty, Frisca Febe Lu 2 2020; 11(12): 407-408 Instract (?mno=28674) 1838/srp.2020.12.66 (http:	DL JENGKOL SEEDS (Pithecellobium lobatum Benth.) ON THE IMPROVEM EVELS OF WHITE RAT Imban Gaol, Intanri Kurniati, Silvia Andriani, Soraya Rahmanisa, Silvia Andriani » PDF (index.php?fulltxt=28674&fulltxtj=196&fulltxtp=196-1604997871.pdf) //dx.doi.org/10.31838/srp.2020.12.66)	E NT OF » doi:
71. APC MYC	PTOSIS INDUCING POTE	ENTIAL AND IMMUNOMODULATION OF COLOCASIA ESCULENTA AGAINS	ЗТ
Mus [.] SRP	tafa M.Farhan, Essa D Alh 2 2020: 11(12): 409-417	theal, Saeed Arrak Turkey, kamal jawad awni	
» Ab 10.3	stract (?mno=4153) 1838/srp.2020.12.67 (http:	» PDF (index.php?fulltxt=4153&fulltxtj=196&fulltxtp=196-1605685488.pdf) //dx.doi.org/10.31838/srp.2020.12.67)	» doi:
72. Brea Zena	ast Fibroadenoma Featur a.M.Al hindawi	es Assessment by Ultrasonography	
» Ab	stract (?mno=28679)	» PDF (index.php?fulltxt=28679&fulltxti=196&fulltxtp=196-1604998017.pdf)	» doi:
10.3	1838/srp.2020.12.68 (http:	//dx.doi.org/10.31838/srp.2020.12.68)	
73. DIFF MES NOF	FERENCE BETWEEN EPI SENCHYMAL STEM CELL RVEGICUS)	THELIAL AND COLLAGEN IN SECOND DEGREE BURNS BETWEEN HUMA . EXTRACT AND SILVER SULFADIAZINE IN THE MALE WHITE RAT (RATTU	IN CORD
Evi ł SRP	<pre>{urniawaty, Efrida Wargane 2 2020: 11(12): 423-426</pre>	egara, Ni Made Ariyuliami Savitri	
» Ab	estract (?mno=28680)	» PDF (index.php?fulltxt=28680&fulltxtj=196&fulltxtp=196-1604998144.pdf)	» doi:

10.31838/srp.2020.12.69 (http://dx.doi.org/10.31838/srp.2020.12.69)

74. The effectiveness analysis of medical care management in the Regional vascular center for patients of elderly and senile age with acute myocardial infarction A.M. Gritsanchuk, O. V.Gridnev, V.V. Fomin, E. V. & 1055;& 1077;& 1089;& 1077;& 1085;& 1085;& 1080;& 1082;& 1086;& 1074;& 1072;, S.S. Kuchits SRP. 2020; 11(12): 427-431 » Abstract (?mno=30277) » PDF (index.php?fulltxt=30277&fulltxtj=196&fulltxtp=196-1605277515.pdf) » doi: 10.31838/srp.2020.12.70 (http://dx.doi.org/10.31838/srp.2020.12.70) 75. Features of the drug labeling system implementation in the Russian Federation: positive and negative experience FGAOU VO First MGMU them. THEM. Sechenova of the Ministry of Health of Russia (Sechenov University) Ekaterina O. Kuchits, Oleg V. Gridnev, Gelya N. Gildeeva, Elena V. Pesennikova, Sergey S. Kuchits SRP. 2020; 11(12): 432-435 » Abstract (?mno=30300) » PDF (index.php?fulltxt=30300&fulltxtj=196&fulltxtp=196-1605277648.pdf) » doi: 10.31838/srp.2020.12.71 (http://dx.doi.org/10.31838/srp.2020.12.71) 76. INFLUENCE OF AN EFFECTIVE CONTRACT ON THE QUALITY OF RENDERING MEDICAL SERVICES IN THE **RUSSIAN FEDERATION** S.S. Kuchits, O.V. Gridnev, E.V. Pesennikova, E.O. Kuchits, & 1040;. & 1052;. Gritsanchuk SRP. 2020; 11(12): 436-439 » Abstract (?mno=30302) » PDF (index.php?fulltxt=30302&fulltxtj=196&fulltxtp=196-1605277818.pdf) » doi: 10.31838/srp.2020.12.72 (http://dx.doi.org/10.31838/srp.2020.12.72) 77. Educational Guidelines for Nurses' Competence Level Caring for Patients with Accidental Chest Trauma during **Emergency Period** Tahany El-Senousy, Fatma Mostafa Mahrous, Shimaa Nabil Abd- Al Salam, Hanem Ismael Zedan SRP. 2020; 11(12): 440-449 » Abstract (?mno=738) » PDF (index.php?fulltxt=738&fulltxtj=196&fulltxtp=196-1605351104.pdf) » doi: 10.31838/srp.2020.12.73 (http://dx.doi.org/10.31838/srp.2020.12.73) 78. Role of Prostatic Artery Embolization in Management of Benign Prostatic Hyperplasia Islam M. Allam, Osama M. Hetta, Ahmed H. Soliman, and Ayman M.Ibrahim SRP. 2020; 11(12): 450-455 » Abstract (?mno=736) » PDF (index.php?fulltxt=736&fulltxtj=196&fulltxtp=196-1605350870.pdf) » doi: 10.31838/srp.2020.12.74 (http://dx.doi.org/10.31838/srp.2020.12.74) 79. Characteristics Growth and Adaptability of Dwarf-odot Elephantgrass (Pennisetum purpureum cv. Mott) Grown in Gorontalo at Established Year Muhammad Mukhtar SRP. 2020; 11(12): 456-462 » Abstract (?mno=5546) » PDF (index.php?fulltxt=5546&fulltxtj=196&fulltxtp=196-1605766416.pdf) » doi: 10.31838/srp.2020.12.75 (http://dx.doi.org/10.31838/srp.2020.12.75) 80. Comparative Study of Dexamethasone Versus Ondansetron as Adjuvants to the Intra-Peritoneal Irrigation of Bupivacaine for Reducing the Postoperative Pain in Patients Undergoing Elective Laparoscopic Cholecystectomy

Basim Herez Ali, Ameer Zuhair Hameed, Raad Jasim Isa

3KP. 2020, 11(12). 403-400		
» Abstract (?mno=5557)	» PDF (index.php?fulltxt=5557&fulltxtj=196&fulltxtp=196-1605767501.pdf)	» doi:
10.31838/srp.2020.12.76 (h	ttp://dx.doi.org/10.31838/srp.2020.12.76)	
31. Relationship of Cybersecu	urity and the National Security of the Country: Iraq Case Study	
Asmaa Khalid Jarjees Al-Tae	e, Hameeda Abdul-Hussain Al-Dhalimi, Adnan Kadhum Jabbar Al-Shaibani	
SRP. 2020; 11(12): 469-476		
» Abstract (?mno=5632)	» PDF (index.php?fulltxt=5632&fulltxtj=196&fulltxtp=196-1605775110.pdf)	» doi:
10.31838/srp.2020.12.77 (h	ttp://dx.doi.org/10.31838/srp.2020.12.77)	
2. The Formation of Homose	xual Behavior in South Sulawesi Province (Study on Young Homosexual in	Makassa
City and Bulukumba Distri	ict)	
Andi Asrina, Sukri Palutturi		
SRP. 2020; 11(12): 477-483		
» Abstract (?mno=5567)	» PDF (index.php?fulltxt=5567&fulltxtj=196&fulltxtp=196-1605767680.pdf)	» doi:
10.31838/srp.2020.12.78 (h	ttp://dx.doi.org/10.31838/srp.2020.12.78)	
3. X-Ray Fluorescence (XRF)) Analysis in Masonry Heritage Building Conservation in Malaysia	
	Ahmad	
Tan Chin Ling and A Ghafar		
Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488		
Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569)	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf)	» doi:
Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (h 34. Deploying Marketing Indic Customer Satisfaction: A (» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) ators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies	» doi: • of
Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (ht 34. Deploying Marketing Indic Customer Satisfaction: A G Effed Darta, Henni Febriawa SRP. 2020: 11(12): 489-499	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) ators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah	» doi: • of
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (http://dx.org/abs/srp.2020.12.79 (http://dx.org/abs/srp.2020.12.79 (http://dx.org/abs/srp.2020; 12.79 (http://dx.org/abs/sr	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) 	» doi: • of » doi:
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (htte://doi.org/10.11838/srp.2020.12.79 4. Deploying Marketing Indic Customer Satisfaction: A deploying Marketing Indic Customer Satisfaction: A deploying SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (htte://doi.org/10.11838/srp.2020.12.80 (htte://doi.org/10.11838/srp.2020.12.80 (htte://doi.org/10.11838/srp.2020.12.80 (http://doi.org/10.11838/srp.2020.12.80 (http://doi.01838/srp.2020.12.80 (httt://doi.01838/srp.2020.12.80 (httt://doi.01838	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) 	» doi: • of » doi:
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (ht/li> 4. Deploying Marketing Indic Customer Satisfaction: A (Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (ht/li> 5. Study of Land Quality and 	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) eators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize 	» doi: • of » doi: Varieties
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (htte://doi.org/10.11838/srp.2020.12.79 4. Deploying Marketing Indic Customer Satisfaction: A Geffed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (htte://doi.org/10.11838/srp.2020.12.80 5. Study of Land Quality and Gorontalo 	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) eators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize 1 	» doi: • of » doi: Varieties
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (ht) 4. Deploying Marketing Indic Customer Satisfaction: A Geffed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (ht) 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemal 	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize Trino, Sudarto 	» doi: • of » doi: Varieties
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hit) 4. Deploying Marketing Indic Customer Satisfaction: A (Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (ht) 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemal SRP. 2020; 11(12): 500-509 	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize Trino, Sudarto 	» doi: • of » doi: Varieties
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (ht) 4. Deploying Marketing Indic Customer Satisfaction: A Geffed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (ht) 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemal SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) eators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize Troo, Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf)	» doi: • of » doi: Varieties » doi:
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (ht) 4. Deploying Marketing Indic Customer Satisfaction: A (Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (ht) 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soeman SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (ht) 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) eators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize True, Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.81)	» doi: • of » doi: Varieties » doi:
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hi 4. Deploying Marketing Indic Customer Satisfaction: A G Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (hi 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemai SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (hi 6. Glycemic Control and Period 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize mo, Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.81) ipheral Nerve Function after Healed of Plantar Foot Diabetic Ulcer in Seman	» doi: • of » doi: Varieties » doi: • ang,
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hi) 4. Deploying Marketing Indic Customer Satisfaction: A (Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (hi) 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemain SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (hi) 6. Glycemic Control and Pering 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize and Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.81) ipheral Nerve Function after Healed of Plantar Foot Diabetic Ulcer in Semaria	» doi: • of » doi: Varieties » doi: rang,
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hi 4. Deploying Marketing Indic Customer Satisfaction: A G Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (hi 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemai SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (hi 6. Glycemic Control and Perilindonesia Maria Suryani, Widiastuti Sa 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) actors to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize Troo, Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.81) ipheral Nerve Function after Healed of Plantar Foot Diabetic Ulcer in Semara amekto, Heri-Nugroho	» doi: • of • doi: • doi: • doi: • ang,
 Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hi 4. Deploying Marketing Indic Customer Satisfaction: A G Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (hi 5. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemai SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (hi 6. Glycemic Control and Peri Indonesia Maria Suryani, Widiastuti Sa SRP. 2020; 11(12): 510-514 	» PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) ators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize in the second	» doi: • of • doi: • doi: • doi: • ang,
Tan Chin Ling and A Ghafar SRP. 2020; 11(12): 484-488 » Abstract (?mno=5569) 10.31838/srp.2020.12.79 (hi 34. Deploying Marketing Indic Customer Satisfaction: A (Effed Darta, Henni Febriawa SRP. 2020; 11(12): 489-499 » Abstract (?mno=5903) 10.31838/srp.2020.12.80 (hi 35. Study of Land Quality and Gorontalo Nurdin, M. L Rayes, Soemal SRP. 2020; 11(12): 500-509 » Abstract (?mno=8009) 10.31838/srp.2020.12.81 (hi 36. Glycemic Control and Peril Indonesia Maria Suryani, Widiastuti Sa SRP. 2020; 11(12): 510-514 » Abstract (?mno=8012)	 » PDF (index.php?fulltxt=5569&fulltxtj=196&fulltxtp=196-1605767752.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.79) sators to Ensure Greater SMEs International Performance in Mediating Role Glance of Indonesian Pharmacies ati, Amrullah » PDF (index.php?fulltxt=5903&fulltxtj=196&fulltxtp=196-1605798560.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.80) Land Characteristics that Determine the Productivity of Composite Maize rno, Sudarto » PDF (index.php?fulltxt=8009&fulltxtj=196&fulltxtp=196-1605969652.pdf) ttp://dx.doi.org/10.31838/srp.2020.12.81) ipheral Nerve Function after Healed of Plantar Foot Diabetic Ulcer in Seman amekto, Heri-Nugroho » PDF (index.php?fulltxt=8012&fulltxtj=196&fulltxtp=196-1605969722.pdf) 	» doi: • of • doi: • doi: • doi: • ang, • doi:

Ary Bastari, Hapzi Ali, Hamidah

SRP. 2020; 11(12): 827-838

» Abstract (?mno=26207) » PDF (index.php?fulltxt=26207&fulltxtj=196&fulltxtp=196-1604498683.pdf) » doi: 10.31838/srp.2020.12.122 (http://dx.doi.org/10.31838/srp.2020.12.122)

88. Improving Employee Performance Through Perceived Organizational Support, Organizational Commitment and Organizational Citizenship Behavior

Muhammad Ridwan, Sitti Rizki Mulyani, Hapzi Ali

SRP. 2020; 11(12): 839-849

» Abstract (?mno=26211) » PDF (index.php?fulltxt=26211&fulltxtj=196&fulltxtp=196-1604498909.pdf) » doi: 10.31838/srp.2020.12.123 (http://dx.doi.org/10.31838/srp.2020.12.123)

89. THE IMPACT OF ASYMMETRIC INFORMATION IN MEDICAL SERVICES: A STUDY IN PROGRESSIVE LAW

Nanik Trihastuti, Stephanie Apsari Putri, Bagoes Widjanarko

SRP. 2020; 11(12): 850-855

» Abstract (?mno=135761) » PDF (index.php?fulltxt=135761&fulltxtj=196&fulltxtp=196-1600334005.pdf) » doi: 10.31838/srp.2020.5.124 (http://dx.doi.org/10.31838/srp.2020.5.124)

90. FULFILLING THE SEXUAL NEEDS TO INCREASE THE MENTAL HEALTH OF PRISONERS IN PENITENTIARY INSTITUTION

Desy Maryani, Nur Rochaeti, Nyoman Serikat Putera Jaya

SRP. 2020; 11(12): 856-858

» Abstract (?mno=135762) » PDF (index.php?fulltxt=135762&fulltxtj=196&fulltxtp=196-1600334086.pdf) » doi: 10.31838/srp.2020.5.125 (http://dx.doi.org/10.31838/srp.2020.5.125)

91. IN VITRO ANTIDIABETIC ACTIVITY OF SARGASSUM HYSTRIX EXTRACT AND ITS ETHYL ACETATE FRACTIONS

HUSNI, Amir, SULISTYO, Ratih Paramadina, RAHMA, Syifa Aulia, NUGRAHENI, Prihati Sih, BUDHIYANTI, Siti Ari SRP. 2020; 11(12): 859-865

» Abstract (?mno=135763) » PDF (index.php?fulltxt=135763&fulltxtj=196&fulltxtp=196-1600334198.pdf) » doi: 10.31838/srp.2020.5.126 (http://dx.doi.org/10.31838/srp.2020.5.126)

92. THE IMPACT OF NEON GAS IN THE MANUFACTURE OF LOW ENERGY SAVING

i ketut wijaya

SRP. 2020; 11(12): 866-872

» Abstract (?mno=135764) » PDF (index.php?fulltxt=135764&fulltxtj=196&fulltxtp=196-1600334360.pdf) » doi: 10.31838/srp.2020.5.127 (http://dx.doi.org/10.31838/srp.2020.5.127)

93. EFFECT OF ATTACKANGLE OF CONCAVE AND CONVEX WINGLET VORTEX GENERATORS ON HEAT

TRANSFER AND PRESSURE DROP IN EVAPORATOR FIN AND TUBE HEAT EXCHANGER WITH FIELD SYNERGY PRINCIPLE USING NUMERICAL SIMULATION

SYAIFUL, YUNIANTO, Bambang, SALSABILA, Carisya Dara, S.U., MSK Tony, F. SOETANTO, Maria SRP. 2020; 11(12): 873-887

» Abstract (?mno=135765) » PDF (index.php?fulltxt=135765&fulltxtj=196&fulltxtp=196-1600334482.pdf) » doi: 10.31838/srp.2020.5.128 (http://dx.doi.org/10.31838/srp.2020.5.128)

94. ASSESSMENT OF COASTAL ENVIRONMENT SUPPORTS IN BATAM'S PEMPING ISLAND-INDONESIA FOR PIPELINE GAS TRANSMISSION

MUBARAK, MUBARAK, NURHUDA, AHMAD, SULAIMAN, ALBERTUS

Most Cited Articles

POLICIES & JOURNAL LINKS

(http://www.ejmanager.com/my/srp)

Copyright © 2019 Systematic Reviews in Pharmacy All Rights Reserved. Subject to Login

change without notice from or liability to Systematic Reviews in Pharmacy.

(https://www.ejmanager.com/reviewers/index.php?) (https://www.ejmanager.com/reviewers/index.php?) (https://www.sysrevpharm.org//?) About Publisher (http://www.sysrevpharm.org//?)

.....

sec=aboutpublisher)

Reviewer Login

- Advertising Policy
- (http://www.sysrevpharm.org//?
- sec=policyadvertising)
- Author's Rights and Obligations
- (http://www.sysrevpharm.org//?
- sec=policyauthorsrights)

Conflict of Interest Policy

(http://www.sysrevpharm.org//?sec=policycois)

Digital Archiving & Preservation Policies

(http://www.sysrevpharm.org//?

sec=digitalarchiving)

Editorial Policies (http://www.sysrevpharm.org//?

sec=policyeditorial)

Peer Review Policy

(http://www.sysrevpharm.org//?

sec=peerreviewpolicy)

Editorial & Peer Review Process

(http://www.sysrevpharm.org//?

sec=editorialprocess)

License Information

	5	5
Copyright	Contact Information	(http://www.sysrevpharm.org//?sec=licenseinfo)
Svstematic	Contact Information	(http://www.svsrevpharm.org//?
Reviews in		sec=policyplagiarism)
Pharmacy		Privacy Policy (http://www.svsrevpharm.org//?
All Rights		sec=policyprivacy)
Reserved	<submissions.srp@gmai< td=""><td>ComProtection of Research Participants (Statement</td></submissions.srp@gmai<>	Com Protection of Research Participants (Statement
Subject to		On Human And Animal Rights)
change		(http://www.svsrevpharm.org//?sec=policyhar)
without		Publication Ethics and Publication Malpractice
notice		Statement (http://www.sysrevpharm.org//?
from or		sec=policypubethics)
liability to		Corrections, Retractions & Expressions of
Systematic		Concern (http://www.sysrevpharm.org//?
Reviews in		sec=correctionretractionconcern)
Pharmacy.		Self-Archiving Policies
For best		(http://www.sysrevpharm.org//?
results,		sec=selfarchivingpolicy)
please use		Statement of Informed Consent
Internet		(http://www.sysrevpharm.org//?
Explorer or		sec=informedconsent)
Google		Terms of Use (http://www.sysrevpharm.org//?
Chrome		sec=policytermofuse)

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Nurdin^{1*}, M. L Rayes², Soemarno², Sudarto²

¹Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 ³Soil Science Departmen, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 *E-mail: <u>nurdin@ung.ac.id</u>

ABSTRACT

The challenge of composite maize developing in the future is the low productivity because the maize is grown on land that is not suitable for land quality. This study aims to determine the land quality and land characteristics that control the composite maize productivity in Gorontalo Province. A total of 33 land units were surveyed and their land observed to obtain data on morphology and soil characteristics, climate and terrain characteristics, as well as composite maize productivity data through ubinan plots and direct interviews with maize farmers. Partial least square of structural equation models (PLS-SEM) analysis has been used to determine the land quality and land characteristics that control the composite maize productivity through variable validity and reliability tests, as well as structural model tests. The results showed that the manifest variables were air temperature, rainfall, wet months, dry months, LGP, drainage, coarse materials, effective depth, pH H₂O, pH KCl, C-organic, total N, available P, available K, ESP, slopes, soil erosion, inundation height, inundation time, surface rock, and rock outcrops were valid and able to explain well the latent variables. Furthermore, the latent variables were temperature, water availability, oxygen availability, nutrient retention, nutrients availability, sodicity, erosion hazard, flood hazard, and land preparation used has good composite reliability and high reliability because of the composite reliability and alpha cronbach >0.6. Land quality that controls the composite maize productivity based on the order of importance were nutrient retention, rooting media, land preparation, and nutrients availability. Meanwhile, land characteristics that control the composite maize productivity based on the order of importance were pH KCI, coarse material, rock outcrops, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH KCI, and available K has a positive relationship and has a significant to very significant effect on the composite maize productivity, while the content of course materials, surface rock, and rock outcrops has a negative relationship and has a significant effect on the composite maize productivity.

INTRODUCTION

Low maize productivity is still a major problem in efforts to increase maize production in Indonesia. Until 2018, maize productivity had only reached 5.24 tons/ha [1], while the potential for maize productivity in Indonesia could reach 10-11 tons/ha [2]. Even though the government has rolled out various programs to increase maize production and productivity in order to achieve national maize selfsufficiency.

Gorontalo Province is one of the maizes producing centers in Indonesia with an average maize productivity achievement until 2019 of 5.03 tons/ha [3] or still far below the average national maize productivity. All this time, maize farmers has been more dominant in planting maize with hybrid and composite varieties. There are no references to the productivity of hybrid or composite maize in this area, so it is assumed that the maize productivity achievements are relatively the same at 5.24 tons/ ha. In fact, specifically the productivity of composite maize can reach 5-6 tons/ha [4] [5] or as much as 8.5 tons/ha [6]. Composite maize, besides its production potential is relatively similar to that of hybrid maize, it also has advantages, including being more adaptive in acid soils [5] and can be used as seeds for the next growing season, while hybrid maize can't be planted again. The use of composite maize can reduce the dependence of maize farmers on hybrid maize seed assistance from the government, so that if the maize seed subsidy is stopped, the farmers can plant the composite maize again.

The challenge ahead in developing composite maize is the

Keywords: Quality, characteristic, land, productivity, maize, composite.

Correspondence:

Nurdin

Doctorate Programe, Agriculture Faculty, Brawijaya University. Jalan Veteran Kota Malang, Jawa Timur-Indonesia. 65145 E-mail: <u>nurdin@ung.ac.id</u>

low productivity of composite maize, so it is necessary to address the root of the problem. Maize planted on land with low productivity potential is one of the causes for the low productivity of maize [7]. Meanwhile, land characteristics and quality have a close relationship with maize productivity [8] and each land quality has a significant effect on land suitability for certain uses [9], especially for maize. Research on land quality that controls the productivity of composite maize has been conducted in the Bogor area using stepwise regression analysis [10]. The use of structural equation modeling (SEM) analysis in determining the characteristics and quality of land that control plant productivity has not been widely published, except for [11] who used SEM analysis on older cocoa plants in Kolaka Timur Regency, Southeast Sulawesi Province. Meanwhile, the use of SEM analysis specifically to determine the relationship between land quality and maize productivity has not been found.

The response of maize plants to the diversity of characteristics and quality of land will vary, so it is important to know the quality and characteristics of the land that control the productivity of maize. The diversity of characteristics and complex quality of land in the field really requires a comprehensive analysis technique that is able to simplify the complexity in one analysis system. One analysis option is to use SEM analysis. SEM analysis is able to analyze how much influence each indicator (manifest) of soil physical and chemical properties (latent) has on production in one analysis unit [11]. The use of SEM is very helpful to

Composite Maize Varieties in Gorontalo

determine the effect on indicators and to produce a model that is better than other multivariate analyzes [12] [13]. Partial Least Square (PLS) is a variant of SEM which has a higher level of flexibility because PLS is based on variants, so that the number of samples used does not need to be large, ranges from 30-100, and does not require normal multivariate assumptions compared to CB-SEM. requires a large data sample size (> 100) and the data must be multivariate normal distribution [14] [15]. Therefore, a research on land quality that controls the productivity of composite maize was carried out using SEM-PLS analysis based on the consideration of complex land characteristics and quality, as well as limited data in the land unit in the study area. The purpose of this study was to determine the quality and characteristics of land that control the productivity of composite maize in Gorontalo.

MATERIALS AND METHOD

This research is located in the Sustainable Agriculture Area of Gorontalo Province (Figure 1) and the Soil Laboratory of the Department of Soil, Faculty of Agriculture, Brawijaya University. The timing of this research was started in December 2019 - May 2020. The tools used included the computer, SmatPLS version 2.0, SPSS, Microsoft Excel, and Microsoft Word. While the materials studied included the morphological data and soil characteristics, climate and terrain characteristics data that had been grouped into their respective land qualities, as well as composite maize productivity data from the study area.

Soil surveys and land observations were carried out to obtain morphological data and soil characteristics, climate and terrain characteristics data from the research area. Meanwhile, composite maize productivity data was obtained from the results of ubinan directly on the land of maize farmers and from direct interviews with maize farmers on 33 land units. Furthermore, the diversity of sizes and data units (ratio data) of land characteristics were converted in the form of interval data which were represented as follows were 1 (very low), 2 (low), 3 (medium), 4 (high), and 5 (very high)). After the data is ready, the analysis process is continued using SEM-PLS (Figure 2).

Figure 2. Research Operational Framework

The latent variable in this study was the quality of the land consisting of: temperature (X1), water availability (X2), oxygen availability (X3), root media (X4), nutrient retention (X5), available nutrients (X6), sodicity (X7), erosion hazard (X8), flood hazard (X9), and land preparation (X10). While the manifest variable was the characteristic of the land which consists of temperature (X1.1), rainfall (X2.1), wet months (X2.2), dry months (X2.3), LGP (X2.4), drainage. (X3.1), texture (X4.1), coarse material (X4.2), effective soil depth (X4.3), pH H2O (X5.1), pH KCI (X5.2), C-organic (X5.3), CEC (X5.4), base saturation (X5.5), total N (X6.1), available P2O5 (X6.2), K available (X6.3), ESP (X7.1), slope (X8.1), soil erosion (X8.2), inundation height (X9.1), length of inundation (X9.2), surface rock (X10.1), and rock outcrop (X10.2). The use of SEM-PLS in this study consisted of testing the validity, reliability of the research variables, and testing the structural model. In summary, the test using SEM-PLS was described as follows:

a. Testing the Validity of Research Variables. The basic evaluation carried out in the SEM-PLS analysis is to evaluate the measurement model (outer model) with the aim of knowing the validity and reliability of indicators in measuring research latent variables through convergent validity, discriminant validity, and composite reliability. Convergent validity testing on SEM-PLS is seen from the size of the outer loading of each indicator on its latent variable. A loading factor value above 0.70 is highly recommended, but a loading factor value of 0.50-0.60 can still be tolerated with a tstatistic value of more than 1.96 or a small p-value of 0.05. The loading factor of an indicator with the highest value is the strongest or most important measure in reflecting the latent variable in question. Discriminant validity is an evaluation of the outer model in SEM-PLS using cross loading values to test valid and reliable indicators in explaining or reflecting latent variables. If the correlation of the latent variable with the measurement core of each indicator is greater than the other latent variables, then the latent variable is able to predict the indicator better than other latent variables and is said to be valid.

- b. Research Variable Reliability Testing. Composite reliability and alpha cronbach were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is> 0.60.
- c. Structural Model Testing. Testing of the structural model (inner model) is carried out after the relationship model is built in accordance with the observed data and the suitability of the overall model (goodness of fit model). Testing of structural models and hypotheses is carried out by looking at the estimated value of the path coefficient and the critical point value (t-statistic) which is significant at $\alpha = 0.05$.

Composite Maize Varieties in Gorontalo

Testing the relationship model and hypothesis between variables can be done by testing the direct correlation coefficient between variables. The results of testing the relationship between the X variables and the Y variable in this study are shown by the correlation coefficient and t-statistic, and also seen in the path diagram.

RESULTS AND DISCUSSION

a. Research Variable Validity

The loading factor value of the research variables, where the loading factor value on the indicators was mostly more than

the critical limit of 0.7 with a confidence level of 95% (Table 1). The value of the loading factor which is below the tolerance value of 0.5 at the 95% confidence level where the t-statistic value of each indicator is smaller than the t-table (1,960) is found in the soil texture indicator of the latent variable root media (X₄) which is only 0.173 is also the CEC indicator (X_{5.4}) and the base saturation indicator (X_{5.5}) of the nutrient retention latent variable (X₅), which are only 0.399 and 0.482 respectively. This means that these indicators have not been able to properly form or explain their latent variables.

Table 1. Outer loading research variables

Effect of indicators on latent variables			Loading factors	Status
Air temperature (X _{1.1})	->	Temperature (X ₁)	1.000	Valid
Rainfall (X _{2.1})	->		0.981	Valid
Wet months (X _{2.2})	->	Watan availability (V)	0.989	Valid
Dry months (X _{2.3})	->	water availability (x ₂)	0.827	Valid
LGP (X _{2.4})	->		0.968	Valid
Drainage (X _{3.1})	->	Oxygen availability (X ₃)	1.000	Valid
Texture (X _{4.1})	->		0.173	Not valid
Coarse material (X _{4.2})	->	Rooting media (X4)	-0.921	Valid
Effective depth (X _{4.3})	->		0.912	Valid
pH H ₂ O (X _{5.1})	->		0.768	Valid
рН КСІ (Х _{5.2})	->		0.772	Valid
C-Organic (X _{5.3})	->	Nutrient retention (X5)	0.710	Valid
CEC (X _{5.4})	->		0.399	Not valid
Base saturation (X _{5.5})	->		0.482	Not valid
N Total (X _{6.1})	->		0.799	Valid
Available P (X _{6.2})	->	Nutrient availability (X ₆)	0.521	Valid
Available K (X _{6.3})	->		0.886	Valid
ESP (X _{7.1})	->	Sodicity (X7)	1.000	Valid
Slope (X _{8.1})	->	Exercise beyond (V.)	0.974	Valid
Soil erosion (X _{8.2})	->	EI OSIOII HAZAI U (X8)	0.957	Valid
Inundation height (X _{9.1})	->	Planding harowd (V.)	0.993	Valid
Inundation period (X _{9.2})	->	Flooding hazard (X9)	0.991	Valid
Surface rock (X _{10.1})	->	Land propagation (V)	0.998	Valid
Rock outcrop (X _{10.2})	->	Lanu preparation (A10)	0.998	Valid
Productivity (Y _{1.1})	->	Local maize productivity (Y_1)	1.000	Valid

			1	able 2. Cross loa	ding of research	variables					
Indikator	Temperature (X1)	Water availability (X ₂)	Oxygen availability (X ₃)	Rooting media (X4)	Nutrient retention (X5)	Nutrient availability (X ₆)	Sodicity (X7)	Erosion hazard (X ₈)	Flooding hazard (X9)	Land preparation (X ₁₀)	Composite maize productivity (Y ₁)
Air temperature $(X_{1.1})$	1	0.952309	0.059098	-0.08736	-0.37805	-0.06653	0.38176	0.016269	-0.10297	0.19833	0.042282
Rainfall (X _{2.1})	0.968555	0.980906	0.114576	0.052348	-0.24379	0.058536	0.356547	-0.0379	-0.04621	0.056015	0.156751
Wet months (X _{2.2})	0.926635	0.989185	0.173659	-0.005903	-0.25644	0.062873	0.374745	-0.06373	-0.04367	0.060342	0.177251
Dry months (X _{2.3})	0.759123	0.82697	0.141078	-0.238735	-0.42612	-0.10563	0.47553	-0.11715	0.027746	0.215367	0.076041
LGP (X _{2.4})	0.900431	0.96821	0.13569	-0.003834	-0.28223	0.056251	0.459669	-0.12209	-0.04398	0.059938	0.193991
Drainage (X _{3.1})	0.059098	0.144225	1	0.129338	-0.24128	0.057861	0.084339	-0.50344	0.236555	-0.22277	0.400657
Texture (X _{4.1})	-0.02057	-0.01261	-0.16957	0.172551	0.242032	0.12283	0.217308	0.196875	-0.00074	-0.02261	0.09248
Coarse material (X _{4.2})	-0.00333	-0.1005	-0.13244	-0.921096	-0.38256	-0.6112	0.18822	0.322934	-0.26391	0.846957	-0.35202
Effective depth (X _{4.3})	-0.17758	-0.09256	0.165016	0.912088	0.3519	0.355112	-0.23141	-0.19005	0.095721	-0.76736	0.180089
pH H ₂ O (X _{5.1})	-0.40346	-0.38437	-0.3719	0.29356	0.767791	0.27088	-0.17175	0.151553	-0.02966	-0.08478	0.186569
pH KCl (X5.2)	-0.25953	-0.22811	-0.44804	0.342269	0.771872	0.272936	-0.02729	0.167533	0.098977	-0.18312	0.268161
C-Organic (X _{5.3})	-0.29516	-0.13852	0.096529	0.248076	0.710022	0.612498	0.073184	-0.4692	0.063874	-0.1793	0.384332
CEC (X _{5.4})	0.066756	0.115697	0.003345	0.084182	0.399393	0.421251	0.373179	-0.05735	0.15285	-0.01387	0.281455
Base saturation $(X_{5.5})$	-0.30026	-0.25724	-0.10527	0.412102	0.481624	0.361795	-0.60079	-0.0895	-0.13592	-0.48759	0.136266
N Total (X _{6.1})	0.002878	0.137879	0.07154	0.268606	0.545283	0.798694	0.030267	-0.37884	-0.10212	-0.2485	0.427705
Available P (X _{6.2})	-0.09821	-0.09791	-0.44547	0.211821	0.409315	0.520984	-0.28705	-0.057	0.033581	-0.26033	-0.02547
Available K(X _{6.3})	-0.09732	-0.01031	0.06693	0.614343	0.51245	0.885686	-0.3292	-0.29441	0.237691	-0.6422	0.49531
ESP (X _{7.1})	0.38176	0.405078	0.084339	-0.186069	-0.06947	-0.21259	1	-0.01035	0.201152	0.361936	-0.0249
Slope (X _{8.1})	-0.02207	-0.12714	-0.51717	-0.295103	-0.1643	-0.40295	-0.03466	0.973779	-0.34215	0.324431	-0.64795
Soil erosion (X _{8.2})	0.064136	-0.00224	-0.44709	-0.166166	-0.11161	-0.32907	0.021581	0.956588	-0.12926	0.257787	-0.48649
Inundation height (X _{9.1})	-0.08956	-0.02635	0.225421	0.194354	0.082178	0.127762	0.193925	-0.26735	0.992798	-0.13415	0.175472
Inundation period (X _{9.2})	-0.11594	-0.06329	0.244833	0.199427	0.048584	0.078386	0.205739	-0.2425	0.991369	-0.11616	0.135302
Surface rock (X _{10.1})	0.212772	0.074279	-0.23401	-0.854273	-0.28568	-0.55023	0.376036	0.319248	-0.13208	0.997623	-0.28655
Rock outcrop (X _{10.2})	0.183196	0.051703	-0.21067	-0.868319	-0.29655	-0.55537	0.34638	0.290608	-0.12053	0.997697	-0.28228
Productivity (Y _{1.1})	0.042282	0.177277	0.400657	0.304774	0.418519	0.534535	-0.0249	-0.59733	0.157534	-0.28507	1

Table 2 Cuses loading of u oarch wariable

The standard of loading factor is greater equal to 0.50 [16] [17] [13]. However, in general, based on the indicated values, it can be concluded that the latent variables of land quality have been able to be well established or explained by each indicator and can be said to be convergent valid on these indicators. The cross-loading value for the indicators of latent variables on average is above the cross-loading value of the indicators for other latent variables (Table 2). That is, the greatest cross loading value on the indicator is found in the latent variable too, except for the texture indicator (X_{4.1}) of the root media variable (X₄), the CEC indicator (X_{5.4}) and base saturation $(X_{5.5})$ of the nutrient retention variable (X_5) whose cross loading value is still smaller (<0.5) than the cross loading value of other latent variables. The standard of loading factor is ≥0.50 [16] [17] [13]. Thus, the indicators of each latent variable are mostly able to explain the latent variable itself better than the other variables, so that the research variables are said to be discriminant valid.

Composite reliability and Cronbach alpha were used to test the reliability value between the indicators of the latent variables that formed them. The composite reliability value and Cronbach's alpha are said to be good, if the value is above 0.60 [18]. The composite reliability value on each research variable is more than the limit value (>0.6), except for the root media variable (Table 3). The composite reliability value and the Cronbach alpha value is greater than 0.6 so that the latent variable has good composite reliability and high reliability. A construct is said to be reliable if the Cronbach Alpha value must be >0.6 [19]. Thus, all indicators used in this study have met the criteria or are feasible to be used in the measurement of all latent variables because they have good validity and high reliability. The results of the evaluation of convergent validity and discriminant validity of indicators or variables as well as composite reliability and alpha Cronbach for indicators or variables can be concluded that the indicators as measures of latent variables are valid and reliable measures respectively.

b. Reliability of Research Variables

Table 3. Composite reliability and Cronbach's Alpha values of research variables

Laten variables	Composite reliability	Alpha Cronbach
Temperature (X ₁)	1.000000	1.000000
Water availability (X ₂)	0.970030	0.965126
Oxygen availability (X ₃)	1.000000	1.000000
Rooting media (X ₄)	0.020314	-1.055192
Nutrient retention (X ₅)	0.770518	0.628062
Nutrient availability (X ₆)	0.788289	0.681393
Sodicity (X ₇)	1.000000	1.000000
Erosion hazard (X ₈)	0.964615	0.927731
Flooding hazard (X ₉)	0.992053	0.984010
Land preparation (X ₁₀)	0.997657	0.995304

b. Structural Model Testing

The structural model (inner model) is evaluated by looking at the coefficient value of the relationship path parameter between latent variables. It seems that the soil quality of the root media, nutrient retention, and available nutrients showed a positive correlation and had a significant effect on the productivity of composite maize (Table 4). The quality of land preparation shows a negative correlation and has a significant effect on the productivity of composite maize. This indicates that the better rooting media, available nutrient and nutrient retention and a decrease in the level of land preparation as the productivity of composite maize increases. The results of this study are slightly different from the research report [8] regarding the quality of soil rooting media which has not affected the productivity of maize in the Bogor area, but the quality of soil nutrient retention and available nutrients has a significant effect on the productivity of maize relatively the same as the results of this study.

Table 4. Path coefficient and significance testing

	Endogenous variables		
Exogenous variables	Composite maize produktivity (Y)		
	Path coeffisient	t-statistics (t _{critics} = 2.00)	
Temperature (X ₁)	0.086	1.531	
Water availability (X ₂)	0.457	-0.491	
Oxygen availability (X ₃)	0.099	0.371	
Rooting media (X ₄)	0.091*	2.250	
Nutrient retention (X ₅)	0.740*	2.291	
Nutrient availability (X ₆)	0.283**	6.509	
Sodicity (X7)	-0.194	-0.036	
Erosion hazard (X ₈)	-0.043	-1.043	
Flooding hazard (X ₉)	0.050	-0.050	
Land preparation (X ₁₀)	-0.386*	-2.339	

Significant on level test of 5%; ** Significant on level test of 1%

c. Land quality and characteristics that controlling of composite maize productivity

Based on the previous structural model testing, the most influential land quality and control of composite maize

productivity based on the order of importance were nutrient retention, root media, land preparation, and available nutrients. This was also based on the results of multiple

Study of Land Quality and Land Characteristics that Determine the Productivity of

Composite Maize Varieties in Gorontalo

regression tests with the best equation (equation 1) of the land quality that affects composite maize production were:

 $Y = 5.892 + 0.430X_1 + 0.453X_2 + 0.248X_3 - 0.443X_4$

r = 0.56

Where: X_1 = root medium, X_2 = nutrient retention, X_3 = nutrients availability, X_4 = land preparation. Furthermore, the land characteristics that most influenced the productivity of composite maize based on the order of importance were pH KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. This was also based on the results of multiple regression tests with the best equation (equation 2) as follows:

 $Y = 4.531 + 0.450X_1 - 0.351X_2 - 0.365X_3 + 0.321X_4 - 0.352X_5 + 0.351X_6 + 0.337X_1...(2)$

Where: $X_1 = pH$ of KCl, $X_2 = coarse$ material, $X_3 = rock$ outcrop, $X_4 = effective$ depth, $X_5 = surface$ rock, $X_6 = K$ available, $X_7 = soil$ texture.

The relationship of each land characteristic and its contribution to land quality in influencing the composite maize productivity was presented in Table 5 and Figure 3. The land characteristics consisting of texture, effective depth, pH of KCI, and available K has a positive relationship and significant effect on the composite maize productivity. This shows that the increasing of the land characteristics by 1%, the composite maize productivity will increase by 30% to 47%. In contrast, the content of coarse material, surface rock, and rock outcrops has a negative relationship and significant effect on the composite maize productivity. This indicates that the decreasing content of coarse material, surface rock, and rock outcrops was 1% each in line with the increase in the composite maize productivity by 42% to 44%. The correlation of each of these land characteristics was quite strong in influencing the composite maize productivity. Coarse material is rock fragments measuring 2 mm in diameter or more which affect soil moisture, infiltration, erosion, and land use [20]. Coarse material <15% is very suitable for maize, while > 55% is not suitable [21] [22] [23]. The most suitable soil texture for maize is a fine or loamy texture [24]. Meanwhile, the deeper effective depth affects root growth and development, so that plants can grow and develop well [25]. Surface rocks and rock outcrops are limiting factors in the suitability of maize land in Saentis Village [26].

Table 5. Coefficient of correlation and contribution level on land quality of the land characteristics and composite maize
productivity

Coefficient of correlation	Contribution on land quality (%)	Coefficient of correlation
Temperature (X _{1.1})	0.127	0.20
Rainfall (X _{2.1})	0.279	17.2
Wet months (X _{2.2})	0.209	13.7
Dry months (X _{2.3})	-0.124	-13.2
LGP (X _{2.4})	0.166	12.2
Drainage (X _{3.1})	0.084	14.1
Texture (X _{4.1})	0.298*	18.4
Coarse material (X _{4.2})	-0.438**	-89.4
Effective depth (X _{4.3})	0.431**	76.1
рН H ₂ O (Х _{5.1})	0.254	32.0
pH KCl (X _{5.2})	0.471**	43.2
C-Organic (X _{5.3})	0.264	41.5
CEC (X _{5.4})	0.123	24.7
Base saturation $(X_{5.5})$	0.216	47.3
N Total (X _{6.1})	0.158	46.7
Available P (X _{6.2})	0.012	33.2
Available K (X _{6.3})	0.368*	77.5
ESP (X _{7.1})	-0.024	-17.1
Slope (X _{8.1})	-0.266	-44.4
Soil erosion (X _{8.2})	-0.158	-28.3
Inundation height (X _{9.1})	0.014	23.1
Inundation period (X _{9.2})	0.010	20.1
Surface rock (X _{10.1})	-0.418**	-83.7
Rock outcrop (X _{10.2})	-0.436**	-85.0

*Significant on level test of 5%; ** Significant on level test of 1%.

CONCLUSION

Land quality that controls the productivity of composite maize based on the order of importance is nutrient retention, root media, land preparation, and available nutrients. Meanwhile, land characteristics that control the productivity of composite maize based on the order of importance are pH of KCI, coarse material, rock outcrop, effective depth, surface rock, available K, and soil texture. Soil texture, effective depth, pH of KCI, and available K had a positive and significant effect on the productivity of composite maize, while the content of coarse material, surface rock, and rock outcrops had a negative relationship and had a significant effect on the productivity of composite maize.

ACKNOWLEDGMENT

Composite Maize Varieties in Gorontalo

Thank you to the Brawijaya University, which provided PNBP research funding for the 2019 budget year. Thanks also went to Ecan Adam, SE, MM and Rival Rahman, SP, MSi for helping with SEM-PLS analysis and mapping of land resources.

REFERENCES

- 1. BPS RI. 2019. Maize production by province (tons), 1993-2018. Central Bureau of Statistics of the Republic of Indonesia, Jakarta.
- Yasin, H. G. M., W. Langgo, dan Faesal. 2014. White seed maize as an alternative staple food. J. Food Crops Science and Technology 9(2): 108 - 117.
- BPS Kabupaten Gorontalo. 2019. Gorontalo regency in 2019 figures. Gorontalo Regencyt Statistics Agency, Limboto.
- Zakaria, A. K. 2011. Anticipatory policy and farmers consolidating strategy toward national maize selfsufficiency. J. Agricultural Policy Analysis 9(3): 261-274.
- Bahtiar and B. Kumontoi. 2015. Challenges of composite maize seed production in Bolaang Mongondow regency, North Sulawesi. Proceedings of the National Cereals Seminar 2015: 596 - 604.
- 6. Murni, A. M and R. W. Arief. 2008. Maize cultivation technology. Center for Agricultural Studies and Development, Bogor.
- Swastika, D. K. S. 2002. Maize self-sufficiency in Indonesia: The past 30 years and future prospect. J. Indonesian Agricultural Research and Development 21(3): 75-83.
- 8. Subardja, D and Sudarsono. 2005. Effect of land quality on maize productivity in volcanic soils and sedimentary rocks in the Bogor area. J. Soils and Climate **23**: 38-47.
- 9. FAO. 1976. A Framework for land evaluation. Food and Agriculture Organization Soil Bull. No. 32, Rome.
- 10. Subardja, D. 2005. Land suitability criteria for land use types based on maize and peanuts in the Bogor area. Dissertation. IPB Postgraduate School, Bogor.
- 11. Syaf, H. 2014. Evaluation of the relationship between land quality, growth and production of aged cocoa in East Kolaka Regency, Southeast Sulawesi Province. J. Bioeducation **3**(1): 267 276.
- Elisanti A. D, W. Purnomo and S. Melaniani. 2013. Implementation of partial least square health status of children under five in Indonesia. J. Biometrics and Population 2(2): 99 – 107.
- Ulum, M., I. M. Tirta, and D. Anggraeni. 2014. Analysis of structural equation modeling (SEM) for small samples using the partial least square (PLS) approach. Proceedings of the National Mathematics Seminar, University of Jember, 19 November 2014: 1 – 15.
- 14. Jaya, I. G. N. M and I. M. Sumertajaya. 2008. Structural equation modeling with partial least square. Proceedings of the National Seminar on Mathematics and Mathematics Education 2008: 119 132.
- 15. Hair, J. F., Ringle, C. M. and M. Sarstedt. 2013. Partial least squares structural equation modeling: Rigorous applications, better results and higher acceptance. J. Long Range Planning **46**: 1 12.
- Igbaria, M., N. Zinatelli, P. Cragg, and A. L. M. Cavaye. 1997. Personal computing acceptance factors in small firms: a structural equation model. J. MIS Quarterly 21(3): 279 – 305.
- 17. Mattjik, A. A and I. M. Sumertajaya. 2011. Investigate multiple variables using SAS. IPB Press, Bogor.

- 18. Sujarweni, V. W. 2014. SPSS for research. Pustaka Baru, Yogyakarta.
- 19. Abdilah, W., and J. Hartono. 2015. Partial least square (PLS): alternative structural equation modeling (SEM) in business research. Andi Offset, Yogyakarta.
- 20. Soil Research Institute. 2004. Technical guidelines for soil observation. Research and Development Center for Soil and Agro-climate, Bogor.
- 21. Djaenudin D, Marwan H, Subagjo H, and A. Hidayat. 2011. Technical guidelines for land evaluation for agricultural commodities. Center for R&D for agricultural land resources, Agricultural R&D Agency, Bogor. 36p.
- 22. Ritung S, K Nugroho, A Mulyani, and E Suryani. 2012. Technical guidelines for land evaluation for agricultural commodities. Center for Research and Development of Land Resources, Agricultural Research and Development Agency, Bogor. 166 p.
- Wahyunto, Hikmatullah, E. Suryani, C. Tafakresnanto, S. Ritung, A. Mulyani, Sukarman, K. Nugroho, Y. Sulaeman, Y. Apriyana, Suciantini, A. Pramudia, Suparto, RE Subandiono, T. Sutriadi, D. Nursyamsi. 2016. Technical guidelines for land suitability assessment guidelines for semi-detailed strategic agricultural commodities at a scale of 1: 50,000. Center for Research and Development of Agricultural Land Resources, Agricultural Research and Development Agency, Bogor. 37 p.
- 24. Sudjana A, A. Rifin and M. Sudjadi. 1991. Maize. Agricultural research and development agency, Bogor.
- 25. Wirosoedarmo R, A. T. Sutanhaji, E. Kurniati and R. Wijayanti. 2011. Evaluation of land suitability for maize using spatial analysis methods. J. Agritech **31**(1): 71-78.
- 26. Elfayetti and Herdi. 2015. Evaluation of land suitability for maize crops in Saentis Village, Percut Sei Tuan. J Social Sciences Education **7**(1): 33-40.

Figure 1. Map of the Research Location

Study of Land Quality and Land Characteristics that Determine the Productivity of Composite Maize Varieties in Gorontalo

Figure 3. Path coefficient diagram of land quality to productivity level of composite maize